

## On a Conjecture of Phadke and Thakare

Jochen Brüning  
*Fachbereich Mathematik der Universität München*  
*Theresienstrasse 39*  
*8000 München 2*  
*Federal Republic of Germany*

and

Ki Hang Kim and Fred N. Roush  
*Mathematics Research Group*  
*Alabama State University*  
*Montgomery, Alabama 36101*

Submitted by Richard A. Brualdi

---

### ABSTRACT

We prove the connectedness of the set of all nonzero bounded linear operators on a complex Hilbert space having a generalized inverse.

---

In a recent paper [3] S. V. Phadke and N. K. Thakare conjectured that in a complex Hilbert space  $H$  the set of operators having a generalized inverse is not connected. The purpose of this note is to disprove this conjecture. We recall that a bounded linear operator  $A \neq 0$  on  $H$  is said to have a generalized inverse if there is a bounded linear operator  $B$  on  $H$  such that

$$ABA = A. \quad (1)$$

As usual we write  $|A| := (A^*A)^{1/2}$  and denote by  $s(|A|)$  the support of  $|A|$ . Then (1) is easily seen to be equivalent to the following condition: there is  $C > 0$  such that

$$A^*A \geq Cs(|A|). \quad (2)$$

The set of all operators with generalized inverse will be denoted by  $GI(H)$ .

THEOREM.  $GI(H)$  is pathwise connected.

*Proof.* Let  $A \neq 0$  be a bounded linear operator on  $H$  with generalized inverse, and let  $U|A|=A$  be the polar decomposition of  $A$ . Then

$$t \mapsto U((1-t)|A| + ts(|A|)), \quad t \in [0, 1],$$

is a path in  $GI(H)$  in view of (2), connecting  $A$  and  $U$ . The operators  $P := UU^*$  and  $Q := U^*U$  are orthogonal projections on  $H$ , and we may assume that  $\dim(1_H - P)(H) \leq \dim(1_H - Q)(H)$ . Now if  $P$  is finite, then these dimensions are equal. Consequently, there exists a partial isometry  $V$  on  $H$  with  $VV^* = 1_H - P$ ,  $V^*V = 1_H - Q$ . But then  $U + V$  is unitary and can be connected with  $U$  through a path in  $GI(H)$ , namely

$$t \mapsto U + tV, \quad t \in [0, 1].$$

Next we assume that  $P$  is infinite. Then we can find a partial isometry  $V$  on  $H$  with  $VV^* = 1_H - P$  and  $V^*V \leq 1_H - Q$ . As before,  $U$  can be connected with  $U + V$  in  $GI(H)$ , so we may assume  $P = 1_H$  from now on. We pick projections  $P_1, P_2$  on  $H$  with  $P_1P_2 = 0$ ,  $P_1 + P_2 = 1_H$ , and  $\dim P_1(H) = \dim P_2(H) = \dim H$ . Then the operators  $Q_i := U^*P_iU$ ,  $i = 1, 2$ , are orthogonal projections, too, satisfying  $Q_1Q_2 = 0$ ,  $Q_1 + Q_2 = Q$ , and  $\dim Q_i(H) = \dim P_i(H) = \dim H$ ,  $i = 1, 2$ . But then also  $\dim(1_H - Q_1)(H) = \dim H$ , implying that there is a partial isometry  $W$  on  $H$  with  $WW^* = P_2$  and  $W^*W = 1_H - Q_1$ . We now define

$$U(t) := \begin{cases} UQ_1 + (1-t)UQ_2, & t \in [0, 1], \\ UQ_1 + (t-1)W, & t \in [1, 2]. \end{cases}$$

Then  $U(0) = U$ , and  $U(2)$  is again unitary. Moreover, using (2), it follows that  $U(t) \in GI(H)$  for  $t \in [0, 2]$ . Since the set of all invertible bounded linear operators on  $H$  is connected [2, p. 70],  $U$  can be connected with  $1_H$  and the theorem is proved. ■

We remark that (1) makes sense in an arbitrary  $W^*$ -algebra. The above statement holds also in this more general case; the details of the proof can be found in [1].

#### REFERENCES

- 1 J. Brüning, Über Windungszahlen in endlichen  $W^*$ -Algebren und verwandte Fragen, Habilitationsschrift, Marburg, 1977.
- 2 P. R. Halmos, *A Hilbert Space Problem Book*. Princeton, Van Nostrand, 1967.
- 3 S. V. Phadke and N. K. Thakare, Generalized inverses and operator equations, *Linear Algebra and Appl.* 23:191-199 (1979).

Received 13 June 1979