

Lectures on Symplectic Field Theory

Chris Wendl

INSTITUT FÜR MATHEMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, UNTER
DEN LINDEN 6, 10099 BERLIN, GERMANY

Email address: wendl@math.hu-berlin.de

Contents

Preface	vii
About the current version	xi
Chapter 1. Introduction	1
1.1. In the beginning, Gromov wrote a paper	1
1.2. Hamiltonian Floer homology	4
1.3. Contact manifolds and the Weinstein conjecture	10
1.4. Symplectic cobordisms and their completions	16
1.5. Contact homology and SFT	20
1.6. Two applications	23
Chapter 2. Basics on holomorphic curves	27
2.1. Linearized Cauchy-Riemann operators	27
2.2. Some useful Sobolev inequalities	27
2.3. The fundamental elliptic estimate	27
2.4. Regularity	27
2.5. Linear local existence and the similarity principle	27
2.6. Simple curves and multiple covers	27
2.7. Nonlinear local existence	27
2.8. The nonlinear equation on push-offs	27
Chapter 3. Asymptotic operators	29
3.1. Stable Hamiltonian structures and Reeb orbits	29
3.2. The linearization in Morse homology	29
3.3. The Hessian of the contact action functional	29
3.4. Spectral flow	29
3.5. The Conley-Zehnder index of a nondegenerate orbit	30
3.6. Winding numbers of eigenfunctions	30
3.7. Elliptic and hyperbolic orbits	30
3.8. CZ = Morse for geodesics	30
3.9. Morse-Bott families	30
Chapter 4. Fredholm theory with cylindrical ends	31
4.1. Cauchy-Riemann operators with punctures	31
4.2. A lemma on semi-Fredholm operators	31
4.3. Some global regularity estimates	31
4.4. Translation-invariant operators on the cylinder	31
4.5. Proof of the semi-Fredholm property	31

4.6.	Exponential decay	31
4.7.	Formal adjoints and proof of the Fredholm property	31
4.8.	The asymptotic formula	31
Chapter 5. The index formula		33
5.1.	Riemann-Roch with punctures	33
5.2.	Some remarks on the formal adjoint	33
5.3.	The index zero case on a torus	33
5.4.	A Weitzenböck formula for Cauchy-Riemann operators	33
5.5.	Large antilinear perturbations and energy concentration	33
5.6.	Two Cauchy-Riemann type problems on the plane	33
5.7.	A linear gluing argument	33
5.8.	Antilinear deformations of asymptotic operators	33
Chapter 6. Symplectic cobordisms and moduli spaces		35
6.1.	Stable Hamiltonian structures	35
6.2.	Almost complex manifolds with cylindrical ends	35
6.3.	Examples of stable Hamiltonian structures	35
6.4.	Moduli spaces of asymptotically cylindrical curves	36
6.5.	Asymptotic regularity	36
6.6.	Simple curves and multiple covers revisited	36
6.7.	Possible generalizations	36
Chapter 7. Asymptotics and compactness		37
7.1.	Removal of singularities	37
7.2.	Finite energy and asymptotics	37
7.3.	Degenerations of holomorphic curves	37
7.4.	The SFT compactness theorem	37
7.5.	Compactness results for biholomorphic maps	37
Chapter 8. Smoothness of the moduli space		39
8.1.	The main result on regular curves	39
8.2.	Functional-analytic setup	39
8.3.	Moduli of complex structures	39
8.4.	Fredholm regularity and the implicit function theorem	39
8.5.	Evaluation and forgetful maps	39
Chapter 9. Transversality		41
9.1.	A paradigm for genericity arguments	41
9.2.	Generic transversality in cobordisms	41
9.3.	Generic transversality in symplectizations	41
9.4.	Transversality of constraints	41
Chapter 10. Gluing		43
10.1.	Moduli spaces of holomorphic buildings	43
10.2.	Deligne-Mumford space is an orbifold	43
Chapter 11. Cylindrical contact homology and the tight 3-tori		45

11.1.	Contact structures on \mathbb{T}^3 and Giroux torsion	45
11.2.	Definition of cylindrical contact homology	45
11.3.	Computing $HC_*(\mathbb{T}^3, \xi_k)$	45
Chapter 12.	Coherent orientations	47
12.1.	Gluing maps and coherence	47
12.2.	Permutations of punctures and bad orbits	47
12.3.	Orienting moduli spaces in general	47
12.4.	The determinant line bundle	47
12.5.	Determinant bundles of moduli spaces	47
12.6.	An algorithm for coherent orientations	47
12.7.	Permutations and bad orbits revisited	47
Chapter 13.	The generating function of SFT	49
13.1.	Some important caveats on transversality	49
13.2.	Auxiliary data, grading and supercommutativity	49
13.3.	The definition of \mathbf{H} and commutators	49
13.4.	Interlude: Orbifolds and branched manifolds	49
13.5.	Cylindrical contact homology revisited	49
13.6.	Combinatorics of gluing	49
13.7.	Some remarks on torsion, coefficients, and conventions	49
Chapter 14.	Contact invariants	51
14.1.	The Eliashberg-Givental-Hofer package	51
14.2.	SFT generating functions for cobordisms	51
14.3.	Full SFT as a BV_∞ -algebra	51
Chapter 15.	Transversality and counting singularities in dimension four	53
15.1.	Automatic transversality	53
15.2.	Curves in symplectizations of 3-manifolds	53
15.3.	Implicit function theorems for local foliations	53
15.4.	Consequences for coherent orientations	53
Chapter 16.	Intersection theory for punctured holomorphic curves	55
16.1.	Prologue	55
16.2.	Homotopy-invariant intersection numbers	55
16.3.	The adjunction formula	55
16.4.	Local foliations: the general case	55
Chapter 17.	Torsion computations and applications	57
17.1.	Some J -holomorphic foliations	57
17.2.	Contact homology of overtwisted contact manifolds	57
17.3.	Examples with higher-order algebraic torsion	57
17.4.	Rigorous obstructions to fillings and cobordisms	57
Appendix A.	Sobolev spaces	59
A.1.	Approximation, extension and embedding theorems	59
A.2.	Products, compositions, and rescaling	64

A.3. Difference quotients	71
A.4. Spaces of sections of vector bundles	75
A.5. Some remarks on domains with cylindrical ends	79
Appendix B. The Floer C_ϵ space	83
Appendix C. Genericity in the space of asymptotic operators	87
Bibliography	93

Preface

This book is an expanded version of the lecture notes I produced for a two-semester course taught at University College London in 2015–16, for Ph.D. students with a background in basic symplectic geometry and interest in symplectic topology and/or geometric analysis. For the most part, each chapter corresponds to a two-hour lecture in the original course, though the reader will quickly notice that in this “expanded” version, most individual chapters contain far more material than can reasonably fit into one lecture (or even two). In reality, much of that material was only sketched or mentioned in passing during lectures, and I ended up using the notes to discuss everything that I would like to have explained if I’d had unlimited time. This includes relatively detailed discussions of several important technical points (e.g. the definition of spectral flow, generic transversality in symplectizations, the punctured Riemann-Roch formula, finite energy and asymptotics with arbitrary stable Hamiltonian structures) which are either incompletely covered by the existing literature or, in my opinion, simply more difficult to learn from other sources than they should be. For topics that are, on the other hand, well covered elsewhere, I have usually not felt obliged to explain every detail, but have tried always to provide adequate references.

One of the interesting features of SFT is that its foundations are—at the time of this writing—not yet complete. When the original “propaganda paper” [EGH00] appeared in 2000, it was widely believed that the technical details would be filled in within a few years, and several papers introducing important applications of SFT to contact topology were written under this assumption. Since then, a certain realization has set in that the results in those papers cannot truly be regarded as “theorems” in the sense of mathematics, and it has become less socially acceptable to preface statements of results with caveats of the form, “this theorem is dependent on the foundations of SFT”. At the same time, the need for a robust perturbation scheme to achieve transversality in SFT spawned the development of a whole new approach to infinite-dimensional differential geometry, the *polyfold* project [Hof06], which is intended for much more general applications but is not yet finished. Opinions vary among symplectic topologists as to how unsatisfied we should all be with this state of affairs, and what could be done about it—among other things, one could make an entire course out of the discussion of such issues, but I have not chosen to do that. My approach is instead to develop the *classical*¹ analysis of pseudoholomorphic curves in symplectizations and symplectic cobordisms, to explain how this would lead to a theory of algebraic contact invariants if transversality for multiple

¹For the purposes of this discussion, the word “classical” may be defined as “not involving the words *polyfold*, *virtual* or *Kuranishi*”.

covers were not an issue, and then to use the tools and insights gained from this discussion to prove *rigorous mathematical theorems* about contact manifolds. Typically, such theorems can be regarded informally as consequences of computations in a (not yet well-defined) theory called SFT, but in a rigorous sense, they are actually consequences of the methods used in those computations. Examples covered in these notes include distinguishing tight contact structures on the 3-torus that are homotopic but not isomorphic (Chapter 11), and the nonexistence of symplectic fillings or symplectic cobordisms between certain pairs of contact manifolds (Chapter 17). The choice of applications is of course biased somewhat toward my own research interests.

Prerequisites. The stated target audience for the original lecture course was “advanced Masters and Ph.D. students in differential geometry or related fields who are not afraid of analysis”. More precisely, the notes assume some knowledge of the following topics:

- Differential geometry: manifolds and vector bundles, differential forms and Stokes’ theorem, connections, basic familiarity with symplectic manifolds;
- Functional analysis: linear operators on Banach spaces, basics of Sobolev spaces, Fredholm operators;
- Differential topology: smooth mapping degree, intersection numbers, Sard’s theorem;
- Algebraic topology: fundamental group, homology and cohomology of manifolds, Poincaré duality, first Chern class, homological intersection numbers.

The following topics are not considered formal prerequisites, but some knowledge of them is likely in any case to be helpful to the reader, who may want to have a good reference for them (as suggested below) within arm’s reach:

- Contact manifolds (e.g. Geiges [Gei08]);
- Differential calculus on Banach spaces and Banach manifolds (e.g. these two books by Lang: [Lan93] and [Lan99]);
- Closed pseudoholomorphic curves (e.g. McDuff-Salamon [MS12] or my other book in preparation [Wenb]);
- Floer homology (e.g. Salamon [Sal99] or Audin-Damian [AD14]).

Acknowledgements. I would like to thank the students who have sat through various iterations of the course that gave rise to this book, notably Alexandru Cioba and Agustín Moreno for their assistance in editing the first several lectures, as well as Adrian Dawid, Milica Đukić, Shah Faisal, Solveig Hepp, Catalina Jurja, and Michael Rothgang for useful comments. My understanding of Taubes’s approach to the Riemann-Roch formula (explained in Chapter 5) and its generalization to the punctured case emerged in part from discussions with Chris Gerig, and I am grateful also to Tim Perutz for helpful hints about Weitzenböck formulas, and Patrick Massot for patient discussions of singular integral operators and elliptic regularity. Thanks also to Michael Hutchings and Janko Latschev for helping me understand the combinatorial factors in Chapter 13, to Jo Nelson for helpful comments on coefficients and orbifold singularities, and to Sam Lisi and Barney Bramham for advice on the

Floer C_ϵ space. And also to Klaus Niederkrüger and Helmut Hofer for enlightening discussions on all manner of things.

About the current version

The version you see in front of you is being revised and updated regularly to accompany a Masters-level special topics course on symplectic field theory at the Humboldt-Universität zu Berlin in the 2026 summer semester.

I have tried to produce a manuscript that is relatively well polished, but I have not tried quite as diligently for that as I do with most of my research papers. As of the beginning of the 2026 summer semester, some of the later chapters that have been in planning for over a decade are not yet complete, and one or two additional chapters exist only as vague plans in my head, so if those chapters exist by the end of the semester, they are unlikely to be error-free. I apologize for any sloppiness that I may have failed so far to expunge. All comments and corrections are welcome,² and may be sent to wendl@math.hu-berlin.de. Updates on the publication of the book will be posted periodically on my website at

<https://www.mathematik.hu-berlin.de/~wendl/publications.html#notes>

Most recent update: **April 15, 2026**

²especially if those corrections are received before the book goes to press

CHAPTER 1

Introduction

Contents

1.1. In the beginning, Gromov wrote a paper	1
1.2. Hamiltonian Floer homology	4
1.3. Contact manifolds and the Weinstein conjecture	10
1.4. Symplectic cobordisms and their completions	16
1.5. Contact homology and SFT	20
1.6. Two applications	23
1.6.1. Tight contact structures on \mathbb{T}^3	23
1.6.2. Filling and cobordism obstructions	24

Symplectic field theory is a general framework for defining invariants of contact manifolds and symplectic cobordisms between them via counts of “asymptotically cylindrical” pseudoholomorphic curves. In this first chapter, we’ll summarize some of the historical background of the subject, and then sketch the basic algebraic formalism of SFT.

1.1. In the beginning, Gromov wrote a paper

Pseudoholomorphic curves first appeared in symplectic geometry in a 1985 paper of Gromov [Gro85]. The development was revolutionary for the field of symplectic topology, but it was not unprecedented: a few years before this, Donaldson had demonstrated the power of using elliptic PDEs in geometric contexts to define invariants of smooth 4-manifolds (see [DK90]). The PDE that Gromov used was a slight generalization of one that was already familiar from complex geometry.

Recall that if M is a smooth $2n$ -dimensional manifold, an **almost complex structure** on M is a smooth linear bundle map $J : TM \rightarrow TM$ such that $J^2 = -\mathbb{1}$. This makes the tangent spaces of M into complex vector spaces and thus induces an orientation on M ; the pair (M, J) is called an **almost complex manifold**. In this context, a **Riemann surface** is an almost complex manifold of real dimension 2 (hence complex dimension 1), and a **pseudoholomorphic curve** (also called **J -holomorphic**) is a smooth map

$$u : \Sigma \rightarrow M$$

satisfying the **nonlinear Cauchy-Riemann equation**

$$(1.1) \quad Tu \circ j = J \circ Tu,$$

where (Σ, j) is a Riemann surface and (M, J) is an almost complex manifold (of arbitrary dimension). The almost complex structure J is called **integrable** if M admits the structure of a complex manifold such that J is multiplication by i in holomorphic coordinate charts. By a basic theorem due to Gauss, every almost complex structure in real dimension two is integrable, hence one can always find local coordinates (s, t) on neighborhoods in Σ such that

$$j\partial_s = \partial_t, \quad j\partial_t = -\partial_s.$$

In these coordinates, (1.1) takes the form

$$\partial_s u + J(u)\partial_t u = 0.$$

The fundamental insight of [Gro85] was that solutions to the equation (1.1) capture information about symplectic structures on M whenever they are related to J in the following way.

DEFINITION 1.1.1. Suppose (M, ω) is a symplectic manifold. An almost complex structure J on M is said to be **tamed** by ω if

$$\omega(X, JX) > 0 \quad \text{for all } X \in TM \text{ with } X \neq 0.$$

Additionally, J is **compatible** with ω if the pairing

$$g(X, Y) := \omega(X, JY)$$

defines a Riemannian metric on M .

EXERCISE 1.1.2. Show that an almost complex structure J is compatible with a symplectic form ω if and only if it is tame and ω is J -invariant.

We shall denote by $\mathcal{J}(M)$ the space of all smooth almost complex structures on M , with the C_{loc}^∞ -topology, and if ω is a symplectic form on M , let

$$\mathcal{J}_\tau(M, \omega), \mathcal{J}(M, \omega) \subset \mathcal{J}(M)$$

denote the subsets consisting of almost complex structures that are tamed by or compatible with ω respectively. Notice that $\mathcal{J}_\tau(M, \omega)$ is an open subset of $\mathcal{J}(M)$, but $\mathcal{J}(M, \omega)$ is not. Proofs of the following may be found in [MS17, §2.5] or [Wenb, §2.2], among other places.

PROPOSITION 1.1.3. *On any symplectic manifold (M, ω) , the spaces $\mathcal{J}_\tau(M, \omega)$ and $\mathcal{J}(M, \omega)$ are each nonempty and contractible.* \square

Tameness implies that the **energy** of a J -holomorphic curve $u : \Sigma \rightarrow M$,

$$E(u) := \int_\Sigma u^* \omega,$$

is always nonnegative, and it is strictly positive unless u is constant. Notice moreover that if the domain Σ is closed, then $E(u)$ depends only on the cohomology class $[\omega] \in H_{\text{dR}}^2(M)$ and the homology class

$$[u] := u_*[\Sigma] \in H_2(M),$$

so in particular, any family of J -holomorphic curves in a fixed homology class satisfies a uniform energy bound. This basic observation is one of the key facts behind

Gromov’s compactness theorem, which states that moduli spaces of closed curves in a fixed homology class are compact up to “nodal” degenerations.

The most famous application of pseudoholomorphic curves presented in [Gro85] is Gromov’s *nonsqueezing theorem*, which was the first known example of an obstruction for embedding symplectic domains that is subtler than the obvious obstruction defined by volume. The technology introduced in [Gro85] also led directly to the development of the *Gromov-Witten invariants* (see [MS12, RT95, RT97]), which follow the same pattern as Donaldson’s earlier smooth 4-manifold invariants: they use counts of J -holomorphic curves to define invariants of symplectic manifolds up to symplectic deformation equivalence.

Here is another sample application from [Gro85]. We denote by

$$A \cdot B \in \mathbb{Z}$$

the intersection number between two homology classes $A, B \in H_2(M)$ in a closed oriented 4-manifold M .

THEOREM 1.1.4. *Suppose (M, ω) is a closed and connected symplectic 4-manifold with the following properties:*

- (i) (M, ω) does not contain any symplectic submanifold $S \subset M$ that is diffeomorphic to S^2 and satisfies $[S] \cdot [S] = -1$.
- (ii) (M, ω) contains two symplectic submanifolds $S_1, S_2 \subset M$ which are both diffeomorphic to S^2 , satisfy

$$[S_1] \cdot [S_1] = [S_2] \cdot [S_2] = 0,$$

and have exactly one intersection point with each other, which is transverse and positive.

Then (M, ω) is symplectomorphic to $(S^2 \times S^2, \sigma_1 \oplus \sigma_2)$, where for $i = 1, 2$, the σ_i are area forms on S^2 satisfying

$$\int_{S^2} \sigma_i = \langle [\omega], [S_i] \rangle.$$

SKETCH OF THE PROOF. Since S_1 and S_2 are both symplectic submanifolds, one can choose a compatible almost complex structure J on M for which both of them are the images of embedded J -holomorphic curves. One then considers the moduli spaces $\mathcal{M}_1(J)$ and $\mathcal{M}_2(J)$ of equivalence classes of J -holomorphic spheres homologous to S_1 and S_2 respectively, where any two such curves are considered equivalent if one is a reparametrization of the other (in the present setting this just means they have the same image). These spaces are both manifestly nonempty, and one can argue via Gromov’s compactness theorem for J -holomorphic curves that both are compact. Moreover, an infinite-dimensional version of the implicit function theorem implies that both are smooth 2-dimensional manifolds, carrying canonical orientations, hence both are diffeomorphic to closed surfaces. Finally, one uses *positivity of intersections* to show that every curve in $\mathcal{M}_1(J)$ intersects every curve in $\mathcal{M}_2(J)$ exactly once, and this intersection is always transverse and positive; moreover, any two curves in the same space $\mathcal{M}_1(J)$ or $\mathcal{M}_2(J)$ are either identical or disjoint. It follows that both moduli spaces are diffeomorphic to S^2 , and both

consist of smooth families of J -holomorphic spheres that foliate M , hence defining a diffeomorphism

$$\mathcal{M}_1(J) \times \mathcal{M}_2(J) \rightarrow M$$

that sends (u_1, u_2) to the unique point in the intersection $\text{im } u_1 \cap \text{im } u_2$. This identifies M with $S^2 \times S^2$ such that each of the submanifolds $S^2 \times \{*\}$ and $\{*\} \times S^2$ are symplectic. The latter observation can be used to determine the symplectic form up to deformation, so that by the Moser stability theorem, ω is determined up to isotopy by its cohomology class $[\omega] \in H_{\text{dR}}^2(S^2 \times S^2)$, which depends only on the evaluation of ω on $[S^2 \times \{*\}]$ and $[\{*\} \times S^2] \in H_2(S^2 \times S^2)$. \square

For a detailed exposition of the above proof of Theorem 1.1.4, see [Wen18, Theorem E].

1.2. Hamiltonian Floer homology

Throughout the following, we write

$$S^1 := \mathbb{R}/\mathbb{Z},$$

so maps on S^1 are the same as 1-periodic maps on \mathbb{R} . One popular version of the *Arnold conjecture* on symplectic fixed points can be stated as follows. Suppose (M, ω) is a closed symplectic manifold and $H : S^1 \times M \rightarrow \mathbb{R}$ is a smooth function. Writing $H_t := H(t, \cdot) : M \rightarrow \mathbb{R}$, H determines a 1-periodic time-dependent Hamiltonian vector field X_t via the relation¹

$$(1.2) \quad \omega(X_t, \cdot) = -dH_t.$$

CONJECTURE 1.2.1 (Arnold conjecture). *If all 1-periodic orbits of X_t are nondegenerate, then the number of these orbits is at least the sum of the Betti numbers of M .*

Here a 1-periodic orbit $\gamma : S^1 \rightarrow M$ of X_t is called **nondegenerate** if, denoting the flow of X_t by φ^t , the linearized time 1 flow

$$d\varphi^1(\gamma(0)) : T_{\gamma(0)}M \rightarrow T_{\gamma(0)}M$$

does not have 1 as an eigenvalue. This can be thought of as a Morse condition for an action functional on the loop space whose critical points are periodic orbits; like Morse critical points, nondegenerate periodic orbits occur in isolation. To simplify our lives, let's restrict attention to *contractible* orbits and also assume that (M, ω) is **symplectically aspherical**, which means

$$[\omega]|_{\pi_2(M)} = 0, \quad \text{i.e.} \quad \langle [\omega], [u] \rangle = 0 \text{ for all continuous maps } u : S^2 \rightarrow M.$$

Then if $C_{\text{contr}}^\infty(S^1, M)$ denotes the space of all smoothly contractible smooth loops in M , the **symplectic action functional** can be defined by

$$\mathcal{A}_H : C_{\text{contr}}^\infty(S^1, M) \rightarrow \mathbb{R} : \gamma \mapsto - \int_{\mathbb{D}} \bar{\gamma}^* \omega + \int_{S^1} H_t(\gamma(t)) dt,$$

¹Elsewhere in the literature, you will sometimes see (1.2) without the minus sign on the right hand side. If you want to know why I strongly believe that the minus sign belongs there, see [Wen18], but to some extent this is just a personal opinion.

where $\bar{\gamma} : \mathbb{D} \rightarrow M$ is any smooth map on the closed unit disk $\mathbb{D} \subset \mathbb{C}$ satisfying

$$\bar{\gamma}(e^{2\pi it}) = \gamma(t),$$

and the symplectic asphericity condition guarantees that $\mathcal{A}_H(\gamma)$ does not depend on the choice of $\bar{\gamma}$.

EXERCISE 1.2.2. The **first variation** of a functional such as $\mathcal{A}_H : C_{\text{contr}}^\infty(S^1, M)$ at $\gamma \in C_{\text{contr}}^\infty(S^1, M)$ is by definition the unique linear map $d\mathcal{A}_H(\gamma) : \Gamma(\gamma^*TM) \rightarrow \mathbb{R}$ such that for any smooth 1-parameter family $\{\gamma_s \in C_{\text{contr}}^\infty(S^1, M)\}_{s \in (-\epsilon, \epsilon)}$ with $\gamma_0 = \gamma$ and $\partial_s \gamma_s|_{s=0} = \eta$, one has

$$\left. \frac{d}{ds} \mathcal{A}_H(\gamma_s) \right|_{s=0} = d\mathcal{A}_H(\gamma)\eta.$$

In other words, if we think of $C_{\text{contr}}^\infty(S^1, M)$ as an infinite-dimensional manifold² with tangent spaces $T_\gamma C_{\text{contr}}^\infty(S^1, M) = \Gamma(\gamma^*TM)$, then $d\mathcal{A}_H(\gamma)$ is simply the differential of \mathcal{A}_H at γ . Prove the formula

$$d\mathcal{A}_H(\gamma)\eta = \int_{S^1} [\omega(\dot{\gamma}, \eta) + dH_t(\eta)] dt = \int_{S^1} \omega(\dot{\gamma} - X_t(\gamma), \eta) dt.$$

Using the nondegeneracy of ω , this shows that the critical points of \mathcal{A}_H are precisely the contractible 1-periodic orbits of X_t .

A few years after Gromov's introduction of pseudoholomorphic curves, Floer proved the most important cases of the Arnol'd conjecture by developing a novel version of infinite-dimensional Morse theory for the functional \mathcal{A}_H . This approach mimicked the homological approach to Morse theory which has since been popularized in books such as [AD14, Sch93], but was apparently only known to experts at the time. In *Morse homology*, one considers a smooth Riemannian manifold (M, g) with a Morse function $f : M \rightarrow \mathbb{R}$, and defines a chain complex whose generators are the critical points of f , graded according to their Morse index. If we denote the generator corresponding to a given critical point $x \in \text{Crit}(f)$ by $\langle x \rangle$, the boundary map on this complex is defined by

$$\partial \langle x \rangle = \sum_{\text{Morse}(y) = \text{Morse}(x) - 1} \#(\mathcal{M}(x, y)/\mathbb{R}) \langle y \rangle,$$

where $\mathcal{M}(x, y)$ denotes the moduli space of negative gradient flow lines $u : \mathbb{R} \rightarrow M$, satisfying $\partial_s u = -\nabla f(u(s))$, $\lim_{s \rightarrow -\infty} u(s) = x$ and $\lim_{s \rightarrow +\infty} u(s) = y$. This space admits a natural \mathbb{R} -action by shifting the variable in the domain, and one can show that for generic choices of f and the metric g , $\mathcal{M}(x, y)/\mathbb{R}$ is a finite set whenever $\text{Morse}(x) - \text{Morse}(y) = 1$. The real magic, however, is contained in the following statement about the case $\text{Morse}(x) - \text{Morse}(y) = 2$:

²At this stage, it is best not to worry so much over exactly what kind of infinite-dimensional manifold $C_{\text{contr}}^\infty(S^1, M)$ is. It is locally modeled on open subsets of the space of smooth sections $\Gamma(\gamma^*TM)$, which is a Fréchet space, but the notion of a “Fréchet manifold” is not so straightforward to define precisely, and one can easily define the term “first variation” without worrying about it.




FIGURE 1.1. One-parameter families of gradient flow lines on a Riemannian manifold degenerate to broken flow lines.

PROPOSITION 1.2.3. *For generic choices of f and g and any two critical points $x, y \in \text{Crit}(f)$ with $\text{Morse}(x) - \text{Morse}(y) = 2$, $\mathcal{M}(x, y)/\mathbb{R}$ is homeomorphic to a finite collection of circles and open intervals whose end points are canonically identified with the finite set*

$$\partial\overline{\mathcal{M}}(x, y) := \bigcup_{\text{Morse}(z)=\text{Morse}(x)-1} \mathcal{M}(x, z) \times \mathcal{M}(z, y).$$

We say that $\mathcal{M}(x, y)$ has a natural **compactification** $\overline{\mathcal{M}}(x, y)$, which has the topology of a compact 1-manifold with boundary, and its boundary is the set of all **broken flow lines** from x to y , cf. Figure 1.1. This set of broken flow lines is precisely what is counted if one computes the $\langle y \rangle$ coefficient of $\partial^2 \langle x \rangle$, hence we deduce

$$\partial^2 = 0$$

as a consequence of the fact that compact 1-manifolds always have zero boundary points when counted with appropriate signs.³ The homology of the resulting chain complex can be denoted by $HM_*(M; g, f)$ and is called the **Morse homology** of M . The well-known Morse inequalities can then be deduced from a fundamental theorem stating that $HM_*(M; g, f)$ is, for generic f and g , isomorphic to the singular homology of M .

With the above notion of Morse homology understood, Floer's approach to the Arnol'd conjecture can now be summarized as follows:

Step 1: Under suitable technical assumptions, construct a homology theory

$$HF_*(M, \omega; H, \{J_t\}),$$

depending *a priori* on the choices of a Hamiltonian $H : S^1 \times M \rightarrow \mathbb{R}$ with all 1-periodic orbits nondegenerate, and a generic S^1 -parametrized family of ω -compatible almost complex structures $\{J_t\}_{t \in S^1}$. The generators of the

³Counting with signs presumes that we have chosen suitable orientations for the moduli spaces $\mathcal{M}(x, y)$, and this can always be done. Alternatively, one can avoid this issue by counting modulo 2, and thus define a homology theory with \mathbb{Z}_2 coefficients.

chain complex are the critical points of the symplectic action functional \mathcal{A}_H , i.e. 1-periodic orbits of the Hamiltonian flow, and the boundary map is defined by counting a suitable notion of gradient flow lines connecting pairs of orbits (more on this below).

Step 2: Prove that $HF_*(M, \omega) := HF_*(M, \omega; H, \{J_t\})$ is a *symplectic invariant*, i.e. it depends on ω , but not on the auxiliary choices H and $\{J_t\}$.

Step 3: Show that if H and $\{J_t\}$ are chosen to be time-independent and H is also C^2 -small, then the chain complex for $HF_*(M, \omega; H, \{J_t\})$ is isomorphic (with a suitable grading shift) to the chain complex for Morse homology $HM_*(M; g, H)$ with $g := \omega(\cdot, J_t \cdot)$. The isomorphism between $HM_*(M; g, H)$ and singular homology thus implies that the Floer complex must have at least as many generators (i.e. periodic orbits) as there are generators of $H_*(M)$, proving the Arnol'd conjecture.

The implementation of Floer's idea required a different type of analysis than what is needed for Morse homology. The moduli space $\mathcal{M}(x, y)$ in Morse homology is simple to understand as the (generically transverse) intersection between the unstable manifold of x and the stable manifold of y with respect to the negative gradient flow. Conveniently, both of those are finite-dimensional manifolds, with their dimensions determined by the Morse indices of x and y . We will see in Chapter 3 that no such thing is true for the symplectic action functional: to the extent that \mathcal{A}_H can be thought of as a Morse function on an infinite-dimensional manifold, its Morse index and its Morse "co-index" at every critical point are both infinite, hence the stable and unstable manifolds are not nearly as nice as finite-dimensional manifolds, providing no reason to expect that their intersection should be. There are additional problems since $C_{\text{contr}}^\infty(S^1, M)$ does not have a Banach space topology: in order to view the negative gradient flow of \mathcal{A}_H as an ODE and make use of the usual local existence/uniqueness theorems (as in [Lan99, Chapter IV]), one would have to extend \mathcal{A}_H to a smooth function on a suitable Hilbert manifold with a Riemannian metric. There is a very limited range of situations in which one can do this and obtain a reasonable formula for $\nabla \mathcal{A}_H$, e.g. [HZ94, §6.2] explains the case $M = \mathbb{T}^{2n}$, in which \mathcal{A}_H can be defined on the Sobolev space $H^{1/2}(S^1, \mathbb{R}^{2n})$ and then studied using Fourier series. This approach is very dependent on the fact that the torus \mathbb{T}^{2n} is a quotient of \mathbb{R}^{2n} . For general symplectic manifolds (M, ω) , one cannot even define $H^{1/2}(S^1, M)$, since functions of class $H^{1/2}$ on S^1 need not be continuous ($H^{1/2}$ is a "Sobolev borderline case" in dimension one).

One of the novelties in Floer's approach was to refrain from viewing the gradient flow as an ODE in a Banach space setting, but instead to write down a formal version of the gradient flow equation and regard it as an elliptic PDE. To this end, let us regard $C_{\text{contr}}^\infty(S^1, M)$ formally as a manifold with tangent spaces

$$T_\gamma C_{\text{contr}}^\infty(S^1, M) := \Gamma(\gamma^* TM),$$

choose a formal Riemannian metric on this manifold (i.e. a smoothly varying family of L^2 -inner products on the spaces $\Gamma(\gamma^* TM)$) and write down the resulting equation for the negative gradient flow. A suitable Riemannian metric can be defined by




FIGURE 1.2. A family of smooth Floer trajectories can degenerate into a broken Floer trajectory.

choosing a smooth S^1 -parametrized family of compatible almost complex structures

$$\{J_t \in \mathcal{J}(M, \omega)\}_{t \in S^1},$$

abbreviated in the following as $\{J_t\}$, and setting

$$\langle \xi, \eta \rangle_{L^2} := \int_{S^1} \omega(\xi(t), J_t \eta(t)) dt$$

for $\xi, \eta \in \Gamma(\gamma^*TM)$. Exercise 1.2.2 then yields the formula

$$d\mathcal{A}_H(\gamma)\eta = \langle J_t(\dot{\gamma} - X_t(\gamma)), \eta \rangle_{L^2},$$

so that it seems reasonable to define the so-called *unregularized* gradient of \mathcal{A}_H by

$$(1.3) \quad \nabla \mathcal{A}_H(\gamma) := J_t(\dot{\gamma} - X_t(\gamma)) \in \Gamma(\gamma^*TM).$$

Let us also think of a path $u : \mathbb{R} \rightarrow C_{\text{contr}}^\infty(S^1, M)$ as a map $u : \mathbb{R} \times S^1 \rightarrow M$, writing $u(s, t) := u(s)(t)$. The negative gradient flow equation $\partial_s u + \nabla \mathcal{A}_H(u(s)) = 0$ then becomes the elliptic PDE

$$(1.4) \quad \partial_s u + J_t(u) (\partial_t u - X_t(u)) = 0.$$

This is called the **Floer equation**, and its solutions are often called **Floer trajectories**. The relevance of Floer homology to our previous discussion of pseudo-holomorphic curves should now be obvious. Indeed, the resemblance of the Floer equation to the nonlinear Cauchy-Riemann equation is not merely superficial—we will see in Chapter 6 that the former can always be viewed as a special case of the latter. In any case, one can use the same set of analytical techniques for both: elliptic regularity theory implies that Floer trajectories are always smooth, Fredholm theory and the implicit function theorem imply that (under appropriate assumptions) they form smooth finite-dimensional moduli spaces. Most importantly, the same “bubbling off” analysis that underlies Gromov’s compactness theorem can be used to prove that spaces of Floer trajectories are compact up to “breaking”, just as in Morse homology (see Figure 1.2)—this is the main reason for the relation $\partial^2 = 0$ in Floer homology.

We should mention one complication that does not arise either in the study of closed holomorphic curves or in finite-dimensional Morse theory. Since the gradient

flow in Morse homology takes place on a closed manifold, it is obvious that every gradient flow line asymptotically approaches critical points at both $-\infty$ and $+\infty$. The following example shows that in the infinite-dimensional setting of Floer theory, this is no longer true.

EXAMPLE 1.2.4. Consider the Floer equation on $M := S^2 = \mathbb{C} \cup \{\infty\}$ with $H := 0$ and J_t defined as the standard complex structure i for every t . Then the orbits of X_t are all constant, and a map $u : \mathbb{R} \times S^1 \rightarrow S^2$ satisfies the Floer equation if and only if it is holomorphic. Identifying $\mathbb{R} \times S^1$ with $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$ via the biholomorphic map $(s, t) \mapsto e^{2\pi(s+it)}$, a solution u approaches periodic orbits as $s \rightarrow \pm\infty$ if and only if the corresponding holomorphic map $\mathbb{C}^* \rightarrow S^2$ extends continuously (and therefore holomorphically) over 0 and ∞ . But this is not true for every holomorphic map $\mathbb{C}^* \rightarrow S^2$, e.g. take any entire function $\mathbb{C} \rightarrow \mathbb{C}$ that has an essential singularity at ∞ .

EXERCISE 1.2.5. Show that in the above example with an essential singularity at ∞ , the symplectic action $\mathcal{A}_H(u(s, \cdot))$ is unbounded as $s \rightarrow \infty$.

EXERCISE 1.2.6. Suppose $u : \mathbb{R} \times S^1 \rightarrow M$ is a solution to the Floer equation with $\lim_{s \rightarrow \pm\infty} u(s, \cdot) = \gamma_{\pm}$ uniformly for a pair of 1-periodic orbits $\gamma_{\pm} \in \text{Crit}(\mathcal{A}_H)$. Show that

$$(1.5) \quad \mathcal{A}(\gamma_-) - \mathcal{A}(\gamma_+) = \int_{\mathbb{R} \times S^1} \omega(\partial_s u, \partial_t u - X_t(u)) ds dt = \int_{\mathbb{R} \times S^1} \omega(\partial_s u, J_t(u) \partial_s u) ds dt.$$

The right hand side of (1.5) is manifestly nonnegative since J_t is compatible with ω , and it is strictly positive unless $\gamma_- = \gamma_+$. It is therefore sensible to call this expression the **energy** $E(u)$ of a Floer trajectory. The following converse of Exercise 1.2.6 plays a crucial role in the compactness theory for Floer trajectories, as it guarantees that all the “levels” in a broken Floer trajectory are asymptotically well behaved. We will prove a variant of this result in the SFT context (see Prop. 1.3.12 below) in Chapter 7.

PROPOSITION 1.2.7. *If $u : \mathbb{R} \times S^1 \rightarrow M$ is a Floer trajectory with $E(u) < \infty$ and all 1-periodic orbits of X_t are nondegenerate, then there exist orbits $\gamma_-, \gamma_+ \in \text{Crit}(\mathcal{A}_H)$ such that $\lim_{s \rightarrow \pm\infty} u(s, \cdot) = \gamma_{\pm}$ uniformly.*

REMARK 1.2.8. It should be emphasized again that we have assumed $[\omega]|_{\pi_2(M)} = 0$ throughout this discussion. Floer homology can also be defined under more general assumptions, but several details become more complicated.

For nice comprehensive treatments of Hamiltonian Floer homology—unfortunately not always with the same sign conventions as used here—see [Sal99, AD14]. Note that this is only one of a few “Floer homologies” that were introduced by Floer in the late 80’s: the others include *Lagrangian intersection Floer homology* [Flo88a] (which has since evolved into the *Fukaya category*, see [Sei08, FOOO09]), and *instanton homology* [Flo88c], an extension of Donaldson’s gauge-theoretic smooth 4-manifold invariants to dimension three. The development of new Floer-type theories has since become a major industry; see [AS19] for a survey.




FIGURE 1.3. A star-shaped hypersurface in Euclidean space

1.3. Contact manifolds and the Weinstein conjecture

A Hamiltonian system on a symplectic manifold (W, ω) is called **autonomous** if the Hamiltonian $H : W \rightarrow \mathbb{R}$ does not depend on time. In this case, the Hamiltonian vector field X_H defined by

$$\omega(X_H, \cdot) = -dH$$

is time-independent and its orbits are confined to level sets of H . The images of these orbits on a given regular level set $H^{-1}(c)$ depend on the geometry of $H^{-1}(c)$, but not on H itself, as they are the integral curves (also known as **characteristics**) of the **characteristic line field** on $H^{-1}(c)$, defined as the unique direction spanned by a vector X such that $\omega(X, Y) = 0$ for all Y tangent to $H^{-1}(c)$. In 1978, Weinstein [Wei78] and Rabinowitz [Rab78] proved that certain kinds of regular level sets in symplectic manifolds are guaranteed to admit closed characteristics, hence implying the existence of periodic Hamiltonian orbits. In particular, this is true whenever $H^{-1}(c)$ is a *star-shaped* hypersurface in the standard symplectic \mathbb{R}^{2n} (see Figure 1.3).

The following symplectic interpretation of the star-shaped condition provides both an intuitive reason to believe Rabinowitz's existence result and motivation for the more general conjecture of Weinstein. In any symplectic manifold (W, ω) , a **Liouville vector field** is a smooth vector field V that satisfies

$$\mathcal{L}_V \omega = \omega.$$

By Cartan's formula for the Lie derivative, the 1-form λ defined by $\lambda := \omega(V, \cdot)$ satisfies $d\lambda = \omega$ if and only if V is a Liouville vector field; moreover, λ then also satisfies $\mathcal{L}_V \lambda = \lambda$, and it is referred to as a **Liouville form**. We sometimes say in this situation that the Liouville form λ and Liouville vector field V are **ω -dual** to each other. A hypersurface $M \subset (W, \omega)$ is said to be of **contact type** if it is transverse to a Liouville vector field defined on a neighborhood of M .

EXAMPLE 1.3.1. Using coordinates $(q_1, p_1, \dots, q_n, p_n)$ on \mathbb{R}^{2n} , the standard symplectic form is written as

$$\omega_{\text{std}} := \sum_{j=1}^n dp_j \wedge dq_j,$$

and the Liouville form $\lambda_{\text{std}} := \frac{1}{2} \sum_{j=1}^n (p_j dq_j - q_j dp_j)$ is dual to the radial Liouville vector field

$$V_{\text{std}} := \frac{1}{2} \sum_{j=1}^n \left(p_j \frac{\partial}{\partial p_j} + q_j \frac{\partial}{\partial q_j} \right).$$

Any star-shaped hypersurface is therefore of contact type.

EXERCISE 1.3.2. Suppose (W, ω) is a symplectic manifold of dimension $2n$, $M \subset W$ is a smoothly embedded and oriented hypersurface, V is a Liouville vector field defined near M and $\lambda := \omega(V, \cdot)$ is the dual Liouville form. Define a 1-form on M by $\alpha := \lambda|_{TM}$.

(a) Show that V is positively transverse to M if and only if α satisfies

$$(1.6) \quad \alpha \wedge (d\alpha)^{n-1} > 0.$$

(b) If V is positively transverse to M , choose $\epsilon > 0$ sufficiently small and consider the embedding

$$\Phi : (-\epsilon, \epsilon) \times M \hookrightarrow W : (r, x) \mapsto \varphi_V^r(x),$$

where φ_V^t denotes the time t flow of V . Show that

$$\Phi^* \lambda = e^r \alpha,$$

hence $\Phi^* \omega = d(e^r \alpha)$.

The above exercise presents any contact-type hypersurface $M \subset (W, \omega)$ as one member of a smooth 1-parameter family of contact-type hypersurfaces $M_r := \varphi_V^r(M) \subset W$, each canonically identified with M such that $\omega|_{TM_r} = e^r d\alpha$. In particular, the characteristic line fields on M_r are the same for all r , thus the existence of a closed characteristic on any of these implies that there also exists one on M . This observation has sometimes been used to prove such existence theorems, e.g. it is used in [HZ94, Chapter 4] to reduce Rabinowitz's result to an "almost existence" theorem based on symplectic capacities. This discussion hopefully makes the following conjecture seem believable.

CONJECTURE 1.3.3 (Weinstein conjecture, symplectic version). *Any closed contact-type hypersurface in a symplectic manifold admits a closed characteristic.*

Weinstein’s conjecture admits a natural rephrasing in the language of contact geometry. A 1-form α on an oriented $(2n - 1)$ -dimensional manifold M is called a (positive) **contact form** if it satisfies (1.6), and the resulting co-oriented hyperplane field

$$\xi := \ker \alpha \subset TM$$

is then called a (positive and co-oriented) **contact structure**.⁴ We call the pair (M, ξ) a **contact manifold**, and refer to a diffeomorphism $\varphi : M \rightarrow M'$ as a **contactomorphism** from (M, ξ) to (M', ξ') if φ_* maps ξ to ξ' and also preserves the respective co-orientations. Equivalently, if ξ and ξ' are defined via contact forms α and α' respectively, this means

$$\varphi^* \alpha' = f \alpha \quad \text{for some } f \in C^\infty(M, (0, \infty)).$$

Contact topology studies the category of contact manifolds (M, ξ) up to contactomorphism. The following basic result provides one good reason to regard ξ rather than α as the geometrically meaningful data, as the result holds for contact *structures*, but not for contact *forms*.

THEOREM 1.3.4 (Gray’s stability theorem). *If M is a closed $(2n - 1)$ -dimensional manifold and $\{\xi_t\}_{t \in [0,1]}$ is a smooth 1-parameter family of contact structures on M , then there exists a smooth 1-parameter family of diffeomorphisms $\{\varphi_t\}_{t \in [0,1]}$ such that $\varphi_0 = \text{Id}$ and $(\varphi_t)_* \xi_0 = \xi_t$.*

PROOF. See [Gei08, §2.2] or [Wenb, Theorem 1.6.12]. □

A corollary is that while the contact form α induced on a contact-type hypersurface $M \subset (W, \omega)$ via Exercise 1.3.2 is not unique, its induced contact structure is unique up to isotopy. Indeed, the space of all Liouville vector fields transverse to M is very large (e.g. one can add to V any sufficiently small Hamiltonian vector field), but it is *convex*, hence any two choices of the induced contact form α on M are connected by a smooth 1-parameter family of contact forms, implying an isotopy of contact structures via Gray’s theorem.

EXERCISE 1.3.5. If α is a nowhere zero 1-form on M and $\xi = \ker \alpha$, show that α is contact if and only if $d\alpha|_\xi$ defines a symplectic vector bundle structure on $\xi \rightarrow M$. Moreover, the orientation of ξ determined by this symplectic bundle structure is compatible with the co-orientation determined by α and the orientation of M for which $\alpha \wedge (d\alpha)^{n-1} > 0$.

The following definition is based on the fact that since $d\alpha|_\xi$ is nondegenerate when α is contact, $\ker d\alpha \subset TM$ is always 1-dimensional and transverse to ξ .

DEFINITION 1.3.6. Given a contact form α on M , the **Reeb vector field** is the unique vector field R_α that satisfies

$$d\alpha(R_\alpha, \cdot) \equiv 0, \quad \text{and} \quad \alpha(R_\alpha) \equiv 1.$$

⁴The adjective “positive” refers to the fact that the orientation of M agrees with the one determined by the volume form $\alpha \wedge (d\alpha)^{n-1}$; we call α a *negative* contact form if these two orientations disagree. It is also possible in general to define contact structures without co-orientations, but contact structures of this type will never appear in this book; for our purposes, the co-orientation is *always* considered to be part of the data of a contact structure.

EXERCISE 1.3.7. Show that the flow of any Reeb vector field R_α preserves both $\xi = \ker \alpha$ and the symplectic vector bundle structure $d\alpha|_\xi$.

CONJECTURE 1.3.8 (Weinstein conjecture, contact version). *On any closed contact manifold (M, ξ) with contact form α , the Reeb vector field R_α admits a periodic orbit.*

To see that this is equivalent to the symplectic version of the conjecture, observe that any contact manifold $(M, \xi = \ker \alpha)$ can be viewed as the contact-type hypersurface $\{0\} \times M$ in the open symplectic manifold

$$(\mathbb{R} \times M, d(e^r \alpha)),$$

called the **symplectization** of (M, ξ) .

EXERCISE 1.3.9. Recall that for any smooth manifold M , the cotangent bundle T^*M carries a tautological 1-form $\lambda_{\text{std}} \in \Omega^1(T^*M)$ that locally takes the form $\lambda_{\text{std}} = \sum_{j=1}^n p_j dq_j$ in any choice of local coordinates (q_1, \dots, q_n) on a neighborhood $\mathcal{U} \subset M$, with (p_1, \dots, p_n) denoting the induced coordinates on the cotangent fibers over \mathcal{U} . (We will discuss cotangent bundles in somewhat more detail in §3.8.) This defines a Liouville form, with $d\lambda_{\text{std}}$ defining the canonical symplectic structure of T^*M . Now if $\xi \subset TM$ is a co-oriented hyperplane field on M , consider the submanifold

$$S_\xi M := \{p \in T^*M \mid \ker p = \xi \text{ and } p(X) > 0 \forall X \in TM \text{ pos. transverse to } \xi\}.$$

Show that ξ is contact if and only if $S_\xi M$ is a symplectic submanifold of $(T^*M, d\lambda_{\text{std}})$, and the Liouville vector field on T^*M dual to λ_{std} is tangent to $S_\xi M$. Moreover, if ξ is contact, then any choice of contact form for ξ determines a diffeomorphism of $S_\xi M$ to $\mathbb{R} \times M$ identifying the Liouville form λ_{std} along $S_\xi M$ with $e^r \alpha$.

REMARK 1.3.10. Exercise 1.3.9 shows that up to symplectomorphism, our definition of the symplectization of (M, ξ) above actually depends only on ξ and not on α .

In 1993, Hofer [Hof93] introduced a new approach to the Weinstein conjecture that was based in part on ideas of Gromov and Floer. Fix a contact manifold (M, ξ) with contact form α , and let

$$\mathcal{J}(\alpha) \subset \mathcal{J}(\mathbb{R} \times M)$$

denote the nonempty and contractible space of all almost complex structures J on $\mathbb{R} \times M$ satisfying the following conditions:

- (1) The natural translation action on $\mathbb{R} \times M$ preserves J ;
- (2) $J\partial_r = R_\alpha$ and $JR_\alpha = -\partial_r$, where r denotes the canonical coordinate on the \mathbb{R} -factor in $\mathbb{R} \times M$;
- (3) $J\xi = \xi$ and $d\alpha(\cdot, J\cdot)|_\xi$ defines a bundle metric on ξ .

It is easy to check that any $J \in \mathcal{J}(\alpha)$ is compatible with the symplectic structure $d(e^r \alpha)$ on $\mathbb{R} \times M$. Moreover, if $\gamma : \mathbb{R} \rightarrow M$ is any periodic orbit of R_α with period $T > 0$, then for any $J \in \mathcal{J}(\alpha)$, the so-called **trivial cylinder**

$$u : \mathbb{R} \times S^1 \rightarrow \mathbb{R} \times M : (s, t) \mapsto (Ts, \gamma(Tt))$$

is a J -holomorphic curve. Following Floer, one version of Hofer’s idea would be to look for J -holomorphic cylinders that satisfy a finite energy condition as in Prop. 1.2.7, forcing them to approach trivial cylinders asymptotically—the existence of such a cylinder would then imply the existence of a closed Reeb orbit, and thus prove the Weinstein conjecture. The first hindrance is that the “obvious” definition of energy in this context,

$$\int_{\mathbb{R} \times S^1} u^* d(e^r \alpha),$$

is not very useful: this integral is infinite if u is a trivial cylinder. To circumvent this, notice that every $J \in \mathcal{J}(\alpha)$ is also compatible with any symplectic structure of the form

$$\omega_\varphi := d(e^{\varphi(r)} \alpha),$$

where φ is a function chosen freely from the set

$$(1.7) \quad \mathcal{T} := \{\varphi \in C^\infty(\mathbb{R}, (-1, 1)) \mid \varphi' > 0\}.$$

Essentially, choosing ω_φ means identifying $\mathbb{R} \times M$ with a subset of the bounded region $(-1, 1) \times M$, in which trivial cylinders have finite symplectic area. Since there is no preferred choice for the function φ , we define the **Hofer energy**⁵ of a J -holomorphic curve $u : \Sigma \rightarrow \mathbb{R} \times M$ by

$$(1.8) \quad E(u) := \sup_{\varphi \in \mathcal{T}} \int_{\Sigma} u^* \omega_\varphi.$$

This has the desired property of being finite for trivial cylinders, and it is also nonnegative, with strict positivity whenever u is not constant.

Another useful observation from [Hof93] was that if the goal is to find periodic orbits, then we need not restrict our attention to J -holomorphic *cylinders* in particular. One can more generally consider curves defined on an arbitrary *punctured* Riemann surface

$$\dot{\Sigma} := \Sigma \setminus \Gamma,$$

where (Σ, j) is a closed connected Riemann surface and $\Gamma \subset \Sigma$ is a finite set of punctures. For any $\zeta \in \Gamma$, one can find coordinates identifying some punctured neighborhood of ζ biholomorphically with the closed punctured disk

$$\dot{\mathbb{D}} := \mathbb{D} \setminus \{0\} \subset \mathbb{C},$$

and then identify this with either the positive or negative half-cylinder

$$Z_+ := [0, \infty) \times S^1, \quad Z_- := (-\infty, 0] \times S^1$$

via the biholomorphic maps

$$Z_+ \rightarrow \dot{\mathbb{D}} : (s, t) \mapsto e^{-2\pi(s+it)}, \quad Z_- \rightarrow \dot{\mathbb{D}} : (s, t) \mapsto e^{2\pi(s+it)}.$$

We will refer to such a choice as a (positive or negative) **holomorphic cylindrical coordinate** system near ζ , and in this way, we can present $(\dot{\Sigma}, j)$ as a *Riemann*

⁵Strictly speaking, the energy defined in (1.8) is not identical to the notion introduced in [Hof93] and used in many of Hofer’s papers, but it is equivalent to it in the sense that uniform bounds on either notion of energy imply uniform bounds on the other.




FIGURE 1.4. An asymptotically cylindrical holomorphic curve in a symplectization, with genus 1, one positive puncture and two negative punctures.

surface with cylindrical ends, i.e. the union of some compact Riemann surface with boundary with a finite collection of half-cylinders Z_{\pm} on which j takes the standard form $j\partial_s = \partial_t$. Note that the standard cylinder $\mathbb{R} \times S^1$ is a special case of this, as it can be identified biholomorphically with $S^2 \setminus \{0, \infty\}$. Another important special case is the plane, $\mathbb{C} = S^2 \setminus \{\infty\}$.

If $u : (\dot{\Sigma}, j) \rightarrow (\mathbb{R} \times M, J)$ is a J -holomorphic curve and $\zeta \in \Gamma$ is one of its punctures, we will say that u is **positively/negatively asymptotic** to a T -periodic Reeb orbit $\gamma : \mathbb{R} \rightarrow M$ at ζ if one can choose holomorphic cylindrical coordinates $(s, t) \in Z_{\pm}$ near ζ such that

$$u(s, t) = \exp_{(T_s, \gamma(Tt))} h(s, t) \quad \text{for } |s| \text{ sufficiently large,}$$

where $h(s, t)$ is a vector field along the trivial cylinder satisfying $h(s, \cdot) \rightarrow 0$ uniformly as $|s| \rightarrow \infty$, and the exponential map is defined with respect to any \mathbb{R} -invariant choice of Riemannian metric on $\mathbb{R} \times M$. We say that $u : (\dot{\Sigma}, j) \rightarrow (\mathbb{R} \times M, J)$ is **asymptotically cylindrical** if it is (positively or negatively) asymptotic to some closed Reeb orbit at each of its punctures. Note that this partitions the finite set of punctures $\Gamma \subset \Sigma$ into two subsets,

$$\Gamma = \Gamma^+ \cup \Gamma^-,$$

the *positive* and *negative* punctures respectively, see Figure 1.4.

EXERCISE 1.3.11. Suppose $u : (\dot{\Sigma}, j) \rightarrow (\mathbb{R} \times M, J)$ is an asymptotically cylindrical J -holomorphic curve, with the asymptotic orbit at each puncture $\zeta \in \Gamma^{\pm}$ denoted by γ_{ζ} , having period $T_{\zeta} > 0$. Show that

$$\sum_{\zeta \in \Gamma^+} T_{\zeta} - \sum_{\zeta \in \Gamma^-} T_{\zeta} = \int_{\dot{\Sigma}} u^* d\alpha \geq 0,$$

with equality if and only if the image of u is contained in that of a trivial cylinder. In particular, u must have at least one positive puncture unless it is constant. Show also that $E(u)$ is finite and satisfies an upper bound determined only by the periods of the positive asymptotic orbits.

The following analogue of Prop. 1.2.7 will be proved in Chapter 7. For simplicity, we shall state a weakened version of what Hofer proved in [Hof93], which did not require any nondegeneracy assumption. A T -periodic Reeb orbit $\gamma : \mathbb{R} \rightarrow M$ is called **nondegenerate** if the Reeb flow φ_α^t has the property that its linearization along the contact bundle (cf. Exercise 1.3.7),

$$d\varphi_\alpha^T(\gamma(0))|_{\xi_{\gamma(0)}} : \xi_{\gamma(0)} \rightarrow \xi_{\gamma(0)}$$

does not have 1 as an eigenvalue. Note that since R_α is not time-dependent, closed Reeb orbits are never completely isolated—they always exist in S^1 -parametrized families—but these families are isolated in the nondegenerate case. A **nondegenerate contact form** is one for which every closed Reeb orbit is nondegenerate. One can show that this condition is generic, meaning for instance that on any closed manifold, the nondegenerate contact forms constitute a C^∞ -dense subset of the space of all contact forms (see Remark 1.3.13 below). The following result is the contact analogue of Proposition 1.2.7.

PROPOSITION 1.3.12. *Suppose (M, ξ) is a closed contact manifold with a nondegenerate contact form α . If $u : (\tilde{\Sigma}, j) \rightarrow (\mathbb{R} \times M, J)$ is a J -holomorphic curve with $E(u) < \infty$ on a punctured Riemann surface such that none of the punctures are removable, then u is asymptotically cylindrical.*

The main results in [Hof93] state that under certain assumptions on a closed contact 3-manifold (M, ξ) , namely if either ξ is *overtwisted* (as defined in [Eli89]) or $\pi_2(M) \neq 0$, one can find for any contact form α on (M, ξ) and any $J \in \mathcal{J}(\alpha)$ a finite-energy J -holomorphic plane. By Proposition 1.3.12, this implies the existence of a contractible periodic Reeb orbit and thus proves the Weinstein conjecture in these settings.

REMARK 1.3.13. The standard genericity result mentioned above for nondegenerate contact forms can be proved in various ways, e.g. it follows from a slightly more general result about generic regular level sets in Hamiltonian systems proved in [Rob70]. A more direct proof via the Sard-Smale theorem that is similar in spirit to the transversality arguments in Chapter 9 may be found in the appendix of [ABW10].

1.4. Symplectic cobordisms and their completions

After the developments described in the previous three sections, it seemed natural that one might define invariants of contact manifolds via a Floer-type theory generated by closed Reeb orbits and counting asymptotically cylindrical holomorphic curves in symplectizations. This theory is what is now called SFT, and its basic structure was outlined in a paper by Eliashberg, Givental and Hofer [EGH00] in 2000, though some of its analytical foundations remain unfinished as of 2026. The term “field theory” is an allusion to “topological quantum field theories,” which associate vector spaces to certain geometric objects and morphisms to cobordisms between those objects. Thus in order to place SFT in its proper setting, we need to introduce symplectic cobordisms between contact manifolds.

Recall that if M_+ and M_- are smooth oriented closed manifolds of the same dimension, an oriented cobordism from M_- to M_+ is a compact smooth oriented manifold W with oriented boundary

$$\partial W \cong -M_- \amalg M_+,$$

where the symbol “ \cong ” in this setting means orientation-preserving diffeomorphism, and $-M_-$ denotes M_- with its orientation reversed. Given positive contact structures ξ_{\pm} on M_{\pm} , we say that a symplectic manifold (W, ω) is a **symplectic cobordism from (M_-, ξ_-) to (M_+, ξ_+)** if W is an oriented cobordism⁶ from M_- to M_+ such that both components of ∂W are contact-type hypersurfaces with induced contact structures isotopic to ξ_{\pm} . Note that our chosen orientation conventions imply that the Liouville vector field chosen near ∂W must point *outward* at M_+ and *inward* at M_- ; we say in this case that M_+ is a symplectically **convex** boundary component, while M_- is symplectically **concave**. As important special cases, (W, ω) is a **symplectic filling** of (M_+, ξ_+) if $M_- = \emptyset$, and it is a **symplectic cap** of (M_-, ξ_-) if $M_+ = \emptyset$. In the literature, fillings and caps are sometimes also referred to as *convex fillings* or *concave fillings* respectively.

The contact-type condition implies the existence of a Liouville form λ near ∂W with $d\lambda = \omega$, such that by Exercise 1.3.2, neighborhoods of M_+ and M_- in W can be identified with the collars (see Figure 1.5)

$$(-\epsilon, 0] \times M_+ \quad \text{or} \quad [0, \epsilon) \times M_-$$

respectively for sufficiently small $\epsilon > 0$, with λ taking the form

$$\lambda = e^r \alpha_{\pm},$$

where $\alpha_{\pm} := \lambda|_{TM_{\pm}}$ are contact forms for ξ_{\pm} , and r as usual denotes the canonical coordinate on the first factor in $\mathbb{R} \times M$. The **symplectic completion** of (W, ω) is the noncompact symplectic manifold $(\widehat{W}, \widehat{\omega})$ defined by attaching cylindrical ends to these collar neighborhoods (Figure 1.6):

$$(1.9) \quad (\widehat{W}, \widehat{\omega}) = ((-\infty, 0] \times M_-, d(e^r \alpha_-)) \cup_{M_-} (W, \omega) \cup_{M_+} ([0, \infty) \times M_+, d(e^r \alpha_+)).$$

In this context, the symplectization $(\mathbb{R} \times M, d(e^r \alpha))$ is symplectomorphic to the completion of the **trivial symplectic cobordism** $([0, 1] \times M, d(e^r \alpha))$ from $(M, \xi = \ker \alpha)$ to itself. More generally, the object in the following easy exercise can also sensibly be called a trivial symplectic cobordism:

EXERCISE 1.4.1. Suppose (M, ξ) is a closed contact manifold with contact form α , and $f_{\pm} : M \rightarrow \mathbb{R}$ is a pair of functions with $f_- < f_+$ everywhere. Show that the domain

$$\{(r, x) \in \mathbb{R} \times M \mid f_-(x) \leq r \leq f_+(x)\} \subset \mathbb{R} \times M$$

defines a symplectic cobordism from (M, ξ) to itself, with a global Liouville form $\lambda = e^r \alpha$ inducing contact forms $e^{f_-} \alpha$ and $e^{f_+} \alpha$ on its concave and convex boundaries respectively.

⁶We assume of course that W is assigned the orientation determined by its symplectic form.




FIGURE 1.5. A symplectic cobordism with concave boundary (M_-, ξ_-) and convex boundary (M_+, ξ_+) , with symplectic collar neighborhoods defined by flowing along Liouville vector fields near the boundary.




FIGURE 1.6. The completion of a symplectic cobordism

We say that (W, ω) is an **exact symplectic cobordism** or **Liouville cobordism** if the Liouville form λ can be extended from a neighborhood of ∂W to define a global primitive of ω on W . Equivalently, this means that ω admits a global Liouville vector field that points inward at M_- and outward at M_+ . An **exact filling** of (M_+, ξ_+) is an exact cobordism whose concave boundary is empty. Observe that if (W, ω) is exact, then its completion $(\widehat{W}, \widehat{\omega})$ also inherits a global Liouville form.

EXERCISE 1.4.2. Use Stokes' theorem to show that there is no such thing as an exact symplectic cap.

The above exercise hints at an important difference between cobordisms in the *symplectic* as opposed to the *oriented smooth* category: symplectic cobordisms are not generally reversible. If W is an oriented cobordism from M_- to M_+ , then reversing the orientation of W produces an oriented cobordism from M_+ to M_- . But one cannot simply reverse orientations in the symplectic category, since the orientation is determined by the symplectic form. For example, many obstructions to the existence of symplectic fillings of given contact manifolds are known—some of them defined in terms of SFT—but there are no obstructions at all to symplectic caps, in fact it is known that all closed contact manifolds admit them (see [EH02, CE20, Laz20]).

The definitions for holomorphic curves in symplectizations in the previous section generalize to completions of symplectic cobordisms in a fairly straightforward way, since these completions look exactly like symplectizations outside of a compact subset. Define

$$\mathcal{J}(W, \omega, \alpha_+, \alpha_-) \subset \mathcal{J}(\widehat{W})$$

as the space of all almost complex structures J on \widehat{W} such that

$$J|_W \in \mathcal{J}(W, \omega), \quad J|_{[0, \infty) \times M_+} \in \mathcal{J}(\alpha_+) \quad \text{and} \quad J|_{(-\infty, 0] \times M_-} \in \mathcal{J}(\alpha_-).$$

Occasionally it is useful to relax the compatibility condition on W to tameness,⁷ i.e. $J|_W \in \mathcal{J}_\tau(W, \omega)$, producing a space that we shall denote by

$$\mathcal{J}_\tau(W, \omega, \alpha_+, \alpha_-) \subset \mathcal{J}(\widehat{W}).$$

As in Prop. 1.1.3, both of these spaces are nonempty and contractible. We can then consider asymptotically cylindrical J -holomorphic curves

$$u : (\dot{\Sigma} = \Sigma \setminus (\Gamma^+ \cup \Gamma^-), j) \rightarrow (\widehat{W}, J),$$

which are proper maps asymptotic to closed orbits of R_{α_\pm} in M_\pm at punctures in Γ^\pm , see Figure 1.7.

One must again tinker with the symplectic form on \widehat{W} in order to define a notion of energy that is finite when we need it to be. We generalize (1.7) as

$$\mathcal{T} := \{ \varphi \in C^\infty(\mathbb{R}, (-1, 1)) \mid \varphi' > 0 \text{ and } \varphi(r) = r \text{ near } r = 0 \},$$

and associate to each $\varphi \in \mathcal{T}$ a symplectic form $\widehat{\omega}_\varphi$ on \widehat{W} defined by

$$\widehat{\omega}_\varphi := \begin{cases} d(e^{\varphi(r)} \alpha_+) & \text{on } [0, \infty) \times M_+, \\ \omega & \text{on } W, \\ d(e^{\varphi(r)} \alpha_-) & \text{on } (-\infty, 0] \times M_-. \end{cases}$$

One can again check that every $J \in \mathcal{J}(W, \omega, \alpha_+, \alpha_-)$ or $\mathcal{J}_\tau(W, \omega, \alpha_+, \alpha_-)$ is tamed by $\widehat{\omega}_\varphi$ for every $\varphi \in \mathcal{T}$. Thus it makes sense to define the **energy** of $u : (\dot{\Sigma}, j) \rightarrow (\widehat{W}, J)$

⁷It seems natural to wonder whether one could not also relax the conditions on the cylindrical ends and require $J|_{\xi_\pm}$ to be tamed by $d\alpha_\pm|_{\xi_\pm}$ instead of compatible with it. I do not currently know whether this works, but in later chapters we will see some reasons to worry that it might not (see §6.7.2).




FIGURE 1.7. An asymptotically cylindrical holomorphic curve in a completed symplectic cobordism, with genus 2, one positive puncture and two negative punctures.

by

$$E(u) := \sup_{\varphi \in \mathcal{T}} \int_{\dot{\Sigma}} u^* \widehat{\omega}_\varphi.$$

It will be a straightforward matter to generalize Proposition 1.3.12 and show that finite energy implies asymptotically cylindrical behavior in completed cobordisms.

EXERCISE 1.4.3. Show that if (W, ω) is an exact cobordism, then every asymptotically cylindrical J -holomorphic curve in \widehat{W} has at least one positive puncture.

1.5. Contact homology and SFT

We can now sketch the algebraic structure of SFT. We shall ignore or suppress several pesky details that are best dealt with later, some of them algebraic, others analytical. Due to analytical problems, some of the “theorems” that we shall (often imprecisely) state in this section are not yet provable at the current level of technology, though we expect that they will be in the foreseeable future. We shall use quotation marks to indicate this caveat wherever appropriate.

The standard versions of SFT all define homology theories with varying levels of algebraic structure which are meant to be invariants of a contact manifold (M, ξ) . The chain complexes always depend on certain auxiliary choices, including a nondegenerate contact form α and a generic $J \in \mathcal{J}(\alpha)$. The generators consist of formal variables q_γ , one for each⁸ closed Reeb orbit γ . In the most straightforward generalization of Hamiltonian Floer homology, the chain complex is simply a graded

⁸Actually, I should be making a distinction here between “good” and “bad” Reeb orbits, but let’s discuss that later; see Chapter 12.

\mathbb{Q} -vector space generated by the variables q_γ , and the boundary map is defined by

$$\partial_{\text{CCH}} q_\gamma = \sum_{\gamma'} \# (\mathcal{M}(\gamma, \gamma')/\mathbb{R}) q_{\gamma'},$$

where $\mathcal{M}(\gamma, \gamma')$ is the moduli space of J -holomorphic cylinders in $\mathbb{R} \times M$ with a positive puncture asymptotic to γ and a negative puncture asymptotic to γ' , and the sum ranges over all orbits γ' for which this moduli space is 1-dimensional. The count $\# (\mathcal{M}(\gamma, \gamma')/\mathbb{R})$ is rational, as it includes rational weighting factors that depend on combinatorial information and are best not discussed right now.⁹

“THEOREM” 1.5.1. *If α admits no contractible Reeb orbits, then $\partial_{\text{CCH}}^2 = 0$, and the resulting homology is independent of the choices of α with this property and generic $J \in \mathcal{J}(\alpha)$.*

The invariant arising from this result is known as **cylindrical contact homology**, and it is sometimes quite easy to work with when it is well defined, though it has the disadvantage of not always being defined. Namely, the relation $\partial_{\text{CCH}}^2 = 0$ can fail if α admits contractible Reeb orbits, because unlike in Floer homology, the compactification of the space of cylinders $\mathcal{M}(\gamma, \gamma')$ generally includes objects that are not broken cylinders. In fact, the objects arising in the “SFT compactification” of moduli spaces of finite-energy curves in completed cobordisms can be quite elaborate, see Figure 1.8. The combinatorics of the situation are not so bad however if the cobordism is exact, as is the case for a symplectization: Exercise 1.4.3 then prevents curves without positive ends from appearing. The only possible degenerations for cylinders then consist of broken configurations whose levels each have *exactly one positive puncture* and arbitrary negative punctures; moreover, all but one of the negative punctures must eventually be capped off by planes, which is why “Theorem” 1.5.1 holds in the absence of planes.

If planes do exist, then one can account for them by defining the chain complex as an *algebra* rather than a vector space, producing the theory known as **contact homology**. For this, the chain complex is taken to be a graded unital algebra over \mathbb{Q} , and we define

$$\partial_{\text{CH}} q_\gamma = \sum_{(\gamma_1, \dots, \gamma_m)} \# (\mathcal{M}(\gamma; \gamma_1, \dots, \gamma_m)/\mathbb{R}) q_{\gamma_1} \cdots q_{\gamma_m},$$

with $\mathcal{M}(\gamma; \gamma_1, \dots, \gamma_m)$ denoting the moduli space of punctured J -holomorphic spheres in $\mathbb{R} \times M$ with a positive puncture at γ and m negative punctures at the orbits $\gamma_1, \dots, \gamma_m$, and the sum ranges over all integers $m \geq 0$ and all m -tuples of orbits for which the moduli space is 1-dimensional. The action of ∂_{CH} is then extended to the whole algebra via a graded Leibniz rule

$$\partial_{\text{CH}}(q_\gamma q_{\gamma'}) := (\partial_{\text{CH}} q_\gamma) q_{\gamma'} + (-1)^{|\gamma|} q_\gamma (\partial_{\text{CH}} q_{\gamma'}).$$

The general compactness and gluing theory for genus zero curves with one positive puncture now implies:

⁹Similar combinatorial factors are hidden behind the symbol “#” in our definitions of ∂_{CH} and \mathbf{H} , and will be discussed in earnest in Chapter 13.




FIGURE 1.8. Degeneration of a sequence u_k of finite energy punctured holomorphic curves with genus 2, one positive puncture and two negative punctures in a symplectic cobordism. The limiting holomorphic building $(v_1^+, v_0, v_1^-, v_2^-, v_3^-)$ in this example has one upper level living in the symplectization $\mathbb{R} \times M_+$, a main level living in \widehat{W} , and three lower levels, each of which is a (possibly disconnected) finite-energy punctured nodal holomorphic curve in $\mathbb{R} \times M_-$. The building has arithmetic genus 2 and the same numbers of positive and negative punctures as u_k .

“THEOREM” 1.5.2. $\partial_{\text{CH}}^2 = 0$, and the resulting homology is (as a graded unital \mathbb{Q} -algebra) independent of the choices α and J .

Maybe you’ve noticed the pattern: in order to accommodate more general classes of holomorphic curves, we need to add more algebraic structure. The **full SFT** algebra counts all rigid holomorphic curves in $\mathbb{R} \times M$, including all combinations of positive and negative punctures and all genera. Here is a brief picture of what it looks like. Counting all the 1-dimensional moduli spaces of J -holomorphic curves modulo \mathbb{R} -translation in $\mathbb{R} \times M$ produces a formal power series

$$\mathbf{H} := \sum \# \left(\mathcal{M}_g(\gamma_1^+, \dots, \gamma_{m_+}^+; \gamma_1^-, \dots, \gamma_{m_-}^-) / \mathbb{R} \right) q_{\gamma_1^-} \cdots q_{\gamma_{m_-}^-} p_{\gamma_1^+} \cdots p_{\gamma_{m_+}^+} \hbar^{g-1},$$

where the sum ranges over all integers $g, m_+, m_- \geq 0$ and tuples of orbits, \hbar and p_γ (one for each orbit γ) are additional formal variables, and

$$\mathcal{M}_g(\gamma_1^+, \dots, \gamma_{m_+}^+; \gamma_1^-, \dots, \gamma_{m_-}^-)$$

denotes the moduli space of J -holomorphic curves in $\mathbb{R} \times M$ with genus g , m_+ positive punctures at the orbits $\gamma_1^+, \dots, \gamma_{m_+}^+$, and m_- negative punctures at the orbits $\gamma_1^-, \dots, \gamma_{m_-}^-$. We can regard \mathbf{H} as an operator on a graded algebra \mathfrak{W} of formal power series in the variables $\{p_\gamma\}$, $\{q_\gamma\}$ and \hbar , equipped with a graded bracket operation that satisfies the quantum mechanical commutation relation

$$[p_\gamma, q_\gamma] = \kappa_\gamma \hbar,$$

where κ_γ is a combinatorial factor that is best ignored for now. Note that due to the signs that accompany the grading, odd elements $\mathbf{F} \in \mathfrak{W}$ need not satisfy $[\mathbf{F}, \mathbf{F}] = 0$, and \mathbf{H} itself is an odd element, thus the following statement is nontrivial; in fact, it is the algebraic manifestation of the general compactness and gluing theory for punctured holomorphic curves in symplectizations.

“THEOREM” 1.5.3. $[\mathbf{H}, \mathbf{H}] = 0$, hence by the graded Jacobi identity, \mathbf{H} determines an operator

$$D_{\text{SFT}} : \mathfrak{W} \rightarrow \mathfrak{W} : \mathbf{F} \mapsto [\mathbf{H}, \mathbf{F}]$$

satisfying $D_{\text{SFT}}^2 = 0$. The resulting homology depends on (M, ξ) but not on the auxiliary choices α and J .

It takes some time to understand how pictures such as Figure 1.8 translate into algebraic relations like $[\mathbf{H}, \mathbf{H}] = 0$, but this is a subject we’ll come back to. There is also an intermediate theory between contact homology and full SFT, called **rational SFT**, which counts only genus zero curves with arbitrary positive and negative punctures. Algebraically, it is obtained from the full SFT algebra as a “semiclassical approximation” by discarding higher-order factors of \hbar , so that the commutation bracket in \mathfrak{W} becomes a graded Poisson bracket. We will discuss all of this in Chapter 13.

1.6. Two applications

We briefly mention two applications that we will be able to establish rigorously using the methods developed in this book. Since SFT itself is not yet well defined in full generality, this sometimes means using SFT for *inspiration*, while proving corollaries via more direct methods.

1.6.1. Tight contact structures on \mathbb{T}^3 . The 3-torus $\mathbb{T}^3 = S^1 \times S^1 \times S^1$ with coordinates (t, θ, ϕ) admits a sequence of contact structures

$$\xi_k := \ker(\cos(2\pi kt) d\theta + \sin(2\pi kt) d\phi),$$

one for each $k \in \mathbb{N}$. These cannot be distinguished from each other by any classical invariants, e.g. they all have the same Euler class, in fact they are all homotopic as co-oriented 2-plane fields. Nonetheless:

THEOREM 1.6.1. *For $k \neq \ell$, (\mathbb{T}^3, ξ_k) and (\mathbb{T}^3, ξ_ℓ) are not contactomorphic.*




FIGURE 1.9. This exact symplectic cobordism does not exist.

We will be able to prove this in Chapter 11 by rigorously defining and computing cylindrical contact homology for a suitable choice of contact forms on (\mathbb{T}^3, ξ_k) .

1.6.2. Filling and cobordism obstructions. Consider a closed connected and oriented surface Σ presented as $\Sigma_+ \cup_{\Gamma} \Sigma_-$, where $\Sigma_{\pm} \subset \Sigma$ are each (not necessarily connected) compact surfaces with a common boundary Γ . By an old result of Lutz [Lut77], the 3-manifold $S^1 \times \Sigma$ admits a unique isotopy class of S^1 -invariant contact structures ξ_{Γ} such that the loops $S^1 \times \{z\}$ are positively/negatively transverse to ξ_{Γ} for $z \in \overset{\circ}{\Sigma}_{\pm}$ and tangent to ξ_{Γ} for $z \in \Gamma$. Now for each $k \in \mathbb{N}$, define

$$(V_k, \xi_k) := (S^1 \times \Sigma, \xi_{\Gamma})$$

where $\Sigma = \Sigma_+ \cup_{\Gamma} \Sigma_-$ is chosen such that Γ has k connected components, Σ_- is connected with genus zero, and Σ_+ is connected with positive genus (see Figure 1.9).

THEOREM 1.6.2. *The contact manifolds (V_k, ξ_k) do not admit any symplectic fillings. Moreover, if $k > \ell$, then there exists no exact symplectic cobordism from (V_k, ξ_k) to (V_{ℓ}, ξ_{ℓ}) .*

For these examples, one can use explicit constructions from [Wen13, Avd21] to show that non-exact cobordisms from (V_k, ξ_k) to (V_{ℓ}, ξ_{ℓ}) do exist, and so do exact cobordisms from (V_{ℓ}, ξ_{ℓ}) to (V_k, ξ_k) , thus both the directionality of the cobordism relation and the distinction between exact and non-exact are crucial. The proof of the theorem, due to the author with Latschev and Hutchings [LW11], uses a numerical contact invariant based on the full SFT algebra—in particular, the curves that cause this phenomenon have multiple positive ends and are thus not seen by

contact homology. We will introduce the relevant numerical invariant in Chapter 14 and compute it for these examples in Chapter 17.

CHAPTER 2

Basics on holomorphic curves

Contents

2.1. Linearized Cauchy-Riemann operators	27
2.2. Some useful Sobolev inequalities	27
2.3. The fundamental elliptic estimate	27
2.4. Regularity	27
2.4.1. The linear case	27
2.4.2. The nonlinear case: bootstrapping	27
2.4.3. The nonlinear case: from $W^{1,p} \cap C^0$ to $W^{1,q}$	27
2.5. Linear local existence and the similarity principle	27
2.6. Simple curves and multiple covers	27
2.7. Nonlinear local existence	27
2.8. The nonlinear equation on push-offs	27

2.1. Linearized Cauchy-Riemann operators

2.2. Some useful Sobolev inequalities

PROPOSITION 2.2.1 (Banach algebra property). *to be written*

PROPOSITION 2.2.2 (C^k -continuity property). *to be written*

PROPOSITION 2.2.3. *to be written*

2.3. The fundamental elliptic estimate

2.4. Regularity

2.4.1. The linear case.

2.4.2. The nonlinear case: bootstrapping.

2.4.3. The nonlinear case: from $W^{1,p} \cap C^0$ to $W^{1,q}$.

2.5. Linear local existence and the similarity principle

2.6. Simple curves and multiple covers

2.7. Nonlinear local existence

2.8. The nonlinear equation on push-offs

CHAPTER 3

Asymptotic operators

Contents

3.1. Stable Hamiltonian structures and Reeb orbits	29
3.2. The linearization in Morse homology	29
3.3. The Hessian of the contact action functional	29
3.4. Spectral flow	29
3.4.1. Geometry in the space of Fredholm operators	29
3.4.2. Symmetric operators of index zero	29
3.4.3. Perturbation of eigenvalues	29
3.4.4. Homotopies of eigenvalues	30
3.5. The Conley-Zehnder index of a nondegenerate orbit	30
3.6. Winding numbers of eigenfunctions	30
3.7. Elliptic and hyperbolic orbits	30
3.8. CZ = Morse for geodesics	30
3.8.1. The unit cotangent bundle	30
3.8.2. The energy functional and its Hessian	30
3.8.3. Conley-Zehnder indices of lifted geodesics	30
3.9. Morse-Bott families	30
3.9.1. Clean intersection conditions	30
3.9.2. The perturbed Conley-Zehnder indices	30
3.9.3. The Robbin-Salamon index	30

3.1. Stable Hamiltonian structures and Reeb orbits

3.2. The linearization in Morse homology

3.3. The Hessian of the contact action functional

EXERCISE 3.3.1. to be written

3.4. Spectral flow

REMARK 3.4.1. to be written

REMARK 3.4.2. to be written

3.4.1. Geometry in the space of Fredholm operators.

3.4.2. Symmetric operators of index zero.

3.4.3. Perturbation of eigenvalues.

3.4.4. Homotopies of eigenvalues.

LEMMA 3.4.3 (see Appendix C). *to be written*

3.5. The Conley-Zehnder index of a nondegenerate orbit**3.6. Winding numbers of eigenfunctions****3.7. Elliptic and hyperbolic orbits****3.8. CZ = Morse for geodesics****3.8.1. The unit cotangent bundle.****3.8.2. The energy functional and its Hessian.****3.8.3. Conley-Zehnder indices of lifted geodesics.****3.9. Morse-Bott families****3.9.1. Clean intersection conditions.****3.9.2. The perturbed Conley-Zehnder indices.****3.9.3. The Robbin-Salamon index.**

CHAPTER 4

Fredholm theory with cylindrical ends

Contents

4.1. Cauchy-Riemann operators with punctures	31
4.2. A lemma on semi-Fredholm operators	31
4.3. Some global regularity estimates	31
4.4. Translation-invariant operators on the cylinder	31
4.5. Proof of the semi-Fredholm property	31
4.6. Exponential decay	31
4.7. Formal adjoints and proof of the Fredholm property	31
4.8. The asymptotic formula	31

4.1. Cauchy-Riemann operators with punctures
4.2. A lemma on semi-Fredholm operators
4.3. Some global regularity estimates
4.4. Translation-invariant operators on the cylinder
4.5. Proof of the semi-Fredholm property
4.6. Exponential decay
4.7. Formal adjoints and proof of the Fredholm property
4.8. The asymptotic formula

CHAPTER 5

The index formula

Contents

5.1. Riemann-Roch with punctures	33
5.2. Some remarks on the formal adjoint	33
5.3. The index zero case on a torus	33
5.4. A Weitzenböck formula for Cauchy-Riemann operators	33
5.5. Large antilinear perturbations and energy concentration	33
5.6. Two Cauchy-Riemann type problems on the plane	33
5.7. A linear gluing argument	33
5.8. Antilinear deformations of asymptotic operators	33

5.1. Riemann-Roch with punctures

5.2. Some remarks on the formal adjoint

5.3. The index zero case on a torus

5.4. A Weitzenböck formula for Cauchy-Riemann operators

5.5. Large antilinear perturbations and energy concentration

5.6. Two Cauchy-Riemann type problems on the plane

5.7. A linear gluing argument

5.8. Antilinear deformations of asymptotic operators

CHAPTER 6

Symplectic cobordisms and moduli spaces

Contents

6.1. Stable Hamiltonian structures	35
6.1.1. Hamiltonian structures and dynamics	35
6.1.2. Collar neighborhoods and cobordisms	35
6.1.3. Stability	35
6.2. Almost complex manifolds with cylindrical ends	35
6.2.1. Symplectizations	35
6.2.2. Completed cobordisms	35
6.3. Examples of stable Hamiltonian structures	35
6.3.1. The contact case	35
6.3.2. The Floer case	36
6.4. Moduli spaces of asymptotically cylindrical curves	36
6.4.1. Curves with fixed asymptotic orbits	36
6.4.2. Asymptotic orbits in families	36
6.5. Asymptotic regularity	36
6.6. Simple curves and multiple covers revisited	36
6.7. Possible generalizations	36
6.7.1. Asymptotically cylindrical ends	36
6.7.2. Tame but not compatible	36
6.7.3. Framed but not stable	36
6.7.4. SFT without symplectic structures?	36

6.1. Stable Hamiltonian structures

6.1.1. Hamiltonian structures and dynamics.

6.1.2. Collar neighborhoods and cobordisms.

6.1.3. Stability.

6.2. Almost complex manifolds with cylindrical ends

6.2.1. Symplectizations.

6.2.2. Completed cobordisms.

6.3. Examples of stable Hamiltonian structures

6.3.1. The contact case.

6.3.2. The Floer case.

6.4. Moduli spaces of asymptotically cylindrical curves

6.4.1. Curves with fixed asymptotic orbits.

6.4.2. Asymptotic orbits in families.

6.5. Asymptotic regularity

6.6. Simple curves and multiple covers revisited

6.7. Possible generalizations

6.7.1. Asymptotically cylindrical ends.

6.7.2. Tame but not compatible.

6.7.3. Framed but not stable.

6.7.4. SFT without symplectic structures?

CHAPTER 7

Asymptotics and compactness

Contents

7.1. Removal of singularities	37
7.2. Finite energy and asymptotics	37
7.3. Degenerations of holomorphic curves	37
7.3.1. Bubbling	37
7.3.2. Breaking	37
7.3.3. The Deligne-Mumford space of Riemann surfaces	37
7.4. The SFT compactness theorem	37
7.4.1. Nodal curves	37
7.4.2. Holomorphic buildings	37
7.4.3. Convergence	37
7.4.4. Symplectizations	37
7.4.5. Stretching the neck	37
7.5. Compactness results for biholomorphic maps	37

7.1. Removal of singularities

7.2. Finite energy and asymptotics

7.3. Degenerations of holomorphic curves

7.3.1. Bubbling.

7.3.2. Breaking.

7.3.3. The Deligne-Mumford space of Riemann surfaces.

7.4. The SFT compactness theorem

7.4.1. Nodal curves.

7.4.2. Holomorphic buildings.

7.4.3. Convergence.

7.4.4. Symplectizations.

7.4.5. Stretching the neck.

7.5. Compactness results for biholomorphic maps

CHAPTER 8

Smoothness of the moduli space

Contents

8.1. The main result on regular curves	39
8.2. Functional-analytic setup	39
8.3. Moduli of complex structures	39
8.3.1. Teichmüller space and automorphism groups	39
8.3.2. Teichmüller slices	39
8.3.3. Adding marked points	39
8.4. Fredholm regularity and the implicit function theorem	39
8.5. Evaluation and forgetful maps	39

8.1. The main result on regular curves

8.2. Functional-analytic setup

8.3. Moduli of complex structures

8.3.1. Teichmüller space and automorphism groups.

8.3.2. Teichmüller slices.

8.3.3. Adding marked points.

8.4. Fredholm regularity and the implicit function theorem

8.5. Evaluation and forgetful maps

CHAPTER 9

Transversality

Contents

9.1. A paradigm for genericity arguments	41
9.2. Generic transversality in cobordisms	41
9.2.1. A theorem for somewhere injective curves	41
9.2.2. The universal moduli space	41
9.2.3. Applying the Sard-Smale theorem	41
9.2.4. From C_ϵ to C^∞	41
9.3. Generic transversality in symplectizations	41
9.3.1. Main results in the \mathbb{R} -invariant setting	41
9.3.2. Injective points of the projected curve	41
9.3.3. Smoothness of the universal moduli space	41
9.4. Transversality of constraints	41
9.4.1. The evaluation and forgetful maps	41
9.4.2. Constraints on derivatives	42

9.1. A paradigm for genericity arguments

LEMMA 9.1.1. *to be written*

9.2. Generic transversality in cobordisms

9.2.1. A theorem for somewhere injective curves.

9.2.2. The universal moduli space.

9.2.3. Applying the Sard-Smale theorem.

9.2.4. From C_ϵ to C^∞ .

9.3. Generic transversality in symplectizations

9.3.1. Main results in the \mathbb{R} -invariant setting.

9.3.2. Injective points of the projected curve.

9.3.3. Smoothness of the universal moduli space.

9.4. Transversality of constraints

9.4.1. The evaluation and forgetful maps.

9.4.2. Constraints on derivatives.

CHAPTER 10

Gluing

Contents

10.1. Moduli spaces of holomorphic buildings	43
10.1.1. Notation and conventions	43
10.1.2. Automorphism groups	43
10.1.3. The deformation operator	43
10.2. Deligne-Mumford space is an orbifold	43

10.1. Moduli spaces of holomorphic buildings

10.1.1. Notation and conventions.

10.1.2. Automorphism groups.

10.1.3. The deformation operator.

10.2. Deligne-Mumford space is an orbifold

CHAPTER 11

Cylindrical contact homology and the tight 3-tori

Contents

11.1. Contact structures on \mathbb{T}^3 and Giroux torsion	45
11.2. Definition of cylindrical contact homology	45
11.2.1. Preliminary remarks	45
11.2.2. A compactness result for cylinders	45
11.2.3. The chain complex	45
11.2.4. The homology	45
11.2.5. Chain maps	45
11.2.6. Chain homotopies	45
11.2.7. Proof of invariance	45
11.3. Computing $HC_*(\mathbb{T}^3, \xi_k)$	45
11.3.1. The Morse-Bott setup	45
11.3.2. A digression on the Floer equation	45
11.3.3. Admissible data for (\mathbb{T}^3, ξ_k)	45

11.1. Contact structures on \mathbb{T}^3 and Giroux torsion

11.2. Definition of cylindrical contact homology

11.2.1. Preliminary remarks.

11.2.2. A compactness result for cylinders.

11.2.3. The chain complex.

11.2.4. The homology.

11.2.5. Chain maps.

11.2.6. Chain homotopies.

11.2.7. Proof of invariance.

11.3. Computing $HC_*(\mathbb{T}^3, \xi_k)$

11.3.1. The Morse-Bott setup.

11.3.2. A digression on the Floer equation.

11.3.3. Admissible data for (\mathbb{T}^3, ξ_k) .

CHAPTER 12

Coherent orientations

Contents

12.1. Gluing maps and coherence	47
12.2. Permutations of punctures and bad orbits	47
12.3. Orienting moduli spaces in general	47
12.4. The determinant line bundle	47
12.5. Determinant bundles of moduli spaces	47
12.6. An algorithm for coherent orientations	47
12.7. Permutations and bad orbits revisited	47

12.1. Gluing maps and coherence

12.2. Permutations of punctures and bad orbits

12.3. Orienting moduli spaces in general

12.4. The determinant line bundle

12.5. Determinant bundles of moduli spaces

12.6. An algorithm for coherent orientations

12.7. Permutations and bad orbits revisited

CHAPTER 13

The generating function of SFT

Contents

13.1. Some important caveats on transversality	49
13.2. Auxiliary data, grading and supercommutativity	49
13.3. The definition of H and commutators	49
13.4. Interlude: Orbifolds and branched manifolds	49
13.4.1. How to count zeroes in an orbifold	49
13.4.2. Multivalued perturbations	49
13.4.3. The inhomogeneous Cauchy-Riemann equation	49
13.5. Cylindrical contact homology revisited	49
13.6. Combinatorics of gluing	49
13.7. Some remarks on torsion, coefficients, and conventions	49
13.7.1. What if $H_1(M)$ has torsion?	49
13.7.2. Combinatorial conventions	49
13.7.3. Coefficients: \mathbb{Q} , \mathbb{Z} or \mathbb{Z}_2 ?	49

13.1. Some important caveats on transversality
13.2. Auxiliary data, grading and supercommutativity
13.3. The definition of H and commutators
13.4. Interlude: Orbifolds and branched manifolds
13.4.1. How to count zeroes in an orbifold.
13.4.2. Multivalued perturbations.
13.4.3. The inhomogeneous Cauchy-Riemann equation.
13.5. Cylindrical contact homology revisited
13.6. Combinatorics of gluing
13.7. Some remarks on torsion, coefficients, and conventions
13.7.1. What if $H_1(M)$ has torsion?
13.7.2. Combinatorial conventions.
13.7.3. Coefficients: \mathbb{Q} , \mathbb{Z} or \mathbb{Z}_2 ?

CHAPTER 14

Contact invariants

Contents

14.1. The Eliashberg-Givental-Hofer package	51
14.1.1. Full SFT as a Weyl superalgebra	51
14.1.2. The semiclassical limit: rational SFT	51
14.1.3. The contact homology algebra	51
14.1.4. Algebraic overtwistedness	51
14.2. SFT generating functions for cobordisms	51
14.2.1. Weak, strong and stable cobordisms	51
14.2.2. Counting disconnected index 0 curves	51
14.3. Full SFT as a BV_∞-algebra	51
14.3.1. Cobordism maps and invariance	51
14.3.2. Algebraic torsion	51

14.1. The Eliashberg-Givental-Hofer package

14.1.1. Full SFT as a Weyl superalgebra.

14.1.2. The semiclassical limit: rational SFT.

14.1.3. The contact homology algebra.

14.1.4. Algebraic overtwistedness.

14.2. SFT generating functions for cobordisms

14.2.1. Weak, strong and stable cobordisms.

14.2.2. Counting disconnected index 0 curves.

14.3. Full SFT as a BV_∞ -algebra

14.3.1. Cobordism maps and invariance.

14.3.2. Algebraic torsion.

CHAPTER 15

Transversality and counting singularities in dimension four

Contents

15.1. Automatic transversality	53
15.1.1. The statement	53
15.1.2. The normal Cauchy-Riemann operator	53
15.1.3. Counting zeroes on line bundles with punctures	53
15.1.4. Proof of Theorem 15.1.1 and generalizations	53
15.1.5. An alternative proof for $g = \#\Gamma_0 = 0$	53
15.2. Curves in symplectizations of 3-manifolds	53
15.3. Implicit function theorems for local foliations	53
15.3.1. Immersed curves with no automorphisms	53
15.3.2. Embedded curves asymptotic to distinct simple orbits	53
15.3.3. Immersed multiply covered curves	53
15.4. Consequences for coherent orientations	53

15.1. Automatic transversality

15.1.1. The statement.

THEOREM 15.1.1. *to be written*

15.1.2. The normal Cauchy-Riemann operator.

15.1.3. Counting zeroes on line bundles with punctures.

15.1.4. Proof of Theorem **15.1.1** and generalizations.

15.1.5. An alternative proof for $g = \#\Gamma_0 = 0$.

15.2. Curves in symplectizations of 3-manifolds

15.3. Implicit function theorems for local foliations

15.3.1. Immersed curves with no automorphisms.

15.3.2. Embedded curves asymptotic to distinct simple orbits.

15.3.3. Immersed multiply covered curves.

15.4. Consequences for coherent orientations

CHAPTER 16

Intersection theory for punctured holomorphic curves

Contents

16.1. Prologue	55
16.2. Homotopy-invariant intersection numbers	55
16.3. The adjunction formula	55
16.4. Local foliations: the general case	55

16.1. Prologue

16.2. Homotopy-invariant intersection numbers

16.3. The adjunction formula

16.4. Local foliations: the general case

CHAPTER 17

Torsion computations and applications

Contents

17.1. Some J-holomorphic foliations	57
17.1.1. Gradient flow lines as holomorphic cylinders	57
17.1.2. Domains with \mathbb{T}^2 -symmetry	57
17.1.3. Open books with holomorphic pages	57
17.2. Contact homology of overtwisted contact manifolds	57
17.3. Examples with higher-order algebraic torsion	57
17.4. Rigorous obstructions to fillings and cobordisms	57

17.1. Some J -holomorphic foliations

17.1.1. Gradient flow lines as holomorphic cylinders.

17.1.2. Domains with \mathbb{T}^2 -symmetry.

17.1.3. Open books with holomorphic pages.

17.2. Contact homology of overtwisted contact manifolds

17.3. Examples with higher-order algebraic torsion

17.4. Rigorous obstructions to fillings and cobordisms

APPENDIX A

Sobolev spaces

Contents

A.1. Approximation, extension and embedding theorems	59
A.2. Products, compositions, and rescaling	64
A.3. Difference quotients	71
A.4. Spaces of sections of vector bundles	75
A.5. Some remarks on domains with cylindrical ends	79

In this appendix, we review some of the standard properties of Sobolev spaces, in particular using them to prove Propositions 2.2.1, 2.2.2 and 2.2.3 from §2.2, and elucidating the construction of Sobolev spaces of sections on vector bundles. A good reference for the necessary background material is [AF03].

A.1. Approximation, extension and embedding theorems

Unless otherwise noted, all functions in the following are assumed to be defined on a nonempty open subset

$$\mathcal{U} \subset \mathbb{R}^n$$

with its standard Lebesgue measure, and taking values in a finite-dimensional normed vector space that will usually not need to be specified, though occasionally we will assume it is \mathbb{R} or \mathbb{C} so that one can define products of functions. The domain \mathcal{U} will also sometimes have additional conditions specified such as boundedness or regularity at the boundary, though we will try not to add too many more restrictions than are really needed. The most useful assumption to impose on \mathcal{U} is known as the **strong local Lipschitz condition**: if \mathcal{U} is bounded, then it means simply that near every boundary point of \mathcal{U} , one can find smooth local coordinates in which \mathcal{U} looks like the region bounded by the graph of a Lipschitz-continuous function, and in this case we call \mathcal{U} a **bounded Lipschitz domain**. If \mathcal{U} is unbounded, then one needs to impose extra conditions guaranteeing e.g. uniformity of Lipschitz constants, and the precise definition becomes a bit lengthy (see [AF03, §4.9]). For our purposes, all we really need to know about the strong local Lipschitz condition is that that it is satisfied both by bounded Lipschitz domains and by relatively tame unbounded domains such as $(0, 1) \times (0, \infty) \subset \mathbb{R}^2$ which have smooth boundary with finitely many corners. We will repeatedly need to use the generalized version of **Hölder's inequality**, which states that for any finite collection of measurable

functions f_1, \dots, f_m ,

$$(A.1) \quad \left\| \prod_{i=1}^m |f_i| \right\|_{L^p} \leq \prod_{i=1}^m \|f_i\|_{L^{p_i}} \quad \text{for } 1 \leq p \leq p_1, \dots, p_m \leq \infty \text{ with } \frac{1}{p} = \sum_{i=1}^m \frac{1}{p_i}.$$

This is an easy corollary of the standard version,

$$\| |f| \cdot |g| \|_{L^1} \leq \|f\|_{L^p} \cdot \|g\|_{L^q} \quad \text{whenever } 1 \leq p, q \leq \infty \text{ and } 1 = \frac{1}{p} + \frac{1}{q}.$$

For an integer $k \geq 0$ and real number $p \in [1, \infty]$, we define $W^{k,p}(\mathcal{U})$ as in §2.2 to be the Banach space of all $f \in L^p(\mathcal{U})$ which have weak partial derivatives $\partial^\alpha f \in L^p(\mathcal{U})$ for all $|\alpha| \leq k$. For $p = 2$, these spaces are also often denoted by

$$H^k(\mathcal{U}) := W^{k,2}(\mathcal{U}),$$

and they admit Hilbert space structures with inner product

$$\langle f, g \rangle_{H^k} = \sum_{|\alpha| \leq k} \langle \partial^\alpha f, \partial^\alpha g \rangle_{L^2}.$$

We denote by

$$W_0^{k,p}(\mathcal{U}) \subset W^{k,p}(\mathcal{U}), \quad H_0^k(\mathcal{U}) \subset H^k(\mathcal{U})$$

the closed subspaces defined as the closures of $C_0^\infty(\mathcal{U})$ with respect to the relevant norms. Since $C_0^\infty(\mathcal{U})$ is dense in $L^p(\mathcal{U})$ for $1 \leq p < \infty$ (see e.g. [LL01, §2.19]), there is no difference between $W^{0,p}(\mathcal{U})$ and $W_0^{0,p}(\mathcal{U})$ for $p < \infty$, but in general $W_0^{k,p}(\mathcal{U}) \neq W^{k,p}(\mathcal{U})$ for $k \geq 1$, with a few notable exceptions such as the case $\mathcal{U} = \mathbb{R}^n$ (cf. Corollary A.1.2 below). Let

$$W_{\text{loc}}^{k,p}(\mathcal{U}) := \{ \text{functions } f \text{ on } \mathcal{U} \mid f \in W^{k,p}(\mathcal{V}) \text{ for all open subsets } \mathcal{V} \subset \mathcal{U} \\ \text{with compact closure } \overline{\mathcal{V}} \subset \mathcal{U} \},$$

and say that a sequence $f_j \in W_{\text{loc}}^{k,p}(\mathcal{U})$ converges in $W_{\text{loc}}^{k,p}$ to $f \in W_{\text{loc}}^{k,p}(\mathcal{U})$ if the restrictions to all precompact open subsets $\mathcal{V} \subset \overline{\mathcal{V}} \subset \mathcal{U}$ converge in $W^{k,p}(\mathcal{V})$. Recall that for $k \in \{0, 1, 2, \dots, \infty\}$, $C^k(\mathcal{U})$ denotes the space of functions on \mathcal{U} with continuous derivatives up to order k , while

$$C^k(\overline{\mathcal{U}}) \subset C^k(\mathcal{U})$$

is the space of $f \in C^k(\mathcal{U})$ such that for all $|\alpha| \leq k$, $\partial^\alpha f$ is bounded and uniformly continuous.

THEOREM A.1.1 ([AF03, §3.17, 3.22]). *For any open subset $\mathcal{U} \subset \mathbb{R}^n$, and any $k \geq 0$, $1 \leq p < \infty$, the subspace*

$$C^\infty(\mathcal{U}) \cap W^{k,p}(\mathcal{U}) \subset W^{k,p}(\mathcal{U})$$

is dense. Moreover, if $\mathcal{U} \subset \mathbb{R}^n$ satisfies the strong local Lipschitz condition, then the space

$$\left\{ f \in C^\infty(\mathcal{U}) \mid f = \tilde{f}|_{\mathcal{U}} \text{ for some } \tilde{f} \in C_0^\infty(\mathbb{R}^n) \right\}$$

is also dense in $W^{k,p}(\mathcal{U})$, so in particular,

$$C^\infty(\overline{\mathcal{U}}) \cap W^{k,p}(\mathcal{U}) \subset W^{k,p}(\mathcal{U})$$

is dense. \square

COROLLARY A.1.2. *The space $C_0^\infty(\mathbb{R}^n)$ is dense in $W^{k,p}(\mathbb{R}^n)$ for every $k \geq 0$ and $p \in [1, \infty)$. \square*

Here is another useful characterization of $W_0^{k,p}(\mathcal{U})$:

THEOREM A.1.3 ([AF03, §5.29]). *Assume $\mathcal{U} \subset \mathbb{R}^n$ is an open subset satisfying the strong local Lipschitz condition. Then a function $f \in W^{k,p}(\mathcal{U})$ belongs to $W_0^{k,p}(\mathcal{U})$ if and only if the function \tilde{f} on \mathbb{R}^n defined to match f on \mathcal{U} and 0 everywhere else belongs to $W^{k,p}(\mathbb{R}^n)$. \square*

While it is obvious from the definitions that functions in $W_0^{k,p}(\mathcal{U})$ always admit extensions of class $W^{k,p}$ over \mathbb{R}^n , this is much less obvious for functions in $W^{k,p}(\mathcal{U})$ in general, and it is not true without sufficient assumptions about the regularity of $\partial\mathcal{U}$. For our purposes it suffices to consider the following case.

THEOREM A.1.4 ([AF03, §5.22]). *Assume $\mathcal{U} \subset \mathbb{R}^n$ is a bounded open subset such that $\partial\overline{\mathcal{U}}$ is a submanifold of class C^m for some $m \in \{1, 2, 3, \dots, \infty\}$. Then there exists a linear operator E that maps functions defined almost everywhere on \mathcal{U} to functions defined almost everywhere on \mathbb{R}^n and has the following properties:*

- For every function f on \mathcal{U} , $Ef|_{\mathcal{U}} \equiv f$ almost everywhere;
- For every nonnegative integer $k \leq m$ and every $p \in [1, \infty)$, E defines a bounded linear operator $W^{k,p}(\mathcal{U}) \rightarrow W^{k,p}(\mathbb{R}^n)$. \square

COROLLARY A.1.5. *Suppose $\mathcal{U}, \mathcal{U}' \subset \mathbb{R}^n$ are open subsets such that \mathcal{U} has compact closure contained in \mathcal{U}' . If \mathcal{U} satisfies the hypothesis of Theorem A.1.4, then the resulting extension operator E can be chosen such that it maps each $W^{k,p}(\mathcal{U})$ for $k \leq m$ and $1 \leq p < \infty$ into $W_0^{k,p}(\mathcal{U}')$.*

PROOF. Choose a smooth function $\rho : \mathcal{U}' \rightarrow [0, 1]$ that has compact support and equals 1 on $\overline{\mathcal{U}}$, then replace the operator E given by Theorem A.1.4 with the operator $f \mapsto \rho \cdot Ef$. \square

To state the Sobolev embedding theorem in its proper generality, recall that for $0 < \alpha \leq 1$, the **Hölder seminorm** of a function f on \mathcal{U} is defined by

$$|f|_{C^\alpha} := |f|_{C^\alpha(\mathcal{U})} := \sup_{x \neq y \in \mathcal{U}} \frac{|f(x) - f(y)|}{|x - y|^\alpha},$$

and $C^{k,\alpha}(\mathcal{U})$ is then defined as the Banach space of functions $f \in C^k(\overline{\mathcal{U}})$ for which the norm

$$\|f\|_{C^{k,\alpha}} := \|f\|_{C^k} + \max_{|\beta|=k} |\partial^\beta f|_{C^\alpha}$$

is finite. In reading the following statement, it is important to remember that elements of $W^{k,p}(\mathcal{U})$ are technically not functions, but rather *equivalence classes* of functions defined almost everywhere. Thus when we say e.g. that there is an inclusion $W^{k,p}(\mathcal{U}) \hookrightarrow C^{m,\alpha}(\mathcal{U})$, the literal meaning is that for every function f representing an element of $W^{k,p}(\mathcal{U})$, one can change the values of f in a unique way

on some set of measure zero in \mathcal{U} so that after this change, $f \in C^{m,\alpha}(\mathcal{U})$. Continuity of the inclusion means that there is a bound of the form

$$\|f\|_{C^{m,\alpha}} \leq c \|f\|_{W^{k,p}}$$

for all $f \in W^{k,p}(\mathcal{U})$, where $c > 0$ is a constant which may in general depend on m , α , k , p and \mathcal{U} , but not on f .

THEOREM A.1.6 ([AF03, §4.12]). *Assume $\mathcal{U} \subset \mathbb{R}^n$ is an open subset satisfying the strong local Lipschitz condition, $k \geq 1$ is an integer and $1 \leq p < \infty$.*

(1) *If $0 < k - n/p \leq 1$, then there exist continuous inclusions*

$$\begin{aligned} W^{k,p}(\mathcal{U}) &\hookrightarrow C^{0,\alpha}(\mathcal{U}) && \text{for each } \alpha \in (0,1) \text{ with } \alpha \leq k - n/p, \\ W^{k,p}(\mathcal{U}) &\hookrightarrow L^q(\mathcal{U}) && \text{for each } q \in [p, \infty]. \end{aligned}$$

(2) *If $kp < n$ and $p^* > p$ is defined by the condition*

$$\frac{1}{p^*} = \frac{1}{p} - \frac{k}{n},$$

then there exist continuous inclusions

$$W^{k,p}(\mathcal{U}) \hookrightarrow L^q(\mathcal{U}), \quad \text{for each } q \in [p, p^*].$$

(3) *If $kp = n$, then there exist continuous inclusions*

$$W^{k,p}(\mathcal{U}) \hookrightarrow L^q(\mathcal{U}), \quad \text{for each } q \in [p, \infty).$$

Moreover, the spaces $W_0^{k,p}(\mathcal{U})$ admit similar inclusions under no assumption on the open subset $\mathcal{U} \subset \mathbb{R}^n$. \square

Under the same assumption on the domain \mathcal{U} , one can apply Theorem A.1.6 to successive derivatives of functions in $W^{k,p}(\mathcal{U})$ and thus obtain the following inclusions for any integer $d \geq 0$:

$$(A.2) \quad W^{k+d,p}(\mathcal{U}) \hookrightarrow C^{d,\alpha}(\mathcal{U}) \quad \text{if } 0 < k - n/p \leq 1, 0 < \alpha < 1 \text{ and } \alpha \leq k - n/p,$$

$$(A.3) \quad W^{k+d,p}(\mathcal{U}) \hookrightarrow W^{d,q}(\mathcal{U}) \quad \text{if } kp > n \text{ and } p \leq q \leq \infty,$$

$$(A.4) \quad W^{k+d,p}(\mathcal{U}) \hookrightarrow W^{d,q}(\mathcal{U}) \quad \text{if } kp < n \text{ and } p \leq q \leq p^*, \text{ with } \frac{1}{p^*} = \frac{1}{p} - \frac{k}{n},$$

$$(A.5) \quad W^{k+d,p}(\mathcal{U}) \hookrightarrow W^{d,q}(\mathcal{U}) \quad \text{if } kp = n \text{ and } p \leq q < \infty.$$

REMARK A.1.7. The embedding theorem suggests that one should intuitively think of $W^{k,p}(\mathcal{U})$ as consisting of functions with “ $k - n/p$ continuous derivatives,” where the number $k - n/p$ may in general be a non-integer and/or negative. This provides a useful mnemonic for results about embeddings of one Sobolev space into another, such as the following.

COROLLARY A.1.8. *Assume $\mathcal{U} \subset \mathbb{R}^n$ is an open subset satisfying the strong local Lipschitz condition, $1 \leq p, q < \infty$, and $k, m \geq 0$ are integers satisfying*

$$k \geq m, \quad p \leq q, \quad \text{and} \quad k - \frac{n}{p} \geq m - \frac{n}{q}.$$

Then there exists a continuous inclusion $W^{k,p}(\mathcal{U}) \hookrightarrow W^{m,q}(\mathcal{U})$. \square

EXERCISE A.1.9. Derive Corollary A.1.8 from Theorem A.1.6 by checking that under the stated conditions, there is a continuous inclusion $W^{k-m,p}(\mathcal{U}) \hookrightarrow L^q(\mathcal{U})$. Show also that the hypothesis $p \leq q$ is unnecessary if $\mathcal{U} \subset \mathbb{R}^n$ has finite measure.

By the Arzelà-Ascoli theorem, the natural inclusion

$$C^{k,\alpha'}(\mathcal{U}) \hookrightarrow C^{k,\alpha}(\mathcal{U})$$

for $\alpha < \alpha'$ is a compact operator whenever $\mathcal{U} \subset \mathbb{R}^n$ is bounded. It follows that if $\mathcal{U} \subset \mathbb{R}^n$ in (A.2) is bounded and α is *strictly* less than the extremal value $k - n/p$, then the inclusion (A.2) is also compact. A similar statement holds for the inclusion (A.4) when $p \leq q < p^*$, and this is known as the **Rellich-Kondrachov compactness theorem**. We summarize these as follows:

THEOREM A.1.10 ([AF03, §6.3]). *Assume $\mathcal{U} \subset \mathbb{R}^n$ is a bounded Lipschitz domain, $k \geq 1$ and $d \geq 0$ are integers and $1 \leq p < \infty$.*

(1) *If $kp > n$ and $k - n/p < 1$, then the inclusions*

$$\begin{aligned} W^{k+d,p}(\mathcal{U}) &\hookrightarrow C^{d,\alpha}(\mathcal{U}) && \text{for } \alpha \in (0, k - n/p), \\ W^{k+d,p}(\mathcal{U}) &\hookrightarrow W^{d,q}(\mathcal{U}) && \text{for } q \in [p, \infty) \end{aligned}$$

are compact.

(2) *If $kp \leq n$ and $p^* \in (p, \infty]$ is defined by the condition $1/p^* = 1/p - k/n$, then the inclusions*

$$W^{k+d,p}(\mathcal{U}) \hookrightarrow W^{d,q}(\mathcal{U}) \quad \text{for } q \in [p, p^*)$$

are compact.

In particular, the continuous inclusion $W^{k,p}(\mathcal{U}) \hookrightarrow W^{m,q}(\mathcal{U})$ in Corollary A.1.8 is compact whenever the inequality $k - n/p \geq m - n/q$ is strict. \square

On connected 1-dimensional domains $\mathcal{U} \subset \mathbb{R}$, the spaces $W^{1,p}(\mathcal{U})$ admit an alternative characterization in terms of classical derivatives defined almost everywhere:

PROPOSITION A.1.11. *For $-\infty < a < b < \infty$, every absolutely continuous function on $[a, b]$ belongs to $W^{1,1}((a, b))$ and has a weak derivative that is equal to its classical derivative almost everywhere. Conversely, every function in $W^{1,1}((a, b))$ is equal almost everywhere to an absolutely continuous function defined on $[a, b]$.*

PROOF. Let us denote the classical derivative of a function f by f'_c and the weak derivative by f'_w whenever there is danger of confusion. If f is absolutely continuous on $[a, b]$, then for every test function $\varphi \in C_0^\infty((a, b))$, $f\varphi$ defines an absolutely continuous function on $[a, b]$ that vanishes at the end points, so the fundamental theorem of calculus implies $\int_{[a,b]} (f\varphi)'_c = \int_{[a,b]} f'_c \varphi + \int_{[a,b]} f \varphi' = 0$, proving that the almost everywhere defined function $f'_c \in L^1([a, b])$ is also the weak derivative f'_w , and thus $f \in W^{1,1}((a, b))$.

Conversely, suppose $f \in W^{1,1}((a, b))$, so it has a weak derivative $f'_w \in L^1((a, b))$. We can then define an absolutely continuous function g on $[a, b]$ by $g(x) := \int_a^x f'_w$, which is differentiable almost everywhere and satisfies $g'_c = f'_w$. By the argument of the previous paragraph, g'_c is also a weak derivative g'_w , thus $g - f$ is a function on

(a, b) with vanishing weak derivative, implying via [LL01, Theorem 6.11] that $g - f$ is equal almost everywhere to a constant. \square

COROLLARY A.1.12. *For $-\infty < a < b < \infty$ and $1 \leq p \leq \infty$, $W^{1,p}((a, b))$ has a canonical identification with the space of absolutely continuous functions on $[a, b]$ whose classical derivatives belong to $L^p([a, b])$. \square*

A.2. Products, compositions, and rescaling

We now restate and prove Propositions 2.2.1, 2.2.2 and 2.2.3 from §2.2. These are all corollaries of the Sobolev embedding theorem, so in particular they hold for the same class of domains $\mathcal{U} \subset \mathbb{R}^n$, and the restrictions on \mathcal{U} can be dropped at the cost of replacing each space $W^{k,p}$ by $W_0^{k,p}$.

We begin by generalizing Prop. 2.2.1, hence we consider Sobolev spaces of functions valued in \mathbb{R} or \mathbb{C} so that pointwise products of functions are well defined almost everywhere. We say that there is a **continuous product map**,

$$W^{k_1, p_1}(\mathcal{U}) \times \dots \times W^{k_m, p_m}(\mathcal{U}) \rightarrow W^{k, p}(\mathcal{U}),$$

or a continuous product **pairing** in the case $m = 2$, if for every set of functions $f_i \in W^{k_i, p_i}(\mathcal{U})$ with $i = 1, \dots, m$, the pointwise product function $f_1 \cdot \dots \cdot f_m$ is in $W^{k, p}(\mathcal{U})$ and there is an estimate of the form

$$\|f_1 \cdot \dots \cdot f_m\|_{W^{k, p}} \leq c \|f_1\|_{W^{k_1, p_1}} \cdot \dots \cdot \|f_m\|_{W^{k_m, p_m}}$$

for some constant $c > 0$ not depending on f_1, \dots, f_m . The case $m = 2$, $k_1 = k_2 = k$ and $p_1 = p_2 = p$ is especially interesting, as the space $W^{k, p}(\mathcal{U})$ is then a **Banach algebra**. More generally, one can ask under what circumstances multiplication by functions of class $W^{k, p}$ defines a bounded linear operator on functions of class $W^{m, q}$. A hint about this comes from the world of classically differentiable functions: multiplication by C^k -smooth functions defines a continuous map $C^m \rightarrow C^m$ if and only if $k \geq m$. The corresponding answer in Sobolev spaces turns out to be that functions of class $W^{k, p}$ need to have strictly more than zero derivatives in the sense of Remark A.1.7, and at least as many derivatives as functions of class $W^{m, q}$.

THEOREM A.2.1. *Assume $\mathcal{U} \subset \mathbb{R}^n$ is an open subset satisfying the strong local Lipschitz condition, $1 \leq p, q < \infty$, and $k, m \geq 0$ are integers satisfying*

$$k \geq m, \quad kp > n, \quad \text{and} \quad k - \frac{n}{p} \geq m - \frac{n}{q}.$$

Then there exists a continuous product pairing

$$W^{k, p}(\mathcal{U}, \mathbb{C}) \times W^{m, q}(\mathcal{U}, \mathbb{C}) \rightarrow W^{m, q}(\mathcal{U}, \mathbb{C}) : (f, g) \mapsto fg.$$

The following preparatory lemma will be useful both for proving the product estimate and for further results below. It is an easy consequence of Theorem A.1.6 and Hölder's inequality.

LEMMA A.2.2. *Assume $\mathcal{U} \subset \mathbb{R}^n$ is an open subset satisfying the strong local Lipschitz condition, $m \geq 2$ is an integer, and we are given positive numbers*

$p_1, \dots, p_m \geq 1$ and integers $k_1, \dots, k_m \geq 0$. Let $I := \{i \in \{1, \dots, m\} \mid k_i p_i \leq n\}$. Then for any $q \geq 1$ satisfying

$$\sum_{i \in I} \left(\frac{1}{p_i} - \frac{k_i}{n} \right) < \frac{1}{q} \leq \sum_{i=1}^m \frac{1}{p_i},$$

there is a continuous product map

$$W^{k_1, p_1}(\mathcal{U}) \times \dots \times W^{k_m, p_m}(\mathcal{U}) \rightarrow L^q(\mathcal{U}).$$

PROOF. By the generalized Hölder inequality (A.1), it suffices to show that for any $q \geq 1$ in the stated range, one can find numbers $q_1, \dots, q_m \in [q, \infty]$ satisfying $1/q = 1/q_1 + \dots + 1/q_m$ for which Theorem A.1.6 provides continuous inclusions

$$W^{k_i, p_i}(\mathcal{U}) \hookrightarrow L^{q_i}(\mathcal{U})$$

for each $i = 1, \dots, m$. Whenever $k_i p_i > n$, this inclusion is valid with q_i chosen freely from the interval $[p_i, \infty]$, so $1/q_i$ can then take any value subject to the constraint

$$0 \leq \frac{1}{q_i} \leq \frac{1}{p_i}.$$

If on the other hand $k_i p_i \leq n$, then we can arrange $1/q_i$ to take any value in the range

$$\frac{1}{p_i} - \frac{k_i}{n} < \frac{1}{q_i} \leq \frac{1}{p_i}.$$

Adding these up, the range of values for $\sum_i \frac{1}{q_i}$ that we can achieve in this way covers the stated interval. \square

PROOF OF THEOREM A.2.1. By density of smooth functions, it suffices to prove that an estimate of the form

$$\|fg\|_{W^{m,q}} \leq c \|f\|_{W^{k,p}} \|g\|_{W^{m,q}}$$

holds for all $f \in C^\infty(\mathcal{U}) \cap W^{k,p}(\mathcal{U})$ and $g \in C^\infty(\mathcal{U}) \cap W^{m,q}(\mathcal{U})$. Equivalently, we need to show that for all f and g of this type and every multiindex α of degree $|\alpha| \leq m$, there is a constant $c > 0$ independent of f and g such that

$$\|\partial^\alpha(fg)\|_{L^q} \leq c \|f\|_{W^{k,p}} \|g\|_{W^{m,q}}.$$

Since f and g are smooth, we are free to use the product rule in computing $\partial^\alpha(fg)$, which will then be a linear combination of terms of the form $\partial^\beta f \cdot \partial^\gamma g$ where $|\alpha| = |\beta| + |\gamma|$, hence we have reduced the problem to proving a bound

$$\|\partial^\beta f \cdot \partial^\gamma g\|_{L^q} \leq c \|f\|_{W^{k,p}} \|g\|_{W^{m,q}}$$

for every pair of multiindices β, γ with $|\beta| + |\gamma| \leq m$. Since $\partial^\beta f \in W^{k-|\beta|,p}(\mathcal{U})$ and $\partial^\gamma g \in W^{m-|\gamma|,q}(\mathcal{U})$, the result follows if we can assume that for every pair of integers $a, b \geq 0$ satisfying $a + b \leq m$, there exists a continuous product pairing

$$(A.6) \quad W^{k-a,p}(\mathcal{U}) \times W^{m-b,q}(\mathcal{U}) \rightarrow L^q(\mathcal{U}).$$

If $(k-a)p > n$, then $W^{k-a,p} \hookrightarrow L^\infty$ and (A.6) is immediate since $W^{m-b,q} \hookrightarrow L^q(\mathcal{U})$. For the remaining cases, we shall apply Lemma A.2.2, noting that the condition $1/q \leq 1/p + 1/q$ is trivially satisfied.

If $(m-b)q > n$ but $(k-a)p \leq n$, then the hypotheses of the lemma are satisfied if and only if

$$\frac{1}{p} - \frac{k-a}{n} < \frac{1}{q}.$$

Since $\frac{1}{p} - \frac{k}{n} \leq \frac{1}{q} - \frac{m}{n}$ by assumption, we have

$$\frac{1}{p} - \frac{k-a}{n} = \frac{1}{p} - \frac{k}{n} + \frac{a}{n} \leq \frac{1}{q} - \frac{m}{n} + \frac{a}{n} \leq \frac{1}{q}$$

since $a \leq m$, and equality holds only if $a = m$, $b = 0$ and $k - n/p = m - n/q$, which implies $mq > n$. In this case $W^{m-b,q} = W^{m,q} \hookrightarrow L^\infty$, and the pairing (A.6) follows because $W^{k-a,p} = W^{k-m,p}$ embeds continuously into L^q : the latter follows from Theorem A.1.6 since $\frac{1}{p} - \frac{k-m}{n} = \frac{1}{q}$.

Finally, when $(k-a)p \leq n$ and $(m-b)q \leq n$, the hypotheses of the lemma are satisfied since

$$\left(\frac{1}{p} - \frac{k-a}{n}\right) + \left(\frac{1}{q} - \frac{m-b}{n}\right) \leq \frac{1}{p} - \frac{k}{n} + \frac{1}{q} - \frac{m}{n} + \frac{m}{n} = \left(\frac{1}{p} - \frac{k}{n}\right) + \frac{1}{q} < \frac{1}{q},$$

where we've used the assumption $kp > n$ and the fact that $a + b \leq m$. \square

REMARK A.2.3. A much simpler argument shows similarly that for any open domain $\mathcal{U} \subset \mathbb{R}^n$, any integer $k \geq 1$ and any $p \in [1, \infty)$, there is a continuous product pairing

$$C^k(\overline{\mathcal{U}}, \mathbb{C}) \times W^{k,p}(\mathcal{U}, \mathbb{C}) \times W^{k,p}(\mathcal{U}, \mathbb{C}).$$

As in Theorem A.2.1, this follows from the density of $C^\infty \cap W^{k,p} \subset W^{k,p}$ after showing that all $f \in C^k(\overline{\mathcal{U}})$ and $g \in C^\infty(\mathcal{U}) \cap W^{k,p}(\mathcal{U})$ satisfy an estimate of the form $\|fg\|_{W^{k,p}} \leq c\|f\|_{C^k}\|g\|_{W^{k,p}}$. The latter follows easily from the definition of the $W^{k,p}$ -norm.

In general it is not straightforward to say when the usual product rule $\partial_i(fg) = \partial_i f \cdot g + f \cdot \partial_i g$ does or does not hold in the sense of weak derivatives. If g and $\partial_i g$ are locally integrable and f is smooth, then there is no trouble: the formula can be derived in this case directly from the definition of weak derivatives, using the observation that for any test function $\varphi \in C_0^\infty(\mathcal{U})$, φf is also in $C_0^\infty(\mathcal{U})$ and satisfies the product rule. If on the other hand f and g are not continuous but have well-defined weak derivatives and a locally integrable product, then there is no guarantee in general that any of $\partial_i(fg)$, $\partial_i f \cdot g$ or $f \cdot \partial_i g$ should be well-defined locally integrable functions. Theorem A.2.1 provides a means of resolving this question whenever f and g belong to suitable Sobolev spaces.

PROPOSITION A.2.4. *Suppose k, m, p, q and $\mathcal{U} \subset \mathbb{R}^n$ satisfy the same conditions as in Theorem A.2.1, and $m \geq 1$. Then for every $f \in W^{k,p}(\mathcal{U}, \mathbb{C})$ and $g \in W^{m,q}(\mathcal{U}, \mathbb{C})$, the weak partial derivatives of $fg \in W^{m,q}(\mathcal{U}, \mathbb{C})$ are given almost everywhere by the usual Leibniz rule $\partial_i(fg) = \partial_i f \cdot g + f \cdot \partial_i g$.*

PROOF. Choose sequences of smooth functions f_j, g_j with $f_j \rightarrow f$ in $W^{k,p}$ and $g_j \rightarrow g$ in $W^{m,q}$. Then since $k \geq m \geq 1$, there is also L^p -convergence $\partial_i f_j \rightarrow \partial_i f$ and L^q -convergence $\partial_i g_j \rightarrow \partial_i g$, so after restricting to a subsequence, we may assume that

all four of the sequences f_j , $\partial_i f_j$, g_j and $\partial_i g_j$ converge pointwise almost everywhere. The continuity of the product pairing $W^{k,p} \times W^{m,q} \rightarrow W^{m,q}$ now implies $W^{m,q}$ -convergence $f_j g_j \rightarrow fg$ and thus L^q -convergence

$$\partial_i(f_j g_j) = \partial_i f_j \cdot g_j + f_j \cdot \partial_i g_j \rightarrow \partial_i(fg).$$

The result follows since $\partial_i f_j \cdot g_j + f_j \cdot \partial_i g_j$ also converges pointwise almost everywhere to $\partial_i f \cdot g + f \cdot \partial_i g$. \square

REMARK A.2.5. A slight simplification of the same argument as in Proposition A.2.4 shows that the product rule also holds (without any assumption on the open domain $\mathcal{U} \subset \mathbb{R}^n$) for $f \in C^m(\overline{\mathcal{U}}, \mathbb{C})$ and $g \in W^{m,p}(\mathcal{U}, \mathbb{C})$ for any $p \in [1, \infty)$ if $m \geq 1$. The key facts here are the continuity of the product pairing $C^m \times W^{m,p} \rightarrow W^{m,p}$ and the density of C^1 in $W^{m,p}$, so that f and g can be approximated by pairs for which the classical product rule holds. Both results can also be extended in a similar manner to prove the expected formula for $\partial^\alpha(fg)$ for any multiindex α of order $|\alpha| \leq m$.

The next result generalizes Proposition 2.2.2 and concerns the following question: if $f : \mathcal{U} \rightarrow \mathbb{R}^m$ is a function of class $W^{k,p}$ whose graph lies in some open subset $\mathcal{V} \subset \mathcal{U} \times \mathbb{R}^m$, and $\Psi : \mathcal{V} \rightarrow \mathbb{R}^N$ is another function, under what conditions can we conclude that the function

$$\mathcal{U} \rightarrow \mathbb{R}^N : x \mapsto \Psi(x, f(x))$$

is in $W^{k,p}(\mathcal{U}, \mathbb{R}^N)$? We will abbreviate this function in the following by $\Psi \circ (\text{Id} \times f)$, and we would also like to know whether it depends continuously (in the $W^{k,p}$ -topology) on f and Ψ . The following theorem is stated rather generally, but on first reading you may prefer to assume $\mathcal{U} \subset \mathbb{R}^n$ is bounded, in which case some of the hypotheses become vacuous. We will say that an open subset $\mathcal{V} \subset \mathcal{U} \times \mathbb{R}^m$ is a **star-shaped neighborhood of $f : \mathcal{U} \rightarrow \mathbb{R}^m$** if it contains the graph of f and

$$(x, v) \in \mathcal{V} \quad \Rightarrow \quad (x, tv + (1-t)f(x)) \in \mathcal{V} \text{ for all } t \in [0, 1].$$

THEOREM A.2.6. *Assume $\mathcal{U} \subset \mathbb{R}^n$ is an open subset satisfying the strong local Lipschitz condition, $p \in [1, \infty)$ and $k \in \mathbb{N}$ satisfy $kp > n$, and $\mathcal{V} \subset \mathcal{U} \times \mathbb{R}^m$ is a star-shaped neighborhood of some function $f_0 \in W^{k,p}(\mathcal{U}, \mathbb{R}^m)$. Assume also $\mathcal{O}^{k,p}(\mathcal{U}; \mathcal{V}) \subset W^{k,p}(\mathcal{U}, \mathbb{R}^m)$ is an open neighborhood of f_0 such that*

$$(x, f(x)) \in \mathcal{V} \quad \text{for all } x \in \mathcal{U} \text{ and } f \in \mathcal{O}^{k,p}(\mathcal{U}; \mathcal{V}),$$

and $C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N) \subset C^k(\overline{\mathcal{V}}, \mathbb{R}^N)$ is a closed linear subspace such that all $\Psi \in C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N)$ have the following properties:¹

- (1) *There exists a bounded subset $\mathcal{K} \subset \mathcal{U}$ such that $\Psi(x, v)$ is independent of x for all $x \in \mathcal{U} \setminus \mathcal{K}$;*
- (2) *$\Psi \circ (\text{Id} \times f_0) \in L^p(\mathcal{U}, \mathbb{R}^N)$.*

Then there is a well-defined and continuous map

$$\begin{aligned} \mathcal{O}^{k,p}(\mathcal{U}; \mathcal{V}) &\xrightarrow{T} \mathcal{L}(C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N), W^{k,p}(\mathcal{U}, \mathbb{R}^N)), \\ T(f)\Psi &:= \Psi \circ (\text{Id} \times f), \end{aligned}$$

¹Both of the conditions on $\Psi \in C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N)$ are vacuous if $\mathcal{U} \subset \mathbb{R}^n$ is bounded.

so in particular the map

$$C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N) \times \mathcal{O}^{k,p}(\mathcal{U}; \mathcal{V}) \rightarrow W^{k,p}(\mathcal{U}, \mathbb{R}^N) : (\Psi, f) \mapsto \Psi \circ (\text{Id} \times f),$$

is well defined and continuous. Moreover, for each $\Psi \in C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N)$ and $f \in W^{k,p}(\mathcal{U}, \mathbb{R}^N)$, the weak partial derivatives of $\Psi \circ (\text{Id} \times f)$ are given almost everywhere by the classical formula

$$\partial_j [\Psi \circ (\text{Id} \times f)](x) = \partial_j \Psi(x, f(x)) + D_2 \Psi(x, f(x)) \partial_j f(x),$$

where $\partial_j \Psi$ denotes the partial derivative of $\Psi(x, v)$ with respect to the j th coordinate in $x \in \mathbb{R}^n$, and $D_2 \Psi$ is its differential with respect to $v \in \mathbb{R}^m$.

PROOF. We will show first that if $f \in \mathcal{O}^{k,p}(\mathcal{U}; \mathcal{V})$ is smooth, then $\Psi \circ (\text{Id} \times f)$ belongs to $W^{k,p}(\mathcal{U}, \mathbb{R}^N)$ for every $\Psi \in C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N)$. Since \mathcal{V} is a star-shaped neighborhood of f_0 , we have

$$\begin{aligned} |\Psi(x, f(x)) - \Psi(x, f_0(x))| &= \left| \int_0^1 \frac{d}{dt} \Psi(x, tf(x) + (1-t)f_0(x)) dt \right| \\ &\leq \left(\int_0^1 |D_2 \Psi(x, tf(x) + (1-t)f_0(x))| dt \right) \cdot |f(x) - f_0(x)| \\ &\leq \|\Psi\|_{C^1(\mathcal{V})} \cdot |f(x) - f_0(x)| \end{aligned}$$

for all $x \in \mathcal{U}$, implying

$$(A.7) \quad \|\Psi \circ (\text{Id} \times f) - \Psi \circ (\text{Id} \times f_0)\|_{L^p} \leq \|\Psi\|_{C^1(\mathcal{V})} \cdot \|f - f_0\|_{L^p},$$

hence $\Psi \circ (\text{Id} \times f) \in L^p(\mathcal{U}, \mathbb{R}^N)$.

For $\ell = 1, \dots, k$, we can regard the ℓ th derivative of Ψ with respect to variables in \mathbb{R}^m as a bounded and uniformly continuous map from \mathcal{V} into the vector space of symmetric ℓ -multilinear maps from \mathbb{R}^m to \mathbb{R}^N , denoting this by

$$D_2^\ell \Psi : \mathcal{V} \rightarrow \text{Hom}((\mathbb{R}^m)^{\otimes \ell}, \mathbb{R}^N).$$

Denote the partial derivatives with respect to variables in $\mathcal{U} \subset \mathbb{R}^n$ by

$$D_1^\beta \Psi : \mathcal{V} \rightarrow \mathbb{R}^N,$$

where β is a multiindex in n variables. Now for any multiindex α with $|\alpha| \leq k$, the derivative $\partial^\alpha (\Psi \circ (\text{Id} \times f))$ is a linear combination of product functions of the form

$$(A.8) \quad (D_1^\gamma D_2^\ell \Psi \circ (\text{Id} \times f))(\partial^{\beta_1} f, \dots, \partial^{\beta_\ell} f) : \mathcal{U} \rightarrow \mathbb{R}^N,$$

where $\ell + |\gamma| \in \{1, \dots, |\alpha|\}$ and $|\beta_1| + \dots + |\beta_\ell| = |\alpha| - |\gamma|$. If $\ell = 0$ but $|\gamma| > 0$, then this expression is clearly in $L^p(\mathcal{U}, \mathbb{R}^N)$ since it is continuous and $D_1^\gamma \Psi(x, v) = 0$ for $x \in \mathcal{U} \setminus \mathcal{K}$, where \mathcal{K} is bounded. For $\ell \geq 1$, it satisfies

$$(A.9) \quad \|(D_1^\gamma D_2^\ell \Psi \circ (\text{Id} \times f))(\partial^{\beta_1} f, \dots, \partial^{\beta_\ell} f)\|_{L^p(\mathcal{U})} \leq \|D_1^\gamma D_2^\ell \Psi\|_{C^0(\mathcal{V})} \cdot \left\| \prod_{j=1}^{\ell} |\partial^{\beta_j} f| \right\|_{L^p(\mathcal{U})}$$

if the product on the right hand side has finite L^p -norm. The latter is trivially true if $\ell = 1$. To deal with the $\ell \geq 2$ case, note that $\partial^{\beta_j} f \in W^{k-|\beta_j|,p}(\mathcal{U})$ for each $j = 1, \dots, \ell$, so the necessary bound will follow from the existence of a continuous product map

$$W^{k-m_1,p}(\mathcal{U}) \times \dots \times W^{k-m_\ell,p}(\mathcal{U}) \rightarrow L^p(\mathcal{U})$$

for $m_j := |\beta_j|$, and we claim that such a product map does exist whenever $kp > n$ and $m_1, \dots, m_\ell \geq 0$ are integers satisfying $m_1 + \dots + m_\ell \leq k$. To see this, note first that since $W^{k-m_j, p} \hookrightarrow L^\infty$ whenever $(k - m_j)p > n$, it suffices to prove the claim under the assumption that $(k - m_j)p \leq n$ for every $j = 1, \dots, \ell$. In this case, Lemma A.2.2 provides the desired product map if the condition

$$\sum_{j=1}^{\ell} \left(\frac{1}{p} - \frac{k - m_j}{n} \right) < \frac{1}{p} \leq \sum_{j=1}^{\ell} \frac{1}{p}$$

is satisfied. And it is: using $kp > n$, $\ell \geq 2$ and $m_1 + \dots + m_\ell \leq k$, we find

$$\begin{aligned} \sum_{j=1}^{\ell} \left(\frac{1}{p} - \frac{k - m_j}{n} \right) &= \ell \left(\frac{1}{p} - \frac{k}{n} \right) + \frac{m_1 + \dots + m_\ell}{n} \\ &\leq \frac{1}{p} + (\ell - 1) \left(\frac{1}{p} - \frac{k}{n} \right) < \frac{1}{p}. \end{aligned}$$

This proves that $\Psi \circ (\text{Id} \times f) \in W^{k,p}(\mathcal{U}, \mathbb{R}^N)$.

An important detail in both of the estimates (A.7) and (A.9) is that on the right hand side, the term depending on Ψ is bounded by something linearly proportional to $\|\Psi\|_{C^k(\mathcal{V})}$, and the same is true of other estimates mentioned below that can be derived in a similar manner. We will not comment on this point any further, but it is the reason why rather than just proving that the map $(\Psi, f) \mapsto \Psi \circ (\text{Id} \times f)$ is continuous, we will obtain the stronger result that the map sending f to the linear operator $\Psi \mapsto \Psi \circ (\text{Id} \times f)$ is continuous with respect to the operator norm.

Next, suppose $f \in \mathcal{O}^{k,p}(\mathcal{U}; \mathcal{V})$ is not necessarily smooth but $f_i \in \mathcal{O}^{k,p}(\mathcal{U}; \mathcal{V})$ is a sequence of smooth functions converging to f in $W^{k,p}$, while $\Psi_i \in C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N)$ converges to $\Psi \in C_0^k(\overline{\mathcal{V}}, \mathbb{R}^N)$ in C^k . Then the same argument we used to estimate $\|\Psi \circ (\text{Id} \times f) - \Psi \circ (\text{Id} \times f_0)\|_{L^p}$ shows that $\Psi_i \circ (\text{Id} \times f_i) \rightarrow \Psi \circ (\text{Id} \times f)$ in L^p , and since f_i is also C^0 -convergent, the compactly supported functions $D_1^\gamma \Psi_i \circ (\text{Id} \times f_i)$ converge to $D_1^\gamma \Psi \circ (\text{Id} \times f)$ in L^p for each multiindex with $1 \leq |\gamma| \leq k$. For $\ell \geq 1$ and $|\gamma| + \ell \leq k$, $D_1^\gamma D_2^\ell \Psi_i \circ (\text{Id} \times f_i)$ converges to $D_1^\gamma D_2^\ell \Psi \circ (\text{Id} \times f)$ in $C^0(\overline{\mathcal{U}}, \mathbb{R}^N)$, and each of the derivatives $\partial^{\beta_j} f_i$ appearing in (A.8) also converges in $L^p(\mathcal{U})$. In light of the continuous product maps discussed above, it follows that each derivative $\partial^\alpha(\Psi_i \circ (\text{Id} \times f_i))$ for $|\alpha| \leq k$ is L^p -convergent, and its limit is necessarily (by Exercise A.2.7 below) the corresponding weak derivative $\partial^\alpha(\Psi \circ (\text{Id} \times f))$, hence $\Psi \circ (\text{Id} \times f) \in W^{k,p}(\mathcal{U}, \mathbb{R}^N)$ and $\Psi_i \circ (\text{Id} \times f_i) \xrightarrow{W^{k,p}} \Psi \circ (\text{Id} \times f)$. Since all sequences in this discussion can also be replaced with subsequences that are pointwise almost everywhere convergent, this also proves that the classical formula for $\partial^\alpha(\Psi_i \circ (\text{Id} \times f_i))$ for each $|\alpha| \leq k$ remains valid for computing the corresponding weak derivative $\partial^\alpha(\Psi \circ (\text{Id} \times f))$. With this understood, one can now repeat the arguments of this paragraph for an arbitrary $W^{k,p}$ -convergent sequence $f_i \rightarrow f$ without assuming the f_i are smooth, thus proving the continuity of the map $(\Psi, f) \mapsto \Psi \circ (\text{Id} \times f)$. \square

EXERCISE A.2.7. Show that if f_i is a sequence of smooth functions on an open set $\mathcal{U} \subset \mathbb{R}^n$ with $f_i \xrightarrow{L^p} f$ and $\partial^\alpha f_i \xrightarrow{L^p} g$ for some multiindex α and functions $f, g \in L^p(\mathcal{U})$, then $\partial^\alpha f = g$ in the sense of distributions.

The following result on coordinate transformations of the domain can be proved in an analogous way to Theorem A.2.6, though it is considerably easier since there is no need to worry about Sobolev product maps (and thus no need to assume $kp > n$ or impose regularity conditions on the domain).

THEOREM A.2.8 ([AF03, §3.41]). *Assume $k \in \mathbb{N}$, $1 \leq p \leq \infty$, and $\mathcal{U}, \mathcal{U}' \subset \mathbb{R}^n$ are open subsets with a C^k -smooth diffeomorphism $\varphi : \mathcal{U} \rightarrow \mathcal{U}'$ such that all derivatives of φ and φ^{-1} up to order k are bounded and uniformly continuous. Then there is a well-defined Banach space isomorphism*

$$W^{k,p}(\mathcal{U}') \rightarrow W^{k,p}(\mathcal{U}) : f \mapsto f \circ \varphi.$$

□

Next, we restate and prove Proposition 2.2.3. Denote by $\mathring{\mathbb{D}}^n$ and $\mathring{\mathbb{D}}_\epsilon^n(x_0)$ the open balls of radius 1 and ϵ about the origin and a point x_0 respectively in \mathbb{R}^n .

THEOREM A.2.9. *Assume $p \in [1, \infty)$ and $k \in \mathbb{N}$ satisfy $kp > n$, and for a given point $x_0 \in \mathring{\mathbb{D}}^n$ with $\epsilon_0 := \text{dist}(x_0, \partial\mathbb{D}^n)$, associate to each $f \in W^{k,p}(\mathring{\mathbb{D}}^n)$ and $\epsilon \in (0, \epsilon_0)$ the function $f_\epsilon \in W^{k,p}(\mathring{\mathbb{D}}^n)$ defined by*

$$f_\epsilon(x) := f(x_0 + \epsilon x).$$

Then for each $\alpha \in (0, 1)$ satisfying $\alpha \leq k - \frac{n}{p}$, there exists a constant $C > 0$ such that the estimate

$$\|f_\epsilon - f_\epsilon(0)\|_{W^{k,p}} \leq C\epsilon^\alpha \|f - f(x_0)\|_{W^{k,p}}$$

holds for all $f \in W^{k,p}(\mathring{\mathbb{D}}^n)$ and $\epsilon \in (0, \epsilon_0)$.

PROOF. To estimate $\|f_\epsilon - f_\epsilon(0)\|_{L^p}$, we use the fact that $f - f(x_0) \in W^{k,p}$ is Hölder continuous, i.e. Theorem A.1.6 embeds $W^{k,p}$ continuously into $C^{0,\alpha}$ for any $\alpha \in (0, 1)$ with $\alpha \leq k - n/p$, thus f satisfies

$$|f(x) - f(x_0)| \leq c \|f - f(x_0)\|_{W^{k,p}(\mathring{\mathbb{D}}^n)} \cdot |x - x_0|^\alpha \quad \text{for all } x \in \mathring{\mathbb{D}}_{\epsilon_0}^n(x_0)$$

for some constant $c > 0$. We therefore have

$$\begin{aligned} \|f_\epsilon - f_\epsilon(0)\|_{L^p}^p &= \int_{\mathring{\mathbb{D}}^n} |f(x_0 + \epsilon x) - f(x_0)|^p \leq c^p \|f - f(x_0)\|_{W^{k,p}}^p \int_{\mathring{\mathbb{D}}^n} |\epsilon x|^{\alpha p} \\ &= c^p \|f - f(x_0)\|_{W^{k,p}}^p \cdot \epsilon^{\alpha p} \int_{\mathring{\mathbb{D}}^n} |x|^{\alpha p} =: C^p \epsilon^{\alpha p} \|f - f(x_0)\|_{W^{k,p}}^p \end{aligned}$$

for a suitable constant $C > 0$, implying $\|f_\epsilon - f_\epsilon(0)\|_{L^p} \leq C\epsilon^\alpha \|f - f(x_0)\|_{W^{k,p}}$.

Next, consider a multiindex β of order $|\beta| = m \in \{1, \dots, k\}$. The functions $\partial^\beta(f - f(x_0)) = \partial^\beta f$ and $\partial^\beta(f_\epsilon - f_\epsilon(0)) = \partial^\beta f_\epsilon$ for each $\epsilon \in (0, \epsilon_0)$ are then in $W^{k-m,p}(\mathring{\mathbb{D}}^n)$, and we need to establish bounds on $\|\partial^\beta f_\epsilon\|_{L^p}$ in terms of the $W^{k,p}$ -norm of $f - f(x_0)$. If $m < k$, then Theorem A.1.6 gives a continuous inclusion

$$(A.10) \quad W^{k-m,p}(\mathring{\mathbb{D}}^n) \hookrightarrow L^q(\mathring{\mathbb{D}}^n)$$

for any $q \in [p, \infty)$ satisfying $1/q \geq 1/p - (k-m)/n$. The same is also trivially true in the case $m = k$, since q and p must then be equal. Notice that if $(k-m)p \geq n$,

then q is allowed to be arbitrarily large. We will therefore assume in general that (A.10) holds with $q \in [p, \infty)$ satisfying

$$\frac{1}{q} + \frac{1}{r} = \frac{1}{p},$$

where $r = \frac{n}{k-m} \in (0, \infty]$ if $(k-m)p < n$ and otherwise $r = p + \delta$ for some $\delta > 0$ which may be chosen arbitrarily small. Given this, we apply change of variables and Hölder's inequality to find

$$\begin{aligned} \|\partial^\beta f_\epsilon\|_{L^p(\mathbb{D}^n)}^p &= \epsilon^{mp} \int_{\mathbb{D}^n} |\partial^\beta f(x_0 + \epsilon x)|^p = \epsilon^{mp-n} \int_{\mathbb{D}_\epsilon^n(x_0)} |\partial^\beta f(x)|^p \\ &\leq \epsilon^{mp-n} \|\partial^\beta f\|_{L^q(\mathbb{D}_\epsilon^n)}^p \|1\|_{L^r(\mathbb{D}_\epsilon^n)}^p \\ &\leq \epsilon^{mp-n} [\text{Vol}(\mathbb{D}_\epsilon^n(x_0))]^{p/r} \|\partial^\beta f\|_{L^q(\mathbb{D}_\epsilon^n)}^p \\ &\leq c\epsilon^{mp-n} [\text{Vol}(\mathbb{D}_\epsilon^n(x_0))]^{p/r} \|\partial^\beta f\|_{W^{k-m,p}(\mathbb{D}_\epsilon^n)}^p \\ &\leq c\epsilon^{mp-n} [\text{Vol}(\mathbb{D}_\epsilon^n(x_0))]^{p/r} \|f - f(x_0)\|_{W^{k,p}(\mathbb{D}_\epsilon^n)}^p \end{aligned}$$

for some constant $c > 0$. Writing $\text{Vol}(\mathbb{D}_\epsilon^n(x_0)) = C\epsilon^n$ for a suitable constant $C > 0$, the exponent on ϵ in this expression becomes $mp - n + \frac{np}{r}$. If $(k-m)p < 0$, this is exactly $kp - n = (k - n/p)p$, and otherwise, taking $r - p > 0$ to be arbitrarily small makes it less than but arbitrarily close to mp . Since $\alpha \leq k - n/p$ and $\alpha < 1 \leq m$, we are now free to replace this exponent with αp and rewrite the established estimate as $\|\partial^\beta f_\epsilon\|_{L^p} \leq C\epsilon^\alpha \|f - f(x_0)\|_{W^{k,p}}$. \square

A.3. Difference quotients

If f is a function on \mathbb{R}^n , then for every $i = 1, \dots, n$ and $h \in \mathbb{R} \setminus \{0\}$, the **difference quotient**

$$D_i^h f(x_1, \dots, x_n) := \frac{f(x_1, \dots, x_{i-1}, x_i + h, x_{i+1}, \dots, x_n) - f(x_1, \dots, x_n)}{h}$$

defines a function $D_i^h f$ on \mathbb{R}^n . The **total difference quotient** of f is then the n -tuple of functions

$$D^h f := (D_1^h f, \dots, D_n^h f),$$

so for example if $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then $D^h f : \mathbb{R}^n \rightarrow \mathbb{R}^{mn}$. The transformation $f \mapsto D_i^h f$ is obviously linear for any fixed number h , and it satisfies a Leibniz rule

$$D_i^h(fg) = D_i^h f \cdot g + f \cdot D_i^h g$$

whenever pointwise products of f and g can be defined (e.g. if both are real or complex valued). It also commutes with differentiation

$$D_i^h(\partial_j f) = \partial_j(D_i^h f)$$

on any function f for which $\partial_j f$ can be defined (weakly or strongly). Clearly if $f \in W^{k,p}(\mathbb{R}^n)$, then $D^h f \in W^{k,p}(\mathbb{R}^n)$ for every $h \in \mathbb{R} \setminus \{0\}$, and if f is supported in an open subset $\mathcal{U} \subset \mathbb{R}^n$, then $D^h f$ is supported in an arbitrarily small neighborhood of $\overline{\mathcal{U}}$ for sufficiently small $|h|$. Moreover, if f is a function defined only on $\mathcal{U} \subset \mathbb{R}^n$,

then on any open subset $\mathcal{V} \subset \mathcal{U}$ with compact closure in \mathcal{U} , $D^h f$ can be defined on \mathcal{V} for any $h \in \mathbb{R} \setminus \{0\}$ satisfying

$$|h| < \text{dist}(\mathcal{V}, \mathbb{R}^n \setminus \mathcal{U}) := \inf \{ |x - y| \mid x \in \mathcal{V} \text{ and } y \in \mathbb{R}^n \setminus \mathcal{U} \}.$$

The following result about difference quotients is useful for proving local regularity of solutions to PDEs, as in §2.4.

THEOREM A.3.1. *Assume $\mathcal{V} \subset \mathcal{U} \subset \mathbb{R}^n$ are open subsets with \mathcal{V} having compact closure contained in \mathcal{U} , $1 \leq p < \infty$, and $k \in \mathbb{N}$.*

(1) *If $f \in W^{k,p}(\mathcal{U})$, then $D^h f$ converges to ∇f in $W^{k-1,p}$ on \mathcal{V} as $h \rightarrow 0$, and*

$$\|D^h f\|_{W^{k-1,p}(\mathcal{V})} \leq \|\nabla f\|_{W^{k-1,p}(\mathcal{U})}$$

for all $h \neq 0$ with $|h| < \text{dist}(\mathcal{V}, \mathbb{R}^n \setminus \mathcal{U})$.

(2) *Suppose $p > 1$, $f \in W^{k-1,p}(\mathcal{U})$ and the difference quotients $D^h f$ satisfy a uniform bound*

$$\|D^h f\|_{W^{k-1,p}(\mathcal{V})} \leq C$$

for all $h \neq 0$ with $|h|$ sufficiently small. Then $f|_{\mathcal{V}} \in W^{k,p}(\mathcal{V})$ and its first derivative satisfies $\|\nabla f\|_{W^{k-1,p}(\mathcal{V})} \leq m_{k,p} C$, where $m_{k,p} \in \mathbb{N}$ is a constant depending only on the definition of the $W^{k-1,p}$ -norm.

The next few results are intended as preparation for the proof of Theorem A.3.1.

LEMMA A.3.2. *For any open subset $\mathcal{U} \subset \mathbb{R}^n$ and continuously differentiable function f on \mathcal{U} , the difference quotients $D_i^h f$ converge to $\partial_i f$ uniformly on compact subsets as $h \rightarrow 0$.*

PROOF. Fix a compact subset $\mathcal{K} \subset \mathcal{U}$. Then for every $x \in \mathcal{K}$ and $h \in \mathbb{R} \setminus \{0\}$ sufficiently small, the mean value theorem gives

$$D_i^h f(x) = \partial_i f(x')$$

where

$$x' := (x_1, \dots, x_{i-1}, x_i + th, x_{i+1}, \dots, x_n) \in \mathcal{U}$$

for some $t \in [0, 1]$, so in particular, $|x' - x| \leq |h|$. We then have $|\partial_i f(x) - D_i^h f(x)| = |\partial_i f(x) - \partial_i f(x')|$, and the result follows since both x and x' may be assumed to lie in a compact subset of \mathcal{U} , on which $\partial_i f$ is uniformly continuous. \square

PROPOSITION A.3.3. *Suppose $1 \leq p < \infty$, $\mathcal{U} \subset \mathbb{R}^n$ is an open subset and $f \in W^{1,p}(\mathcal{U})$. Then for any open subset $\mathcal{V} \subset \mathcal{U}$ with compact closure in \mathcal{U} , $\|D^h f\|_{L^p(\mathcal{V})} \leq \|\nabla f\|_{L^p(\mathcal{U})}$ for every $h \neq 0$ with $|h| < \text{dist}(\mathcal{V}, \mathbb{R}^n \setminus \mathcal{U})$, and $D^h f \rightarrow \nabla f$ in L^p on \mathcal{V} as $h \rightarrow 0$.*

PROOF. We show first that for any $f \in W^{1,p}(\mathcal{U})$,

$$(A.11) \quad \|D_i^h f\|_{L^p(\mathcal{V})} \leq \|\partial_i f\|_{L^p(\mathcal{U})}, \quad i = 1, \dots, n$$

for every $\mathcal{V} \subset \mathcal{U}$ with compact closure in \mathcal{U} and every $h \neq 0$ with $|h| < \text{dist}(\mathcal{V}, \mathbb{R}^n \setminus \mathcal{U})$. Indeed, if $f \in W^{1,p}(\mathcal{U}) \cap C^\infty(\mathcal{U})$, then denoting the standard basis of \mathbb{R}^n by (e_1, \dots, e_n) ,

we have

$$\begin{aligned} |D_i^h f(x)| &= \left| \frac{f(x + he_i) - f(x)}{h} \right| = \left| \frac{1}{h} \int_0^1 \frac{d}{dt} f(x + the_i) dt \right| \\ &= \left| \int_0^1 \partial_i f(x + the_i) dt \right| \leq \int_0^1 |\partial_i f(x + the_i)| dt. \end{aligned}$$

Then since any measurable function $g : [0, 1] \rightarrow \mathbb{R}$ satisfies

$$\left(\int_0^1 |g(t)| dt \right)^p \leq \int_0^1 |g(t)|^p dt$$

by Jensen's inequality, this gives

$$\begin{aligned} \|D_i^h f\|_{L^p(\mathcal{V})}^p &= \int_{\mathcal{V}} |D_i^h f(x)|^p d\mu(x) \leq \int_{\mathcal{V}} \left(\int_0^1 |\partial_i f(x + the_i)| dt \right)^p d\mu(x) \\ &\leq \int_{\mathcal{V}} \int_0^1 |\partial_i f(x + the_i)|^p dt d\mu(x) = \int_0^1 \int_{\mathcal{V}} |\partial_i f(x + the_i)|^p d\mu(x) dt \\ &\leq \int_0^1 \|\partial_i f\|_{L^p(\mathcal{U})}^p dt = \|\partial_i f\|_{L^p(\mathcal{U})}^p. \end{aligned}$$

This estimate extends to every $f \in W^{1,p}(\mathcal{U})$ by density of smooth functions.

Next, suppose $f \in W^{1,p}(\mathcal{U})$ and $\epsilon > 0$ is given. Choose a smooth approximation $f_\epsilon \in W^{1,p}(\mathcal{U}) \cap C^\infty(\mathcal{U})$ with $\|f - f_\epsilon\|_{W^{1,p}(\mathcal{U})} < \epsilon/3$. By Lemma A.3.2, $D_i^h f_\epsilon \rightarrow \partial_i f_\epsilon$ in C_{loc}^0 on \mathcal{U} as $h \rightarrow 0$, and since \mathcal{V} has finite measure, this implies we can find $\delta > 0$ such that $|h| < \delta$ implies $\|D_i^h f_\epsilon - \partial_i f_\epsilon\|_{L^p(\mathcal{V})} < \epsilon/3$. Now by (A.11),

$$\|D_i^h f_\epsilon - D_i^h f\|_{L^p(\mathcal{V})} \leq \|\partial_i f_\epsilon - \partial_i f\|_{L^p(\mathcal{U})} \leq \|f_\epsilon - f\|_{W^{1,p}(\mathcal{U})} < \epsilon/3,$$

so combining these estimates gives $\|D_i^h f - \partial_i f\|_{L^p(\mathcal{V})} < \epsilon$ whenever $|h| < \delta$. \square

The proof of the next proposition will require the following standard result from real analysis, known as the **Banach-Alaoglu theorem**. It follows easily from the separability of L^p -spaces for $p < \infty$ together with the duality of L^p and L^q for $1/p + 1/q = 1$; see for instance [LL01, §2.18].

THEOREM A.3.4 (Banach-Alaoglu). *For any measurable subset $\mathcal{U} \subset \mathbb{R}^n$, if $1 < p < \infty$, then every bounded sequence $f_j \in L^p(\mathcal{U})$ has a weakly convergent subsequence, i.e. after passing to a subsequence, one can find a function $f_\infty \in L^p(\mathcal{U})$ such that for every $\varphi \in L^q(\mathcal{U})$ with $1/p + 1/q = 1$, $\int_{\mathcal{U}} f_j \varphi \rightarrow \int_{\mathcal{U}} f_\infty \varphi$.* \square

REMARK A.3.5. One popular way of summarizing the Banach-Alaoglu theorem is the statement that ‘‘closed balls in L^p are weakly compact’’; indeed, if $f_j \in L^p(\mathcal{U})$ satisfies the bound $\|f_j\|_{L^p} \leq C$, then the weak limit f_∞ provided by Theorem A.3.4 also satisfies $\|f_\infty\|_{L^p} \leq C$. The latter follows from the general fact that for any sequence $f_j \in L^p(\mathcal{U})$ converging weakly to some $f_\infty \in L^p(\mathcal{U})$,

$$\|f_\infty\|_{L^p(\mathcal{U})} \leq \liminf \|f_j\|_{L^p(\mathcal{U})}.$$

The proof of this is not hard; see e.g. [LL01, §2.11].

PROPOSITION A.3.6. *Suppose $\mathcal{V} \subset \mathcal{U} \subset \mathbb{R}^n$ are open subsets such that \mathcal{V} has compact closure contained in \mathcal{U} , $1 < p < \infty$, f is a measurable function on \mathcal{U} with $\|f\|_{L^p(\mathcal{V})} < \infty$, and there exist constants $C > 0$ and $\delta > 0$ such that*

$$\|D_i^h f\|_{L^p(\mathcal{V})} \leq C \quad \text{whenever } 0 < |h| < \delta.$$

Then $f|_{\mathcal{V}}$ has a weak partial derivative $\partial_i f \in L^p(\mathcal{V})$ satisfying $\|\partial_i f\|_{L^p(\mathcal{V})} \leq C$.

PROOF. For any sequence $h_j \rightarrow 0$ of sufficiently small nonzero real numbers, the sequence $D_i^{h_j} f$ satisfies $\|D_i^{h_j} f\|_{L^p(\mathcal{V})} \leq C$, thus the Banach-Alaoglu theorem implies that after passing to a subsequence, one finds a function $g \in L^p(\mathcal{V})$ with $\|g\|_{L^p(\mathcal{V})} \leq C$ such that

$$\int_{\mathcal{V}} (D_i^{h_j} f) \varphi \rightarrow \int_{\mathcal{V}} g \varphi$$

for all $\varphi \in L^q(\mathcal{V})$, where $1/p + 1/q = 1$. In particular, this is true for all test functions $\varphi \in C_0^\infty(\mathcal{V})$, and in this case there is an ‘‘integration by parts’’ relation

$$\begin{aligned} \int_{\mathcal{V}} (D_i^{h_j} f) \varphi &= \int_{\mathcal{V}} \frac{f(x + h_j e_i) - f(x)}{h_j} \varphi(x) d\mu(x) \\ &= - \int_{\mathcal{V}} f(x) \frac{\varphi(x - h_j e_i) - \varphi(x)}{-h_j} d\mu(x) = - \int_{\mathcal{V}} f D_i^{-h_j} \varphi. \end{aligned}$$

By Lemma A.3.2, $D_i^{-h_j} \varphi \rightarrow \partial_i \varphi$ uniformly on \mathcal{V} and thus also in $L^q(\mathcal{V})$, so taking the limit of the integrals, we’ve shown

$$\int_{\mathcal{V}} g \varphi = - \int_{\mathcal{V}} f \partial_i \varphi \quad \text{for all } \varphi \in C_0^\infty(\mathcal{V}),$$

or in other words, $\partial_i f = g \in L^p(\mathcal{V})$. □

PROOF OF THEOREM A.3.1. The two statements in the theorem follow by applying Propositions A.3.3 and A.3.6 respectively to $\partial^\alpha f$ for every multiindex α with $|\alpha| \leq k-1$, using the fact that $D^h(\partial^\alpha f) = \partial^\alpha(D^h f)$. For the bound on $\|\nabla f\|_{W^{k-1,p}(\mathcal{V})}$, we observe that by assumption,

$$\|D^h f\|_{W^{k-1,p}(\mathcal{V})} = \sum_{|\alpha| \leq k-1} \|\partial^\alpha(D^h f)\|_{L^p(\mathcal{V})} = \sum_{|\alpha| \leq k-1} \|D^h(\partial^\alpha f)\|_{L^p(\mathcal{V})} \leq C,$$

thus each individual term in this sum satisfies $\|D^h(\partial^\alpha f)\|_{L^p(\mathcal{V})} \leq C$, implying $\|\nabla(\partial^\alpha f)\|_{L^p(\mathcal{V})} \leq C$ and thus

$$\begin{aligned} \|\nabla f\|_{W^{k-1,p}(\mathcal{V})} &= \sum_{|\alpha| \leq k-1} \|\partial^\alpha(\nabla f)\|_{L^p(\mathcal{V})} = \sum_{|\alpha| \leq k-1} \|\nabla(\partial^\alpha f)\|_{L^p(\mathcal{V})} \\ &\leq \sum_{|\alpha| \leq k-1} C =: m_{k,p} C. \end{aligned}$$

□

A.4. Spaces of sections of vector bundles

In this section, fix a field

$$\mathbb{F} := \mathbb{R} \text{ or } \mathbb{C},$$

assume M is a smooth n -dimensional manifold, possibly with boundary, and $\pi : E \rightarrow M$ is a smooth vector bundle of rank m over \mathbb{F} . This comes with a “bundle atlas” $\mathcal{A}(\pi)$, a set whose elements $\alpha \in \mathcal{A}(\pi)$ each consist of the following data:

- (1) An open subset $\mathcal{U}_\alpha \subset M$;
- (2) A smooth local coordinate chart $\varphi_\alpha : \mathcal{U}_\alpha \xrightarrow{\cong} \Omega_\alpha$, where Ω_α is an open subset of $\mathbb{R}_+^n := \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n \geq 0\}$;
- (3) A smooth local trivialization $\Phi_\alpha : E|_{\mathcal{U}_\alpha} \xrightarrow{\cong} \mathcal{U}_\alpha \times \mathbb{F}^m$.

Smoothness of φ_α and Φ_α means as usual that for every pair $\alpha, \beta \in \mathcal{A}(\pi)$, the coordinate transformations

$$\varphi_{\beta\alpha} := \varphi_\beta^{-1} \circ \varphi_\alpha : \Omega_{\alpha\beta} \xrightarrow{\cong} \Omega_{\beta\alpha}, \quad \Omega_{\alpha\beta} := \varphi_\alpha(\mathcal{U}_\alpha \cap \mathcal{U}_\beta)$$

and transition maps

$$g_{\beta\alpha} : \mathcal{U}_\alpha \cap \mathcal{U}_\beta \rightarrow \text{GL}(m, \mathbb{F}) \quad \text{such that} \quad \Phi_\beta \circ \Phi_\alpha^{-1}(x, v) = (x, g_{\beta\alpha}(x)v) \\ \text{for } x \in \mathcal{U}_\alpha \cap \mathcal{U}_\beta, v \in \mathbb{F}^m$$

are smooth, and we shall assume the bundle atlas is maximal in the sense that any triple $(\mathcal{U}, \varphi, \Phi)$ that is smoothly compatible with every $\alpha \in \mathcal{A}(\pi)$ also belongs to $\mathcal{A}(\pi)$.

Any $\alpha \in \mathcal{A}(\pi)$ now associates to sections $\eta : M \rightarrow E$ their local coordinate representatives

$$\eta^\alpha := \text{pr}_2 \circ \Phi_\alpha \circ \eta \circ \varphi_\alpha^{-1} : \Omega_\alpha \rightarrow \mathbb{F}^m,$$

where $\text{pr}_2 : \mathcal{U}_\alpha \times \mathbb{F}^m \rightarrow \mathbb{F}^m$ is the projection, and the representatives with respect to two distinct $\alpha, \beta \in \mathcal{A}(\pi)$ are related by

$$\eta^\beta = (g_{\beta\alpha} \circ \varphi_\beta^{-1})(\eta^\alpha \circ \varphi_{\alpha\beta}) \quad \text{on } \Omega_{\beta\alpha} \subset \Omega_\beta.$$

For $p \in [1, \infty]$ and each integer $k \geq 0$, we then define the topological vector space of sections of class $W_{\text{loc}}^{k,p}$ by

$$W_{\text{loc}}^{k,p}(E) := \left\{ \eta : M \rightarrow E \mid \begin{array}{l} \text{sections such that } \eta^\alpha \in W_{\text{loc}}^{k,p}(\mathring{\Omega}_\alpha, \mathbb{F}^m) \\ \text{for all } \alpha \in \mathcal{A}(\pi) \end{array} \right\},$$

where convergence $\eta_j \rightarrow \eta$ in $W_{\text{loc}}^{k,p}(E)$ means that $\eta_j^\alpha \rightarrow \eta^\alpha$ in $W_{\text{loc}}^{k,p}(\mathring{\Omega}_\alpha, \mathbb{F}^m)$ for all $\alpha \in \mathcal{A}(\pi)$. Note that Ω_α is not necessarily an open subset of \mathbb{R}^n since it may contain points in $\partial\mathbb{R}_+^n = \mathbb{R}^{n-1} \times \{0\}$, but its interior $\mathring{\Omega}_\alpha$ is open in \mathbb{R}^n , and $W_{\text{loc}}^{k,p}(\mathring{\Omega}_\alpha)$ is thus defined as in §A.1. Strictly speaking, elements of $\eta \in W_{\text{loc}}^{k,p}(E)$ are not sections but *equivalence classes* of sections defined almost everywhere—the latter notion is defined with respect to any measure arising from a smooth volume element on M , and it does not depend on this choice.

It turns out that $W_{\text{loc}}^{k,p}(E)$ can be given the structure of a Banach space if M is compact. This follows from the fact that M can then be covered by a finite subset of the atlas $\mathcal{A}(\pi)$, but we must be a little bit careful: not all charts in $\mathcal{A}(\pi)$ are equally

suitable for defining $W^{k,p}$ -norms on sections, because e.g. even a nice smooth section $\eta \in \Gamma(E)$ may have $\|\eta^\alpha\|_{W^{k,p}(\hat{\Omega}_\alpha)} = \infty$ if $\Omega_\alpha \subset \mathbb{R}_+^n$ is unbounded. One way to deal with this is as follows: we will say that $\alpha \in \mathcal{A}(\pi)$ is a **precompact chart** if there exists $\alpha' \in \mathcal{A}(\pi)$ and a compact subset $\mathcal{K} \subset M$ such that

$$\mathcal{U}_\alpha \subset \mathcal{K} \subset \mathcal{U}_{\alpha'}.$$

When this is the case, $\Omega_\alpha \subset \mathbb{R}_+^n$ is necessarily bounded, and the transition maps between two precompact charts necessarily have bounded derivatives of all orders, as they are restrictions to precompact subsets of maps that are smooth on larger domains. If M is compact, then one can always find a finite subset $I \subset \mathcal{A}(\pi)$ consisting of precompact charts such that $M = \bigcup_{\alpha \in I} \mathcal{U}_\alpha$.

DEFINITION A.4.1. Suppose $E \rightarrow M$ is a smooth vector bundle over a compact manifold M , and $I \subset \mathcal{A}(\pi)$ is a finite set of precompact charts such that $\{\mathcal{U}_\alpha\}_{\alpha \in I}$ is an open cover of M . We then define $W^{k,p}(E)$ as the vector space of all sections $\eta : M \rightarrow E$ for which the norm

$$\|\eta\|_{W^{k,p}} := \|\eta\|_{W^{k,p}(E)} := \sum_{\alpha \in I} \|\eta^\alpha\|_{W^{k,p}(\hat{\Omega}_\alpha)}$$

is finite.

The norm in the above definition depends on auxiliary choices, but it is easy to see that the resulting definition of the space $W^{k,p}(E)$ and its topology do not. In fact:

PROPOSITION A.4.2. *If M is compact, then $W^{k,p}(E) = W_{\text{loc}}^{k,p}(E)$, and a sequence η_j converges to η in $W_{\text{loc}}^{k,p}(E)$ if and only if the norm given in Definition A.4.1 satisfies $\|\eta_j - \eta\|_{W^{k,p}(E)} \rightarrow 0$.*

The proposition is an immediate consequence of the following.

LEMMA A.4.3. *Suppose M is a smooth manifold, $\pi : E \rightarrow M$ is a smooth vector bundle, $\{\beta\} \cup J \subset \mathcal{A}(\pi)$ is a finite collection of charts such that $M = \bigcup_{\alpha \in J} \mathcal{U}_\alpha$ and all coordinate transformations and transition maps relating any two charts in the collection $\{\beta\} \cup J$ have bounded derivatives of all orders (e.g. it suffices to assume all are precompact). Then there exists a constant $c > 0$ such that*

$$\|\eta^\beta\|_{W^{k,p}(\hat{\Omega}_\beta)} \leq c \sum_{\alpha \in J} \|\eta^\alpha\|_{W^{k,p}(\hat{\Omega}_\alpha)}$$

for all sections $\eta : M \rightarrow E$ with $\eta^\alpha \in W^{k,p}(\hat{\Omega}_\alpha)$ for every $\alpha \in J$.

PROOF. Choose a partition of unity $\{\rho_\alpha : M \rightarrow [0, 1]\}_{\alpha \in J}$ subordinate to the finite open cover $\{\mathcal{U}_\alpha\}_{\alpha \in J}$. Now $\eta = \sum_{\alpha \in J} \rho_\alpha \eta$, and each $\rho_\alpha \eta$ is supported in \mathcal{U}_α , so $(\rho_\alpha \eta)^\beta$ has support in $\Omega_{\beta\alpha} = \varphi_\beta(\mathcal{U}_\alpha \cap \mathcal{U}_\beta)$. Thus using Theorem A.2.8 with the fact that $g_{\beta\alpha}$, φ_β^{-1} , $\varphi_{\alpha\beta}$ and $\varphi_{\beta\alpha} = \varphi_{\alpha\beta}^{-1}$ are all smooth functions with bounded derivatives

of all orders on the domains in question, we find

$$\begin{aligned}
\|\eta^\beta\|_{W^{k,p}(\hat{\Omega}_\beta)} &= \left\| \sum_{\alpha \in J} (\rho_\alpha \eta)^\beta \right\|_{W^{k,p}(\hat{\Omega}_\beta)} \leq \sum_{\alpha \in J} \|(\rho_\alpha \eta)^\beta\|_{W^{k,p}(\hat{\Omega}_{\beta\alpha})} \\
&= \sum_{\alpha \in J} \|(\rho_\alpha \circ \varphi_\beta^{-1})(g_{\beta\alpha} \circ \varphi_\beta^{-1})(\eta^\alpha \circ \varphi_{\alpha\beta})\|_{W^{k,p}(\hat{\Omega}_{\beta\alpha})} \\
&\leq c \sum_{\alpha \in J} \|\eta^\alpha\|_{W^{k,p}(\hat{\Omega}_{\alpha\beta})} \leq c \sum_{\alpha \in J} \|\eta^\alpha\|_{W^{k,p}(\hat{\Omega}_\alpha)}.
\end{aligned}$$

□

COROLLARY A.4.4. *If M is compact, then the norm on $W^{k,p}(E)$ given by Definition A.4.1 is independent of all auxiliary choices up to equivalence of norms.* □

THEOREM A.4.5. *For any smooth vector bundle $\pi : E \rightarrow M$ over a compact manifold M , $W^{k,p}(E)$ is a Banach space.*

PROOF. If $\eta_j \in W^{k,p}(E)$ is a Cauchy sequence, then for some chosen finite collection $I \subset \mathcal{A}(\pi)$ of precompact charts covering M , the sequences η_j^α for $\alpha \in I$ are Cauchy in $W^{k,p}(\hat{\Omega}_\alpha)$ and thus have limits $\xi^{(\alpha)} \in W^{k,p}(\hat{\Omega}_\alpha, \mathbb{F}^m)$. Choosing a partition of unity $\{\rho_\alpha : M \rightarrow [0, 1]\}_{\alpha \in I}$ subordinate to $\{\mathcal{U}_\alpha\}_{\alpha \in I}$, we can now associate to each $\alpha \in I$ a section $\eta_{\infty,\alpha} \in W^{k,p}(E)$ characterized uniquely by the condition that it vanishes outside of \mathcal{U}_α and is represented in the trivialization on \mathcal{U}_α by

$$\eta_{\infty,\alpha}^\alpha = (\rho_\alpha \circ \varphi_\alpha^{-1})\xi^{(\alpha)}.$$

We claim that $\rho_\alpha \eta_j \rightarrow \eta_{\infty,\alpha}$ in $W^{k,p}(E)$ for each $\alpha \in I$. Indeed, we have

$$(\rho_\alpha \eta_j)^\alpha = (\rho_\alpha \circ \varphi_\alpha^{-1})\eta_j^\alpha \rightarrow (\rho_\alpha \circ \varphi_\alpha^{-1})\xi^{(\alpha)} = \eta_{\infty,\alpha}^\alpha \quad \text{in } W^{k,p}(\hat{\Omega}_\alpha)$$

since $\eta_j^\alpha \rightarrow \xi^{(\alpha)}$. For all other $\beta \in I$ not equal to α , $(\rho_\alpha \eta_j)^\beta - \eta_{\infty,\alpha}^\beta \in W^{k,p}(\hat{\Omega}_{\beta\alpha}, \mathbb{F}^m)$ has support in $\Omega_{\beta\alpha} = \varphi_\beta(\mathcal{U}_\alpha \cap \mathcal{U}_\beta)$, thus

$$\|(\rho_\alpha \eta_j)^\beta - \eta_{\infty,\alpha}^\beta\|_{W^{k,p}(\hat{\Omega}_{\beta\alpha})} = \|(\rho_\alpha \eta_j)^\beta - \eta_{\infty,\alpha}^\beta\|_{W^{k,p}(\hat{\Omega}_{\beta\alpha})} \leq c \|(\rho_\alpha \eta_j)^\alpha - \eta_{\infty,\alpha}^\alpha\|_{W^{k,p}(\hat{\Omega}_\alpha)},$$

where the inequality comes from Lemma A.4.3 after replacing M with \mathcal{U}_α , and \mathcal{U}_β with $\mathcal{U}_\beta \cap \mathcal{U}_\alpha$ (note that the lemma does not require M to be compact). With the claim established, we have

$$\eta_j = \sum_{\alpha \in I} \rho_\alpha \eta_j \rightarrow \sum_{\alpha \in I} \eta_{\infty,\alpha} \quad \text{in } W^{k,p}(E).$$

□

REMARK A.4.6. One can use exactly the same approach to show that when M is compact, the space $C^k(E)$ of C^k -smooth sections $\eta : M \rightarrow E$ has a canonical (up to equivalence of norms) Banach space structure for each finite integer $k \geq 0$ such that convergence in the C^k -norm is equivalent to uniform convergence of all derivatives up to order k .

EXERCISE A.4.7. For $\mathcal{U} \subset \mathbb{R}^n$ an open subset, the space $W_{\text{loc}}^{k,p}(\mathcal{U})$ was defined in §A.1, but one can give it an alternative definition in the present context by viewing functions on \mathcal{U} as sections of a trivial vector bundle over \mathcal{U} , with the latter viewed as a noncompact smooth n -manifold. Show that these two definitions of $W_{\text{loc}}^{k,p}(\mathcal{U})$ are equivalent.

EXERCISE A.4.8. Suppose $\mathcal{U} \subset \mathbb{R}^n$ is a bounded open subset with smooth boundary, so its closure $\overline{\mathcal{U}} \subset \mathbb{R}^n$ is a smooth compact submanifold with boundary, and let $E \rightarrow \overline{\mathcal{U}}$ be a trivial vector bundle. Show that there is a canonical Banach space isomorphism between $W^{k,p}(\mathcal{U})$ as defined in §A.1 and $W^{k,p}(E)$ as defined in the present section. *Hint: Recall that sections in $W^{k,p}(E)$ are only required to be defined almost everywhere, so in particular if the domain M is a manifold with boundary, they need not be well defined on ∂M .*

In light of Exercise A.4.8, the natural generalization of $W_0^{k,p}(\mathcal{U})$ in the present setting is

$$W_0^{k,p}(E) := \overline{C_0^\infty(E|_{M \setminus \partial M})},$$

i.e. it is the closure in the $W^{k,p}$ -norm of the space of smooth sections that vanish near the boundary. Density of smooth sections will imply that this is the same as $W^{k,p}(E)$ if M is closed, but in general $W_0^{k,p}(E)$ is a closed subspace of $W^{k,p}(E)$.

The partition of unity argument in Theorem A.4.5 contains all the essential ideas needed to generalize results about Sobolev spaces on domains in \mathbb{R}^n to compact manifolds. We now state the essential results, leaving the proofs as exercises.

THEOREM A.4.9. *Assume M is a smooth compact n -manifold, possibly with boundary, $\pi : E \rightarrow M$ is a smooth vector bundle of finite rank, $k \geq 0$ is an integer and $1 \leq p < \infty$. Then the Banach space $W^{k,p}(E)$ has the following properties.*

- (1) *The space $\Gamma(E)$ of smooth sections is dense in $W^{k,p}(E)$.*
- (2) *If $kp > n$, then for each integer $d \geq 0$, there exists a continuous and compact inclusion*

$$W^{k+d,p}(E) \hookrightarrow C^d(E).$$

- (3) *The natural inclusion*

$$W^{k+1,p}(E) \hookrightarrow W^{k,p}(E)$$

is compact.

- (4) *Suppose $F, G \rightarrow M$ are smooth vector bundles such that there exists a smooth bundle map*

$$E \otimes F \rightarrow G : \eta \otimes \xi \mapsto \eta \cdot \xi.$$

Then if $kp > n$ and $0 \leq m \leq k$, there exists a continuous product pairing

$$W^{k,p}(E) \times W^{m,p}(F) \rightarrow W^{m,p}(G) : (\eta, \xi) \mapsto \eta \cdot \xi.$$

In particular, products of $W^{k,p}$ sections give $W^{k,p}$ sections whenever $kp > n$.

- (5) *Suppose $F \rightarrow M$ is another smooth vector bundle, $\mathcal{V} \subset E$ is an open subset that intersects every fiber of E , and we consider the spaces*

$$W^{k,p}(\mathcal{V}) := \{ \eta \in W^{k,p}(E) \mid \eta(M) \subset \mathcal{V} \}$$

and

$$C_M^k(\mathcal{V}, F) := \{ \Phi : \mathcal{V} \rightarrow F \mid \text{fiber-preserving maps of class } C^k \},$$

where the latter is assigned the topology of C^k -convergence on compact subsets. If $kp > n$, then $W^{k,p}(\mathcal{V})$ is an open subset of $W^{k,p}(E)$, and the map

$$C_M^k(\mathcal{V}, F) \times W^{k,p}(\mathcal{V}) \rightarrow W^{k,p}(F) : (\Phi, \eta) \mapsto \Phi \circ \eta$$

is well defined and continuous.

(6) If N is another smooth compact manifold and $\varphi : N \rightarrow M$ is a smooth diffeomorphism, then there is a Banach space isomorphism

$$W^{k,p}(E) \rightarrow W^{k,p}(\varphi^* E) : \eta \mapsto \eta \circ \varphi.$$

□

REMARK A.4.10. It is sometimes useful to extend the definitions and results of this section to vector bundles that are not smooth, e.g. vector bundles of class C^k or $W^{k,p}$, for which all transition maps are required to be of class C^k or $W^{k,p}$ respectively. The latter makes sense in general only if $kp > n$, so that transition maps are at least continuous. Given a bundle of this type, one can enhance the arguments of this section with the aid of Theorem A.2.1 to show that $W^{m,p}(E)$ is a well-defined Banach space for every $m \leq k$, though it would not be well defined if $m > k$. Such spaces arise frequently in global analysis, e.g. if f is a non-smooth element in the Banach manifold \mathcal{B} of $W^{k,p}$ -smooth maps of M into another manifold N , then $f^*TN \rightarrow M$ is in general a vector bundle of class $W^{k,p}$, and $T_f\mathcal{B} = W^{k,p}(f^*TN)$.

A.5. Some remarks on domains with cylindrical ends

For bundles $\pi : E \rightarrow M$ with M noncompact, $W^{k,p}(E)$ is not generally well defined without making additional choices. When $M = \dot{\Sigma} = \Sigma \setminus \Gamma$ is a punctured Riemann surface and $\pi : E \rightarrow \dot{\Sigma}$ is equipped with an asymptotically Hermitian structure $\{(E_z, J_z, \omega_z)\}_{z \in \Gamma}$ as defined in Chapter 4, one nice way to define $W^{k,p}(E)$ was introduced in §4.1: one takes it to be the space of sections in $W_{\text{loc}}^{k,p}(E)$ whose $W^{k,p}$ -norms on each cylindrical end are finite with respect to a choice of asymptotic trivialization. This definition requires the convenient fact that complex vector bundles over S^1 are always trivial, though one can also do without this by using the ideas in the previous section. Indeed, any collection of local trivializations on the asymptotic bundle $E_z \rightarrow S^1$ covering S^1 gives rise via the asymptotically Hermitian structure to a collection of trivializations on E covering the corresponding cylindrical end \dot{U}_z . The key fact is then that S^1 is compact, hence one can always choose such a covering to be finite: combining this with a finite covering of $\dot{\Sigma}$ in the complement of its cylindrical ends by precompact charts, we obtain a covering of $\dot{\Sigma}$ by a finite collection of bundle charts that are not all precompact, but nonetheless have the property that all transition maps have bounded derivatives of all orders. This is enough to define a $W^{k,p}$ -norm for sections of $E \rightarrow \dot{\Sigma}$ as in Definition A.4.1 and to prove that it does not depend on the choices of charts or local trivializations, though it does depend on the asymptotically Hermitian structure.

With this definition understood, one can easily generalize the Sobolev embedding theorem and other important statements in Theorem A.4.9 to the setting of an asymptotically Hermitian bundle over a punctured Riemann surface. We shall leave the details of this generalization as an exercise, but take the opportunity to point out a few important differences from the compact case.

First, since $\dot{\Sigma}$ is not compact, neither are the inclusions

$$W^{k+d,p}(E) \hookrightarrow C^d(E), \quad W^{k+1,p}(E) \hookrightarrow W^{k,p}(E).$$

The proof of compactness fails due to the fact that cylindrical ends require local trivializations over unbounded domains of the form $(0, \infty) \times (0, 1) \subset \mathbb{R}^2$, for which Theorem A.1.10 does not hold. And indeed, considering unbounded shifts on the infinite cylinder $\dot{\Sigma} = \mathbb{R} \times S^1$, it is easy to find a sequence of $W^{k,p}$ -bounded functions with $kp > 2$ that do not have a C^0 -convergent subsequence. That is the bad news.

The good news is that if $\eta \in W^{k+d,p}(E)$ for $kp > 2$, then one can say considerably more about η than just that it is C^d -smooth. Indeed, restricting to one of the cylindrical ends $[0, \infty) \times S^1 \subset \dot{\Sigma}$, notice that the finiteness of the $W^{k+d,p}$ -norm over $\dot{\Sigma}$ implies

$$\|\eta\|_{W^{k+d,p}((R,\infty) \times S^1)} \rightarrow 0 \quad \text{as } R \rightarrow \infty.$$

Since these domains are all naturally diffeomorphic for different values of R , the C^d -norm of η over $(R, \infty) \times S^1$ is bounded by the $W^{k+d,p}$ -norm via a constant that does not depend on R , so this implies an asymptotic decay condition

$$\|\eta\|_{C^d([R,\infty) \times S^1)} \rightarrow 0 \quad \text{as } R \rightarrow \infty$$

for every $\eta \in W^{k+d,p}(E)$.

Here is another useful piece of good news: since $\dot{\Sigma}$ does not have boundary, $W^{k,p}(E) = W_0^{k,p}(E)$.

THEOREM A.5.1. *Given an asymptotically Hermitian bundle E over a punctured Riemann surface $\dot{\Sigma}$, the space $C_0^\infty(E)$ of smooth sections with compact support is dense in $W^{k,p}(E)$ for all $k \geq 0$ and $1 \leq p < \infty$.*

PROOF. We can assume as in Definition A.4.1 that the $W^{k,p}$ -norm for sections η of E is given by

$$\|\eta\|_{W^{k,p}} = \sum_{\alpha \in I} \|\eta^\alpha\|_{W^{k,p}(\Omega_\alpha)},$$

where $I \subset \mathcal{A}(\pi)$ is a finite collection of bundle charts

$$\alpha = \left(\varphi_\alpha : \mathcal{U}_\alpha \xrightarrow{\cong} \Omega_\alpha, \Phi_\alpha : E|_{\mathcal{U}_\alpha} \xrightarrow{\cong} \mathcal{U}_\alpha \times \mathbb{C}^n \right)$$

such that each of the open sets $\Omega_\alpha \subset \mathbb{C}$ is either bounded or (for charts over the cylindrical ends) of the form

$$\Omega_\alpha = (0, \infty) \times \omega_\alpha \subset \mathbb{R}^2 = \mathbb{C}$$

for some bounded open subset $\omega_\alpha \subset \mathbb{R}$. Now given $\eta \in W^{k,p}(E)$, Theorem A.1.1 provides for each $\alpha \in I$ a sequence $\eta_j^\alpha \in W^{k,p}(\Omega_\alpha)$ of smooth functions with bounded

support such that $\eta_j^\alpha \rightarrow \eta^\alpha$ in $W^{k,p}(\Omega_\alpha)$. Choose a partition of unity $\{\rho_\alpha : \dot{\Sigma} \rightarrow [0, 1]\}_{\alpha \in I}$ subordinate to the open cover $\{\mathcal{U}_\alpha\}_{\alpha \in I}$ and let

$$\eta_j := \sum_{\alpha \in I} \rho_\alpha(\eta_j^\alpha \circ \varphi_\alpha) \in W^{k,p}(E).$$

These sections are smooth and have compact support since the η_j^α have bounded support in Ω_α , and they converge in $W^{k,p}$ to η . \square

APPENDIX B

The Floer C_ϵ space

The C_ϵ -topology for functions was introduced by Floer [Flo88b] to provide a Banach manifold of perturbed geometric structures without departing from the smooth category: it is a way to circumvent the annoying fact that spaces of smooth functions which arise naturally in geometric settings are not Banach spaces. The construction of C_ϵ spaces generally depends on several arbitrary choices and is thus far from canonical, but this detail is unimportant since the C_ϵ space itself is never the main object of interest. What is important is merely the properties that it has, namely that it not only embeds continuously into C^∞ and contains an abundance of non-trivial functions, but also is a separable Banach space and can therefore be used in the Sard-Smale theorem for genericity arguments. We shall prove these facts in this appendix.

Fix a smooth finite-rank vector bundle $\pi : E \rightarrow M$ over a finite-dimensional compact manifold M , possibly with boundary. For each integer $k \geq 0$, we denote by $C^k(E)$ the Banach space of C^k -smooth sections of E ; note that the norm on $C^k(E)$ depends on various auxiliary choices but is well defined up to equivalence of norms since M is compact. Now if $\epsilon = (\epsilon_k)_{k=0}^\infty$ is a sequence of positive numbers with $\epsilon_k \rightarrow 0$, set

$$C_\epsilon(E) = \{ \eta \in \Gamma(E) \mid \|\eta\|_{C_\epsilon} < \infty \},$$

where the C_ϵ -norm is defined by

$$(B.1) \quad \|\eta\|_{C_\epsilon} = \sum_{k=0}^{\infty} \epsilon_k \|\eta\|_{C^k}.$$

The norm for $C_\epsilon(E)$ is somewhat more delicate than for $C^k(E)$, e.g. its equivalence class is not obviously independent of auxiliary choices. This remark is meant as a sanity check, but it should not cause extra concern since, in practice, the space $C_\epsilon(E)$ is typically regarded as an auxiliary choice in itself. In many applications, one fixes an open subset $\mathcal{U} \subset M$ and considers the closed subspace

$$C_\epsilon(E; \mathcal{U}) = \{ \eta \in C_\epsilon(E) \mid \eta|_{M \setminus \mathcal{U}} \equiv 0 \}.$$

REMARK B.0.1. The requirement for M to be compact can be relaxed as long as $\mathcal{U} \subset M$ has compact closure: e.g. in one situation of frequent interest in this book, we take M to be the noncompact completion of a symplectic cobordism. In this case $C_\epsilon(E; \mathcal{U})$ can be defined as a closed subspace of $C_\epsilon(E|_{M_0})$ where $M_0 \subset M$ is any compact manifold with boundary that contains the closure of \mathcal{U} . For this reason, we lose no generality in continuing under the assumption that M is compact.

In order to prove things about $C_\epsilon(E)$, we will need to specify a more precise definition of the C^k -norms. To this end, define a sequence of vector bundles $E^{(k)} \rightarrow M$ for integers $k \geq 0$ inductively by

$$E^{(0)} := E, \quad E^{(k+1)} := \text{Hom}(TM, E^{(k)}).$$

Choose connections and bundle metrics on both TM and E ; these induce connections and bundle metrics on each of the $E^{(k)}$, so that for any section $\xi \in \Gamma(E^{(k)})$, the covariant derivative $\nabla\xi$ is now a section of $E^{(k+1)}$. In particular for $\eta \in \Gamma(E)$, we can define the “ k th covariant derivative” of η as a section

$$\nabla^k \eta \in \Gamma(E^{(k)}).$$

Using the bundle metrics to define C^0 -norms for sections of $E^{(k)}$, we can then define

$$\|\eta\|_{C^k(E)} = \sum_{m=0}^k \|\nabla^m \eta\|_{C^0(E^{(m)})},$$

where by convention $\nabla^0 \eta := \eta$. We will assume throughout the following that the C^k -norms appearing in (B.1) are defined in this way.

THEOREM B.0.2. $C_\epsilon(E)$ is a Banach space.

PROOF. We need to show that C_ϵ -Cauchy sequences converge in the C_ϵ -norm. It is clear from the definitions that if $\eta_j \in C_\epsilon(E)$ is Cauchy, then η_j is also C^k -Cauchy for every $k \geq 0$, hence its derivatives $\nabla^k \eta_j$ for every k are C^0 -convergent to continuous sections ξ^k of $E^{(k)}$. This convergence implies that $\xi^{k+1} = \nabla \xi^k$ in the sense of distributions, hence by the equivalence of classical and distributional derivatives (see e.g. [LL01, §6.10]), $\eta_\infty := \xi^0$ is smooth with $\nabla^k \eta_\infty = \xi^k$, so that $\nabla^k \eta_j \rightarrow \nabla^k \eta_\infty$ in $C^0(E^{(k)})$ for all k .

We claim $\eta_\infty \in C_\epsilon(E)$. Choose $N > 0$ such that $\|\eta_i - \eta_j\|_{C_\epsilon} < 1$ for all $i, j \geq N$. Then for every $m \in \mathbb{N}$ and every $i \geq N$,

$$\begin{aligned} \sum_{k=0}^m \epsilon_k \|\eta_i\|_{C^k} &\leq \sum_{k=0}^m \epsilon_k \|\eta_i - \eta_N\|_{C^k} + \sum_{k=0}^m \epsilon_k \|\eta_N\|_{C^k} \\ &\leq \|\eta_i - \eta_N\|_{C_\epsilon} + \|\eta_N\|_{C_\epsilon} < 1 + \|\eta_N\|_{C_\epsilon}. \end{aligned}$$

Fixing m and letting $i \rightarrow \infty$, we then have

$$\sum_{k=0}^m \epsilon_k \|\eta_\infty\|_{C^k} \leq 1 + \|\eta_N\|_{C_\epsilon}$$

for all m , so we can now let $m \rightarrow \infty$ and conclude $\|\eta_\infty\|_{C_\epsilon} \leq 1 + \|\eta_N\|_{C_\epsilon} < \infty$.

The argument that $\|\eta_j - \eta_\infty\|_{C_\epsilon} \rightarrow 0$ as $j \rightarrow \infty$ is similar: pick $\epsilon > 0$ and N such that $\|\eta_i - \eta_j\|_{C_\epsilon} < \epsilon$ for all $i, j \geq N$. Then for a fixed $m \in \mathbb{N}$, we can let $i \rightarrow \infty$ in the expression $\sum_{k=0}^m \epsilon_k \|\eta_i - \eta_j\|_{C^k} < \epsilon$, giving

$$\sum_{k=0}^m \epsilon_k \|\eta_\infty - \eta_j\|_{C^k} \leq \epsilon.$$

This is true for every m , so we can take $m \rightarrow \infty$ and conclude $\|\eta_\infty - \eta_j\|_{C_\epsilon} \leq \epsilon$ for all $j \geq N$. \square

To show that $C_\epsilon(E)$ is also separable, we will follow a hint¹ from [HS95] and embed it isometrically into another Banach space that can be more easily shown to be separable. For each integer $k \geq 0$, define the vector bundle

$$F^{(k)} = E^{(0)} \oplus \dots \oplus E^{(k)},$$

and let X_ϵ denote the vector space of all sequences

$$\xi := (\xi^0, \xi^1, \xi^2, \dots) \in \prod_{k=0}^{\infty} C^0(F^{(k)})$$

such that

$$\|\xi\|_{X_\epsilon} := \sum_{k=0}^{\infty} \epsilon_k \|\xi^k\|_{C^0} < \infty.$$

EXERCISE B.0.3. Adapt the proof of Theorem B.0.2 to show that X_ϵ is also a Banach space.

LEMMA B.0.4. X_ϵ is separable.

PROOF. Since $C^0(F^{(k)})$ is separable for each $k \geq 0$, we can fix countable dense subsets $P^k \subset C^0(F^{(k)})$. The set

$$P := \{(\xi^0, \dots, \xi^N, 0, 0, \dots) \in X_\epsilon \mid N \geq 0 \text{ and } \xi^k \in P^k \text{ for all } k = 0, \dots, N\}$$

is then countable and dense in X_ϵ . \square

THEOREM B.0.5. $C_\epsilon(E)$ is separable.

PROOF. Consider the injective linear map

$$C_\epsilon(E) \hookrightarrow X_\epsilon : \eta \mapsto (\eta, (\eta, \nabla\eta), (\eta, \nabla\eta, \nabla^2\eta), \dots).$$

This is an isometric embedding and thus presents $C_\epsilon(E)$ as a closed linear subspace of X_ϵ , hence the theorem follows from Lemma B.0.4 and the fact that subspaces of separable metric spaces are always separable. \square

Note that given any open subset $\mathcal{U} \subset M$, Theorems B.0.2 and B.0.5 also hold for $C_\epsilon(E; \mathcal{U})$, as a closed subspace of $C_\epsilon(E)$. So far in this discussion, however, there has been no guarantee that $C_\epsilon(E)$ or $C_\epsilon(E; \mathcal{U})$ contains anything other than the zero-section, though it is clear that in theory, one should always be able to enlarge the space by choosing new sequences ϵ_k that converge to zero faster. The following result says that $C_\epsilon(E; \mathcal{U})$ can always be made large enough to be useful in applications.

THEOREM B.0.6. Given an open subset $\mathcal{U} \subset M$, the sequence ϵ_k can be chosen to have the following properties:

- (1) $C_\epsilon(E; \mathcal{U})$ is dense in the space of continuous sections vanishing outside \mathcal{U} .
- (2) Given any point $p \in \mathcal{U}$, a neighborhood $\mathcal{N}_p \subset \mathcal{U}$ of p , a number $\delta > 0$ and a continuous section η_0 of E , there exists a section $\eta \in \Gamma(E)$ and a smooth compactly supported function $\beta : \mathcal{N}_p \rightarrow [0, 1]$ such that

$$\beta\eta \in C_\epsilon(E; \mathcal{U}), \quad \beta(p)\eta(p) = \eta_0(p), \quad \text{and} \quad \|\eta - \eta_0\|_{C^0} < \delta.$$

¹Thanks to Sam Lisi for explaining to me what the hint in [HS95] was referring to.

PROOF. Note first that it suffices to find two separate sequences ϵ_k and ϵ'_k that have the first and second property respectively, as the sequence of minima $\min(\epsilon_k, \epsilon'_k)$ will then have both properties.

The following construction for the first property is based on a suggestion by Barney Bramham. Observe first that the space $C^0(E; \mathcal{U})$ of continuous sections vanishing outside \mathcal{U} is a closed subspace of $C^0(E)$ and is thus separable, so we can choose a countable C^0 -dense subset $P \subset C^0(E; \mathcal{U})$. Moreover, the space of *smooth* sections vanishing outside \mathcal{U} is dense in $C^0(E; \mathcal{U})$, hence we can assume without loss of generality that the sections in P are smooth. Now write $P = \{\eta_1, \eta_2, \eta_3, \dots\}$ and define $\epsilon_k > 0$ for every integer $k \geq 0$ to have the property

$$\epsilon_k < \frac{1}{2^k} \min \left\{ \frac{1}{\|\eta_1\|_{C^k}}, \dots, \frac{1}{\|\eta_k\|_{C^k}} \right\}.$$

Then every η_j is in $C_\epsilon(E; \mathcal{U})$, as

$$\|\eta_j\|_{C_\epsilon} < \sum_{k=0}^{j-1} \epsilon_k \|\eta_j\|_{C^k} + \sum_{k=j}^{\infty} \frac{1}{2^k} < \infty.$$

The second property is essentially local, so it can be deduced from Lemma B.0.7 below. \square

LEMMA B.0.7. *Suppose $\beta : \mathring{\mathbb{D}}^n \rightarrow [0, 1]$ is a smooth function with compact support on the open unit ball $\mathring{\mathbb{D}}^n \subset \mathbb{R}^n$ and $\beta(0) = 1$. One can choose a sequence of positive numbers $\epsilon_k \rightarrow 0$ such that for every $\eta_0 \in \mathbb{R}^m$ and $r > 0$, the function $\eta : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by*

$$\eta(p) := \beta(p/r)\eta_0$$

satisfies $\sum_{k=0}^{\infty} \epsilon_k \|\eta\|_{C^k} < \infty$.

PROOF. Define $\epsilon_k > 0$ so that for $k \geq 1$,

$$\epsilon_k = \frac{1}{k^k \|\beta\|_{C^k}}.$$

Then

$$\sum_{k=1}^{\infty} \epsilon_k \|\eta\|_{C^k} \leq \sum_{k=1}^{\infty} \frac{1}{k^k \|\beta\|_{C^k}} \frac{\|\beta\|_{C^k}}{r^k} = \sum_{k=1}^{\infty} \left(\frac{1}{r}\right)^k < \infty.$$

\square

APPENDIX C

Genericity in the space of asymptotic operators

The purpose of this appendix is to prove Lemma 3.4.3, which was needed for our definition of spectral flow in §3.4. The proof combines some ideas from that section with the technique used in Chapter 9 to prove generic transversality of moduli spaces via the Sard-Smale theorem. Some knowledge of that technique should thus be considered a prerequisite for this appendix; if you have never seen it before and were directed here after reading the statement of Lemma 3.4.3, you might want to skip this for now and come back after you've read as far as Chapter 9.

Recalling the notation from Chapter 3, we fix the real Hilbert spaces

$$\mathcal{H} = L^2(S^1, \mathbb{R}^{2n}), \quad \mathcal{D} = H^1(S^1, \mathbb{R}^{2n}),$$

the symmetric index 0 Fredholm operator

$$\mathbf{T}_{\text{ref}} = -J_0 \partial_t : \mathcal{D} \rightarrow \mathcal{H}$$

and, given a bounded family of symmetric matrices $S \in L^\infty(S^1, \text{End}^{\text{sym}}(\mathbb{R}^{2n}))$, refer to any operator of the form

$$\mathbf{A} = -J_0 \partial_t - S : \mathcal{D} \rightarrow \mathcal{H}$$

as an **asymptotic operator**. Such operators belong to the space of symmetric compact perturbations of \mathbf{T}_{ref} ,

$$\text{Fred}_{\mathbb{R}}^{\text{sym}}(\mathcal{D}, \mathcal{H}, \mathbf{T}_{\text{ref}}) = \{ \mathbf{T}_{\text{ref}} + \mathbf{K} : \mathcal{D} \rightarrow \mathcal{H} \mid \mathbf{K} \in \mathcal{L}_{\mathbb{R}}^{\text{sym}}(\mathcal{H}) \},$$

which we regard as a smooth Banach manifold via its obvious identification with the space $\mathcal{L}_{\mathbb{R}}^{\text{sym}}(\mathcal{H})$ of symmetric bounded linear operators on \mathcal{H} . For $k \in \mathbb{N}$, we denote by

$$\text{Fred}_{\mathbb{R}}^{\text{sym},k}(\mathcal{D}, \mathcal{H}, \mathbf{T}_{\text{ref}}) \subset \text{Fred}_{\mathbb{R}}^{\text{sym}}(\mathcal{D}, \mathcal{H}, \mathbf{T}_{\text{ref}})$$

the finite-codimensional submanifold determined by the condition $\dim_{\mathbb{R}} \ker \mathbf{A} = \dim_{\mathbb{R}} \text{coker } \mathbf{A} = k$.

Here is the statement of Lemma 3.4.3 again.

LEMMA. *Fix a smooth path $[-1, 1] \rightarrow L^\infty(S^1, \text{End}^{\text{sym}}(\mathbb{R}^{2n})) : s \mapsto S_s$ and consider the 1-parameter family of symmetric index 0 Fredholm operators*

$$\mathbf{A}_s := -J_0 \partial_t - S_s : H^1(S^1, \mathbb{R}^{2n}) \rightarrow L^2(S^1, \mathbb{R}^{2n})$$

for $s \in [-1, 1]$, assuming $\mathbf{A}_{\pm 1}$ are isomorphisms. Then after replacing S_s by a family of the form $\tilde{S}_s(t) := S_s(t) + B(s, t)$ for some smooth function $B : [-1, 1] \rightarrow \text{End}^{\text{sym}}(\mathbb{R}^{2n})$ that vanishes for $s = \pm 1$ and may be assumed arbitrarily C^∞ -small, one can arrange that the following conditions hold:

- (1) For each $s \in (-1, 1)$, all eigenvalues of \mathbf{A}_s are simple.
(2) All intersections of the smooth path

$$(-1, 1) \rightarrow \text{Fred}_{\mathbb{R}}^{\text{sym}}(\mathcal{D}, \mathcal{H}, \mathbf{T}_{\text{ref}}) : s \mapsto \mathbf{A}_s$$

with $\text{Fred}_{\mathbb{R}}^{\text{sym}, 1}(\mathcal{D}, \mathcal{H}, \mathbf{T}_{\text{ref}})$ are transverse.

We shall now prove this by constructing a Floer-type space of C_ϵ -smooth (see Appendix B) perturbed families of asymptotic operators, and using the Sard-Smale theorem to find a countable collection of comeager subsets whose intersection contains perturbations achieving the desired conditions.

Choose a sequence of positive numbers $\epsilon = (\epsilon_k)_{k=0}^\infty$ with $\epsilon_k \rightarrow 0$ to define a separable Banach space

$$\mathcal{A}_\epsilon := \{B \in C^\infty([-1, 1] \times S^1, \text{End}^{\text{sym}}(\mathbb{R}^{2n})) \mid \|B\|_{C_\epsilon} < \infty \text{ and } B(\pm 1, \cdot) \equiv 0\},$$

and assume via Theorem B.0.6 that \mathcal{A}_ϵ is dense in the Banach space of continuous functions $[-1, 1] \times S^1 \rightarrow \text{End}^{\text{sym}}(\mathbb{R}^{2n})$ vanishing at $\{\pm 1\} \times S^1$. We then consider perturbed 1-parameter families of asymptotic operators of the form

$$\mathbf{A}_s^B := \mathbf{A}_s + B(s, \cdot) : \mathcal{D} \rightarrow \mathcal{H}$$

for $B \in \mathcal{A}_\epsilon$, $s \in [-1, 1]$. Remarks 3.4.1 and 3.4.2 imply that the perturbed family defines a smooth path in $\text{Fred}_{\mathbb{R}}^{\text{sym}}(\mathcal{D}, \mathcal{H}, \mathbf{T}_{\text{ref}})$ as long as the original path $s \mapsto \mathbf{A}_s$ is smooth in $L^\infty(S^1, \text{End}^{\text{sym}}(\mathbb{R}^{2n}))$. For each $k \in \mathbb{N}$ and $B \in \mathcal{A}_\epsilon$, define the set

$$\mathcal{V}^k(B) = \{(s, \lambda) \in (-1, 1) \times \mathbb{R} \mid \dim_{\mathbb{R}} \ker(\mathbf{A}_s^B - \lambda) = k\}.$$

To show that eigenvalues are generically simple, we need to show that for a comeager set of choices of $B \in \mathcal{A}_\epsilon$, $\mathcal{V}^k(B)$ is empty for all $k \geq 2$. Given $(s_0, \lambda_0) \in \mathcal{V}^k(B)$, recall from §3.4 that there exist decompositions

$$\mathcal{D} = V \oplus K, \quad \mathcal{H} = W \oplus K$$

where $K = \ker(\mathbf{A}_{s_0}^B - \lambda_0)$, $W = \text{im}(\mathbf{A}_{s_0}^B - \lambda_0)$ is the L^2 -orthogonal complement of K , and $V = W \cap \mathcal{D}$, so that any symmetric bounded linear operator \mathbf{T} in a sufficiently small neighborhood $\mathcal{O} \subset \mathcal{L}_{\mathbb{R}}^{\text{sym}}(\mathcal{D}, \mathcal{H})$ of $\mathbf{A}_{s_0}^B - \lambda_0$ can be written in block form

$$\mathbf{T} = \begin{pmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{pmatrix}$$

with $\mathbf{A} : V \rightarrow W$ invertible. This gives rise to a smooth map

$$\Phi : \mathcal{O} \rightarrow \text{End}_{\mathbb{R}}^{\text{sym}}(K) : \mathbf{T} \mapsto \mathbf{D} - \mathbf{C}\mathbf{A}^{-1}\mathbf{B}$$

whose zero set is precisely the set of nearby symmetric operators with k -dimensional kernel. A neighborhood of (s_0, λ_0) in $\mathcal{V}^k(B)$ can thus be identified with the zero set of the map

$$\Psi_B(s, \lambda) := \Phi(\mathbf{A}_s^B - \lambda) \in \text{End}_{\mathbb{R}}^{\text{sym}}(K),$$

defined for $(s, \lambda) \in (-1, 1) \times \mathbb{R}$ sufficiently close to (s_0, λ_0) . Notice that the derivative $d\Psi_B(s, \lambda) : \mathbb{R} \oplus \mathbb{R} \rightarrow \text{End}_{\mathbb{R}}^{\text{sym}}(K)$ is Fredholm since its domain and target are both finite dimensional, and it can only ever be surjective when $k = \dim_{\mathbb{R}} K = 1$.

The following space will now play the role of a “universal moduli space” as in Chapter 9: let

$$\mathcal{V}^k = \{(s, \lambda, B) \in (-1, 1) \times \mathbb{R} \times \mathcal{A}_\epsilon \mid (s, \lambda) \in \mathcal{V}^k(B)\}.$$

The proof that this is a smooth Banach manifold depends on the following algebraic lemma.

LEMMA C.0.1. *Fix an asymptotic operator $\mathbf{A} = -J_0 \partial_t - S$ and a linear transformation*

$$\Upsilon : \ker \mathbf{A} \rightarrow \ker \mathbf{A}$$

that is symmetric with respect to the L^2 -product. Then there exists a continuous loop $B : S^1 \rightarrow \text{End}^{\text{sym}}(\mathbb{R}^{2n})$ such that

$$\langle \eta, B\xi \rangle_{L^2} = \langle \eta, \Upsilon\xi \rangle_{L^2}$$

for all $\eta, \xi \in \ker \mathbf{A}$.

PROOF. Note first that every nontrivial loop $\eta \in \ker \mathbf{A} \subset H^1(S^1, \mathbb{R}^{2n})$ is continuous and nowhere zero due to the generalized existence/uniqueness result for solutions to linear ODEs in Exercise 3.3.1. It follows that if we fix a basis (η_1, \dots, η_k) for $\ker \mathbf{A}$, then the vectors $\eta_1(t), \dots, \eta_k(t) \in \mathbb{R}^{2n}$ are also linearly independent for all $t \in S^1$ and thus span a continuous S^1 -family of k -dimensional subspaces $V_t \subset \mathbb{R}^{2n}$, each equipped with a distinguished basis. There is therefore a unique continuous S^1 -family of linear transformations $\hat{B}(t) : V_t \rightarrow V_t$ such that for every $\eta \in \ker \mathbf{A}$, $\hat{B}(t)\eta(t) = (\Upsilon\eta)(t)$ for all t . Extend $\hat{B}(t)$ arbitrarily to a continuous family of linear maps on \mathbb{R}^{2n} .

The matrices $\hat{B}(t) \in \text{End}(\mathbb{R}^{2n})$ need not be symmetric, but they do satisfy

$$\langle \eta, \hat{B}\xi \rangle_{L^2} = \langle \eta, \Upsilon\xi \rangle_{L^2} \quad \text{for all } \eta, \xi \in \ker \mathbf{A}.$$

Since Υ is symmetric, this implies moreover that for all $\eta, \xi \in \ker \mathbf{A}$,

$$\langle \eta, \Upsilon\xi \rangle_{L^2} = \langle \xi, \Upsilon\eta \rangle_{L^2} = \langle \xi, \hat{B}\eta \rangle_{L^2} = \langle \eta, \hat{B}^T\xi \rangle_{L^2}.$$

The loop $B := \frac{1}{2}(\hat{B} + \hat{B}^T)$ thus has the desired properties. \square

Now using the previously described construction in the space of symmetric Fredholm operators, a neighborhood of any point (s_0, λ_0, B_0) in \mathcal{V}^k can be identified with the zero set of a smooth map of the form

$$\Psi(s, \lambda, B) := \Psi_B(s, \lambda) \in \text{End}_{\mathbb{R}}^{\text{sym}}(K),$$

defined for all (s, λ, B) sufficiently close to (s_0, λ_0, B_0) in $(-1, 1) \times \mathbb{R} \times \mathcal{A}_\epsilon$, where $K = \ker(\mathbf{A}_{s_0}^{B_0} - \lambda_0)$. The partial derivative of Ψ with respect to the third variable at (s_0, λ_0, B_0) is then a linear map

$$\mathbf{L} := D_3\Psi(s_0, \lambda_0, B_0) : \mathcal{A}_\epsilon \rightarrow \text{End}_{\mathbb{R}}^{\text{sym}}(K)$$

of the form

$$(C.1) \quad \mathbf{L}B : K \rightarrow K : \eta \mapsto \pi_K(B(s_0, \cdot)\eta),$$

where $\pi_K : W \oplus K \rightarrow K$ is the orthogonal projection. We claim that \mathbf{L} is surjective. Indeed, for any $\Upsilon \in \text{End}_{\mathbb{R}}^{\text{sym}}(K)$, Lemma C.0.1 provides a continuous loop $C_0 : S^1 \rightarrow \text{End}^{\text{sym}}(\mathbb{R}^{2n})$ such that

$$\pi_K(C_0\eta) = \Upsilon\eta \quad \text{for all } \eta \in K,$$

and this can be extended to a continuous function $C : [-1, 1] \times S^1 \rightarrow \text{End}^{\text{sym}}(\mathbb{R}^{2n})$ satisfying $C(s_0, \cdot) \equiv C_0$ and $C(\pm 1, \cdot) \equiv 0$ since $s_0 \neq \pm 1$. The function C might fail to be of class C_ϵ , but since it can be approximated arbitrarily well in the C^0 -norm by functions in \mathcal{A}_ϵ , we conclude that the image of \mathbf{L} is dense in $\text{End}_{\mathbb{R}}^{\text{sym}}(K)$. Since the latter is finite dimensional, the claim follows.

The implicit function theorem now gives \mathcal{V}^k the structure of a smooth Banach submanifold of $(-1, 1) \times \mathbb{R} \times \mathcal{A}_\epsilon$, and it is separable since the latter is also separable. Consider the projection

$$(C.2) \quad \pi : \mathcal{V}^k \rightarrow \mathcal{A}_\epsilon : (s, \lambda, B) \mapsto B,$$

which is a smooth map of separable Banach manifolds whose fibers $\pi^{-1}(B)$ are the spaces $\mathcal{V}^k(B)$. Using Lemma 9.1.1, the fact that each map Ψ_B is Fredholm implies that π is also a Fredholm map, so the Sard-Smale theorem implies that the regular values of π form a comeager subset

$$\mathcal{A}_\epsilon^{\text{reg}, k} \subset \mathcal{A}_\epsilon.$$

The intersection

$$\mathcal{A}_\epsilon^{\text{reg}} := \bigcap_{k \in \mathbb{N}} \mathcal{A}_\epsilon^{\text{reg}, k}$$

is then another comeager subset of \mathcal{A}_ϵ , with the property that for each $B \in \mathcal{A}_\epsilon^{\text{reg}}$ and every $k \in \mathbb{N}$ and $(s, \lambda) \in \mathcal{V}^k(B)$, $d\Psi_B(s, \lambda)$ is (by Lemma 9.1.1) surjective. As was observed previously, this is impossible for dimensional reasons if $k \geq 2$, implying that $\mathcal{V}^k(B)$ is then empty.

To find perturbations that also achieve the transversality condition, we use a similar argument: define for each $B \in \mathcal{A}_\epsilon$ the subset

$$\mathcal{V}^0(B) = \{s \in (-1, 1) \mid \dim_{\mathbb{R}} \ker \mathbf{A}_s^B = 1\},$$

along with the corresponding universal set

$$\mathcal{V}^0 = \{(s, B) \in (-1, 1) \times \mathcal{A}_\epsilon \mid s \in \mathcal{V}^0(B)\}.$$

A neighborhood of any (s_0, B_0) in \mathcal{V}^0 is then the zero set of a smooth map of the form

$$\Psi(s, B) = \Phi(\mathbf{A}_s^B) \in \text{End}_{\mathbb{R}}^{\text{sym}}(\ker \mathbf{A}_{s_0}^{B_0}),$$

defined for all $(s, B) \in (-1, 1) \times \mathcal{A}_\epsilon$ close enough to (s_0, B_0) . For a fixed $B \in \mathcal{A}_\epsilon$ near B_0 and $s_1 \in \mathcal{V}^0(B)$ near s_0 , a neighborhood of s_1 in $\mathcal{V}^0(B)$ is then the zero set of $\Psi_B(s) := \Psi(s, B)$, and the intersection of the path $s \mapsto \mathbf{A}_s^B \in \text{Fred}_{\mathbb{R}}^{\text{sym}}(\mathcal{D}, \mathcal{H}, \mathbf{T}_{\text{ref}})$ with $\text{Fred}_{\mathbb{R}}^{\text{sym}, 1}(\mathcal{D}, \mathcal{H}, \mathbf{T}_{\text{ref}})$ at $s = s_1$ is transverse if and only if

$$d\Psi_B(s_1) : \mathbb{R} \rightarrow \text{End}_{\mathbb{R}}^{\text{sym}}(\ker \mathbf{A}_{s_0}^{B_0})$$

is surjective. At (s_0, B_0) , the partial derivative of Ψ with respect to B is again the same operator

$$\mathbf{L} = D_2\Psi(s_0, B_0) : \mathcal{A}_\epsilon \rightarrow \text{End}_{\mathbb{R}}^{\text{sym}}(\ker \mathbf{A}_{s_0}^{B_0})$$

as in (C.1), which we've already seen is surjective due to Lemma C.0.1. Thus one can apply the Sard-Smale theorem to the projection

$$\mathcal{V}^0 \rightarrow \mathcal{A}_\epsilon : (s, B) \mapsto B,$$

obtaining a comeager subset $\mathcal{A}_\epsilon^{\text{reg},0} \subset \mathcal{A}_\epsilon$ such that all paths $\mathbf{A}_s + B(s, \cdot)$ for $B \in \mathcal{A}_\epsilon^{\text{reg},0}$ satisfy the required transversality condition. The comeager subset $\mathcal{A}_\epsilon^{\text{reg},0} \cap \mathcal{A}_\epsilon^{\text{reg}} \subset \mathcal{A}_\epsilon$ thus consists of perturbed families of operators for which all desired conditions are satisfied, and it contains a sequence converging in the C^∞ -topology to 0. This concludes the proof of Lemma 3.4.3.

Bibliography

- [AS19] A. Abbondandolo and F. Schlenk, *Floer homologies, with applications*, Jahresber. Dtsch. Math.-Ver. **121** (2019), no. 3, 155–238.
- [AF03] R. A. Adams and J. J. F. Fournier, *Sobolev spaces*, 2nd ed., Pure and Applied Mathematics (Amsterdam), vol. 140, Elsevier/Academic Press, Amsterdam, 2003.
- [ABW10] P. Albers, B. Bramham, and C. Wendl, *On nonseparating contact hypersurfaces in symplectic 4-manifolds*, Algebr. Geom. Topol. **10** (2010), no. 2, 697–737.
- [AD14] M. Audin and M. Damian, *Morse theory and Floer homology*, Universitext, Springer, London; EDP Sciences, Les Ulis, 2014. Translated from the 2010 French original by Reinie Ern e.
- [Avd21] R. Avdek, *Liouville hypersurfaces and connect sum cobordisms*, J. Symplectic Geom. **19** (2021), no. 4, 865–957.
- [CE20] J. Conway and J. B. Etnyre, *Contact surgery and symplectic caps*, Bull. Lond. Math. Soc. **52** (2020), no. 2, 379–394.
- [DK90] S. K. Donaldson and P. B. Kronheimer, *The geometry of four-manifolds*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, New York, 1990. Oxford Science Publications.
- [Eli89] Y. Eliashberg, *Classification of overtwisted contact structures on 3-manifolds*, Invent. Math. **98** (1989), no. 3, 623–637.
- [EGH00] Y. Eliashberg, A. Givental, and H. Hofer, *Introduction to symplectic field theory*, Geom. Funct. Anal., Special Volume (2000), 560–673.
- [EH02] J. B. Etnyre and K. Honda, *On symplectic cobordisms*, Math. Ann. **323** (2002), no. 1, 31–39.
- [Flo88a] A. Floer, *Morse theory for Lagrangian intersections*, J. Differential Geom. **28** (1988), no. 3, 513–547.
- [Flo88b] ———, *The unregularized gradient flow of the symplectic action*, Comm. Pure Appl. Math. **41** (1988), no. 6, 775–813.
- [Flo88c] ———, *An instanton-invariant for 3-manifolds*, Comm. Math. Phys. **118** (1988), no. 2, 215–240.
- [FOOO09] K. Fukaya, Y.-G. Oh, H. Ohta, and K. Ono, *Lagrangian intersection Floer theory: anomaly and obstruction. Part I*, AMS/IP Studies in Advanced Mathematics, vol. 46, American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2009.
- [Gei08] H. Geiges, *An introduction to contact topology*, Cambridge Studies in Advanced Mathematics, vol. 109, Cambridge University Press, Cambridge, 2008.
- [Gro85] M. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), no. 2, 307–347.
- [Hof93] H. Hofer, *Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjecture in dimension three*, Invent. Math. **114** (1993), no. 3, 515–563.
- [Hof06] H. Hofer, *A general Fredholm theory and applications*, Current developments in mathematics, 2004, Int. Press, Somerville, MA, 2006, pp. 1–71.
- [HS95] H. Hofer and D. A. Salamon, *Floer homology and Novikov rings*, The Floer memorial volume, Progr. Math., vol. 133, Birkh user, Basel, 1995, pp. 483–524.

- [HZ94] H. Hofer and E. Zehnder, *Symplectic invariants and Hamiltonian dynamics*, Birkhäuser Verlag, Basel, 1994.
- [Lan93] S. Lang, *Real and functional analysis*, 3rd ed., Springer-Verlag, New York, 1993.
- [Lan99] ———, *Fundamentals of differential geometry*, Springer-Verlag, New York, 1999.
- [LW11] J. Latschev and C. Wendl, *Algebraic torsion in contact manifolds*, *Geom. Funct. Anal.* **21** (2011), no. 5, 1144–1195. With an appendix by Michael Hutchings.
- [Laz20] O. Lazarev, *Maximal contact and symplectic structures*, *J. Topol.* **13** (2020), no. 3, 1058–1083.
- [LL01] E. H. Lieb and M. Loss, *Analysis*, 2nd ed., Graduate Studies in Mathematics, vol. 14, American Mathematical Society, Providence, RI, 2001.
- [Lut77] R. Lutz, *Structures de contact sur les fibrés principaux en cercles de dimension trois*, *Ann. Inst. Fourier (Grenoble)* **27** (1977), no. 3, ix, 1–15 (French, with English summary).
- [MS12] D. McDuff and D. Salamon, *J-holomorphic curves and symplectic topology*, 2nd ed., American Mathematical Society Colloquium Publications, vol. 52, American Mathematical Society, Providence, RI, 2012.
- [MS17] ———, *Introduction to symplectic topology*, 3rd ed., Oxford University Press, 2017.
- [Rab78] P. H. Rabinowitz, *Periodic solutions of Hamiltonian systems*, *Comm. Pure Appl. Math.* **31** (1978), no. 2, 157–184.
- [Rob70] R. C. Robinson, *A global approximation theorem for Hamiltonian systems*, *Global Analysis (Proc. Sympos. Pure Math., Vol. XIV, Berkeley, Calif., 1968)*, Amer. Math. Soc., Providence, R.I., 1970, pp. 233–243.
- [RT95] Y. Ruan and G. Tian, *A mathematical theory of quantum cohomology*, *J. Differential Geom.* **42** (1995), no. 2, 259–367.
- [RT97] ———, *Higher genus symplectic invariants and sigma models coupled with gravity*, *Invent. Math.* **130** (1997), 455–516.
- [Sal99] D. Salamon, *Lectures on Floer homology*, *Symplectic geometry and topology (Park City, UT, 1997)*, IAS/Park City Math. Ser., vol. 7, Amer. Math. Soc., Providence, RI, 1999, pp. 143–229.
- [Sch93] M. Schwarz, *Morse homology*, *Progress in Mathematics*, vol. 111, Birkhäuser Verlag, Basel, 1993.
- [Sei08] P. Seidel, *Fukaya categories and Picard-Lefschetz theory*, *Zurich Lectures in Advanced Mathematics*, European Mathematical Society (EMS), Zürich, 2008.
- [Wei78] A. Weinstein, *Periodic orbits for convex Hamiltonian systems*, *Ann. of Math. (2)* **108** (1978), no. 3, 507–518.
- [Wen13] C. Wendl, *Non-exact symplectic cobordisms between contact 3-manifolds*, *J. Differential Geom.* **95** (2013), no. 1, 121–182.
- [Wena] ———, *Signs (or how to annoy a symplectic topologist)*. Blog post, available at <https://symplecticfieldtheorist.wordpress.com/2015/08/23/>.
- [Wenb] ———, *Lectures on holomorphic curves in symplectic and contact geometry*. Version 3.3, http://www.math.hu-berlin.de/~wendl/pub/jhol_bookv33.pdf.
- [Wen18] ———, *Holomorphic curves in low dimensions: from symplectic ruled surfaces to planar contact manifolds*, *Lecture Notes in Mathematics*, vol. 2216, Springer-Verlag, 2018.