

---

Exercise sheet, Theme “Convex Optimization”

**Aufgabe 1** (Properties of convex functions)

Let  $f$  be a real-valued function defined on a convex set  $\mathcal{X} \subset \mathbb{R}^d$ .

- Prove: a convex function  $f$  is always continuous on  $\text{Int}(\mathcal{X})$  (and upper semi-continuous on  $\partial\mathcal{X}$ )
- For this question only,  $f$  is not assumed convex. Prove: if for all  $x \in \mathcal{X}$ ,  $\partial f(x) \neq \emptyset$ , then  $f$  is a convex function.
- Prove: if  $f$  is convex, then for all  $x \in \text{Int}(\mathcal{X})$ ,  $\partial f(x) \neq \emptyset$ . *Hint:* one can use without justification the “supporting hyperplane” theorem: if  $\mathcal{C}$  is a convex set and  $x_0 \in \partial\mathcal{C}$ , then there exists a vector  $w \neq 0$  such that for all  $x \in \mathcal{C}$ ,  $\langle x - x_0, w \rangle \geq 0$ . Use the fact that the epigraph of  $f$  is convex.
- Prove: if  $f$  is convex and differentiable at  $x$ , then  $\nabla f(x) \in \partial f(x)$  and is even its unique element if  $x \in \text{Int}(\mathcal{X})$ .
- Prove:  $f(x^*)$  is a local minimum of  $f$  iff  $f(x^*)$  is a global minimum, iff  $0 \in \partial f(x^*)$ .
- Prove: if  $f$  is convex and differentiable, then  $f(x^*)$  is minimum of  $f$  iff for all  $y \in \mathcal{X}$  it holds  $\langle \nabla f(x^*), y - x^* \rangle \geq 0$ .

**Aufgabe 2** (Brunn-Minkowski inequality)

In this exercise we will prove a fundamental theorem which will be used in the next one. Let the ambient space be  $\mathbb{R}^d$  and denote for two measurable sets  $A, B$  the “Minkowski sum” of  $A$  and  $B$  as

$$A + B = \{a + b, a \in A, b \in B\}.$$

The Brunn-Minkowski inequality states the following: for any measurable  $A, B$  of finite volume and  $\lambda \in [0, 1]$  such that  $(1 - \lambda)A + \lambda B$  is measurable, it holds

$$\mathcal{V}_d((1 - \lambda)A + \lambda B)^{\frac{1}{d}} \geq (1 - \lambda)\mathcal{V}_d(A)^{\frac{1}{d}} + \lambda\mathcal{V}_d(B)^{\frac{1}{d}},$$

where  $\mathcal{V}_d$  denotes volume ( $d$ -dimensional Lebesgue measure). In this sense the function  $A \mapsto \mathcal{V}_d(A)^{\frac{1}{d}}$  is “concave”.

Below we call “cuboid” a  $d$ -dimensional rectangular (axis-aligned) box (=cartesian product of intervals along each coordinate).

- Prove the inequality when both  $A$  and  $B$  are cuboids.
- Prove the inequality when  $A$  and  $B$  are disjoint finite unions of cuboids, by recursion on the number of boxes.  
*Hint:* assume for example that  $A$  is a disjoint union of at least two cuboids and prove that there exists at least an axis-aligned hyperplane  $H$  that separates two cuboids of  $A$ . (*Hadwiger-Ohmann cut*). Apply recursively the inequality for  $A$  and  $B$  intersected with both half-spaces defined by  $H$ .
- Conclude by approximating arbitrary measurable sets by finite union of cuboids.

**Aufgabe 3** (Grünbaum's Lemma)

We proceed to proving the following property which was used in the lecture: if  $K$  is a compact convex set of  $\mathbb{R}^d$ , and  $H$  a hyperplane going through the center of gravity of  $K$ , then the intersection of  $K$  with either half-space defined by  $H$  has volume at least  $\frac{1}{2}\mathcal{V}_d(K)$ .

Without loss of generality, we assume  $K$  has center of gravity at the origin and  $H = \{x : x_1 = 0\}$ . For any set  $A$ , we denote  $A_t := A \cap \{x : x_1 = t\}$ ;  $A_+ := A \cap \{x : x_1 \geq 0\}$  and  $A_- := A \cap \{x : x_1 \leq 0\}$ .

- a) Construct a “symmetrized” version  $K'$  of  $K$  (“Schwarzsche Abrundung”) as follows. For any  $t \in \mathbb{R}$ ,  $K'_t$  is the  $(d-1)$ -dimensional ball  $B(0, r_t)$  with  $r_t$  chosen so that  $\mathcal{V}_{d-1}(B(0, r_t)) = \mathcal{V}_{d-1}(K_t)$ .

Prove that  $K'$  is convex. (*Hint*: use the Brunn-Minkowski inequality to establish that  $t \mapsto r_t$  is concave.) Prove that  $\mathcal{V}_d(K'_+) = \mathcal{V}_d(K_+)$  and  $\mathcal{V}_d(K'_-) = \mathcal{V}_d(K_-)$ .

- b) Now consider a second transformation by “conification” of  $K'$ . Consider a cone  $C$  defined as follows:  $C_t$  is the  $(d-1)$ -dimensional ball  $B(0, r'_t)$  with  $r'_t = (r_0 - \alpha t)\mathbf{1}\{r_0\alpha^{-1} \geq t \geq t_-\}$ , with  $\alpha$  chosen so that  $\mathcal{V}_d(C_+) = \mathcal{V}_d(K'_+)$  and  $t_-$  chosen so that  $\mathcal{V}_d(C_-) = \mathcal{V}_d(K'_-)$  (note that  $C_0 = K'_0$  by construction).

Prove that the center of gravity of  $C$  must have nonnegative first coordinate  $g_C$  (and zero other coordinates)

- c) Deduce from the previous question that the cone is a worst-case situation. Compute the position of  $g_C$  of the center of gravity of a cone of height  $h$  and its volume, and conclude.