

KOSZUL MODULES AND GREEN'S CONJECTURE

MARIAN APRODU, GAVRIL FARKAS, ȘTEFAN PAPADIMA[†], CLAUDIU RAICU, AND JERZY WEYMAN

ABSTRACT. We prove a strong vanishing result for finite length Koszul modules, and use it to derive Green's conjecture for every g -cuspidal rational curve over an algebraically closed field \mathbf{k} , with $\text{char}(\mathbf{k}) = 0$ or $\text{char}(\mathbf{k}) \geq \frac{g+2}{2}$. As a consequence, we deduce that the general canonical curve of genus g satisfies Green's conjecture in this range. Our results are new in positive characteristic, whereas in characteristic zero they provide a different proof for theorems first obtained in two landmark papers by Voisin. Our strategy involves establishing two key results of independent interest: (1) we describe an explicit, characteristic-independent version of Hermite reciprocity for \mathfrak{sl}_2 -representations; (2) we completely characterize, in arbitrary characteristics, the (non-)vanishing behavior of the syzygies of the tangential variety to a rational normal curve.

1. INTRODUCTION

Formulated in 1984, Green's Conjecture [13, Conjecture 5.1] predicts that one can recognize in a precise way the intrinsic complexity of a smooth algebraic curve from the syzygies of its canonical embedding. If $C \hookrightarrow \mathbb{P}^{g-1}$ is a non-hyperelliptic canonically embedded curve of genus g , we denote by $K_{i,j}(C, \omega_C)$ the Koszul cohomology group of i -th syzygies of weight j . Green's Conjecture predicts the equivalence

$$K_{i,1}(C, \omega_C) = 0 \iff i \geq g - \text{Cliff}(C) - 1,$$

where $\text{Cliff}(C)$ is the Clifford index of C . Equivalently, $K_{i,2}(C, \omega_C) = 0$ if and only if $i < \text{Cliff}(C)$. Although for arbitrary curves the conjecture remains wide open, Green's Conjecture for a *general* curve of every genus has been resolved using geometric methods in two landmark papers by Voisin [28, 29]. In this case one has $\text{Cliff}(C) = \lfloor \frac{g-1}{2} \rfloor$ and Green's Conjecture reduces to a single vanishing statement

$$(1) \quad K_{\lfloor \frac{g-1}{2} \rfloor, 1}(C, \omega_C) = 0.$$

More direct approaches have been proposed over the years to solve the Generic Green Conjecture (even prior to Voisin's papers), but none has been so far brought to fruition. One of them is described in Eisenbud's paper [7, Section 3.I] and relies on degeneration and proving the vanishing (1) for a general canonically embedded g -cuspidal rational curve. This approach reduces the Generic Green Conjecture to a rather impenetrable looking problem in \mathfrak{sl}_2 -representation theory. Using a novel perspective on these questions essentially inspired from topology, and which we have already put to work in [2] to attack questions in geometric group theory, we complete this approach by reducing the Generic Green Conjecture to a strong vanishing result for finite length Koszul modules. In characteristic zero, this gives a new, relatively short proof of Green's Conjecture for generic curves of any genus. Compared to Voisin's approach, we do not use the geometry of $K3$ surfaces and Hilbert schemes of 0-dimensional subschemes, but instead we view the Generic Green Conjecture as a question on the geometry of the Grassmannian of lines, which we then solve by using general vanishing results for homogeneous bundles (Bott's Theorem). This approach also works in positive characteristic, as we shall explain.

Let $\mathcal{T} \subseteq \mathbf{P}^g$ denote the tangential variety of the degree g rational normal curve $\Gamma \subseteq \mathbf{P}^g$, over some algebraically closed field \mathbf{k} . A general hyperplane section C of \mathcal{T} is a canonically embedded g -cuspidal

[†]Ștefan Papadima passed away on January 10, 2018.

rational curve, the cusps corresponding to the points of intersection of C with Γ . Using a form of the Lefschetz hyperplane principle [3, Lemma 2.19], one has the isomorphism

$$K_{i,j}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) \cong K_{i,j}(C, \omega_C)$$

for all i, j . In characteristic zero, it is a consequence of the theory of limit linear series of Eisenbud and Harris [9] that all g -cuspidal rational curves verify the Brill–Noether Theorem, and in particular they satisfy $\text{Cliff}(C) = \lfloor \frac{g-1}{2} \rfloor$. In this way, Green’s Conjecture for C is turned into a concrete question about the syzygies of \mathcal{T} . We prove the following:

Theorem 1.1. *If $g \geq 3$ and $\text{char}(\mathbf{k}) = 0$ or $\text{char}(\mathbf{k}) \geq \frac{g+2}{2}$, then $K_{\lfloor \frac{g}{2} \rfloor, 1}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) = 0$.*

As a consequence of this result we give a new proof of Voisin’s Theorem [28, 29] on the Generic Green Conjecture in characteristic zero, as well as an effective bound, in terms of the genus of the curve for the characteristics in which the Generic Green Conjecture holds.

Theorem 1.2. *Let $g \geq 3$ and suppose that $\text{char}(\mathbf{k}) = 0$ or $\text{char}(\mathbf{k}) \geq \frac{g+2}{2}$. For a general curve C of genus g we have that $K_{\lfloor \frac{g}{2} \rfloor, 1}(C, \omega_C) = 0$, and C verifies Green’s Conjecture.*

Early on, Schreyer [25] observed that Green’s Conjecture fails for some positive characteristics, for instance for genus 7 and characteristics 2, or genus 9 and characteristics 3. More recently, Eisenbud and Schreyer [10, Conjecture 0.1] predicted that Green’s Conjecture should hold for general curves of genus g whenever $\text{char}(\mathbf{k}) \geq \frac{g-1}{2}$. A version of Green’s Conjecture involving arbitrary smooth curves in positive characteristic has been put forward recently by Bopp and Schreyer [4].

Our Theorem 1.2 gives an almost complete answer to the Eisenbud–Schreyer conjecture and leaves at most one characteristic open. If $2 \leq \text{char}(\mathbf{k}) \leq \frac{g+1}{2}$ then the tangential variety \mathcal{T} is contained in a rational normal scroll of codimension at least $\lfloor \frac{g}{2} \rfloor$. Therefore $K_{\lfloor \frac{g}{2} \rfloor, 1}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) \neq 0$ (see Remark 5.17) and the same non-vanishing holds for the general hyperplane section C . This shows that cuspidal curves cannot be used to prove the Eisenbud–Schreyer conjecture when $\frac{g-1}{2} \leq \text{char}(\mathbf{k}) \leq \frac{g+1}{2}$. One key ingredient in our proof of Theorem 1.1 comes from a solid understanding of the Hilbert function of finite length Koszul modules, as explained next.

Koszul modules. Suppose that V is an n -dimensional \mathbf{k} -vector space and fix a subspace $K \subseteq \bigwedge^2 V$ with $\dim(K) = m$. We denote by $S := \text{Sym } V$ the symmetric algebra over V and consider the Koszul complex resolving the residue field \mathbf{k} :

$$\cdots \longrightarrow \bigwedge^3 V \otimes S \xrightarrow{\delta_3} \bigwedge^2 V \otimes S \xrightarrow{\delta_2} V \otimes S \xrightarrow{\delta_1} S.$$

Truncating this complex to the last three terms, and restricting δ_2 along the inclusion $\iota : K \hookrightarrow \bigwedge^2 V$ we obtain a 3-term complex

$$(2) \quad K \otimes S \xrightarrow{\delta_2|_{K \otimes S}} V \otimes S \xrightarrow{\delta_1} S.$$

Following [23, §2] as well as [2], we define the *Koszul module* associated to the pair (V, K) to be the middle homology of the complex (2). We make the convention that K is placed in degree zero, so that $W(V, K)$ is a graded S -module generated in degree zero. Using the exactness of the Koszul differentials $\delta_{i,q} : \bigwedge^i V \otimes \text{Sym}^q V \rightarrow \bigwedge^{i-1} V \otimes \text{Sym}^{q+1} V$, we have the following alternative identification of the q -th graded piece of the Koszul module $W_q(V, K) = \text{Ker}(\delta_{1,q+1})/\delta_{2,q}(K \otimes \text{Sym}^q V)$:

$$(3) \quad W_q(V, K) = \text{Coker} \left(\bigwedge^3 V \otimes \text{Sym}^{q-1} V \longrightarrow \frac{\bigwedge^2 V}{K} \otimes \text{Sym}^q V \right) \cong \frac{\bigwedge^2 V \otimes \text{Sym}^q V}{K \otimes \text{Sym}^q V + \text{Ker}(\delta_{2,q})}.$$

The formation of the Koszul module $W(V, K)$ is natural in the following sense. An inclusion $K \subseteq K'$ induces a surjective morphism of graded S -modules

$$(4) \quad W(V, K) \twoheadrightarrow W(V, K'),$$

that is, *bigger* subspaces $K \subseteq \bigwedge^2 V$ correspond to *smaller* Koszul modules. For instance, we have that $W(V, K) = 0$ if and only if $K = \bigwedge^2 V$. We shall be interested more generally in studying Koszul modules that are finite dimensional as \mathbf{k} -vector spaces, that is, those that satisfy $W_q(V, K) = 0$ for $q \gg 0$. Since $W(V, K)$ is generated in degree zero, the vanishing $W_q(V, K) = 0$ for some $q \geq 0$ implies that $W_{q'}(V, K) = 0$ for all $q' \geq q$.

We write $\iota^\vee : \bigwedge^2 V^\vee \twoheadrightarrow K^\vee$ for the dual to the inclusion ι , and let $K^\perp := \text{Ker}(\iota^\vee) \subseteq \bigwedge^2 V^\vee$. It is shown in [23, Lemma 2.4] that the set-theoretic support of $W(V, K)$ in the affine space V^\vee is given by the *resonance variety* $\mathcal{R}(V, K)$, defined as

$$(5) \quad \mathcal{R}(V, K) := \left\{ a \in V^\vee : \text{there exists } b \in V^\vee \text{ such that } a \wedge b \in K^\perp \setminus \{0\} \right\} \cup \{0\}.$$

In particular, $W(V, K)$ has finite length if and only if $\mathcal{R}(V, K) = \{0\}$. In view of (5), this last condition is equivalent to the fact that the linear subspace $\mathbb{P}K^\perp \subseteq \mathbb{P}(\bigwedge^2 V^\vee)$ is disjoint from the Grassmann variety

$$\mathbf{G} := \text{Gr}_2(V^\vee)$$

in its Plücker embedding, which can happen only when $m = \text{codim}(\mathbb{P}K^\perp) > \dim(\mathbf{G}) = 2n - 4$. Summarizing, we have the following equivalences:

$$(6) \quad \mathbb{P}(K^\perp) \cap \mathbf{G} = \emptyset \iff \mathcal{R}(V, K) = \{0\} \iff \dim_{\mathbf{k}} W(V, K) < \infty.$$

Moreover, if the equivalent statements in (6) hold, then $m \geq 2n - 3$.

We provide a sharp effective bound for the vanishing of the graded components of a finite length Koszul module, extending the main result from [2] to (sufficiently) positive characteristics. Recalling that $n = \dim_{\mathbf{k}}(V)$, we prove the following:

Theorem 1.3. *Suppose that $n \geq 3$. If $\text{char}(\mathbf{k}) = 0$ or $\text{char}(\mathbf{k}) \geq n - 2$, then we have the equivalence*

$$(7) \quad \mathcal{R}(V, K) = \{0\} \iff W_q(V, K) = 0 \text{ for } q \geq n - 3.$$

Experiments in small characteristics suggest that the assumption $\text{char}(\mathbf{k}) \geq n - 2$ is probably necessary in order for (7) to hold, but we do not know this in general. As the next theorem shows, the vanishing range $q \geq n - 3$ is on the other hand optimal, since $W_{n-4}(V, K) \neq 0$ when $m = 2n - 3$ and $n \geq 4$.

Theorem 1.4. *Suppose $\text{char}(\mathbf{k}) = 0$ or $\text{char}(\mathbf{k}) \geq n - 2$, and fix a subspace $K \subseteq \bigwedge^2 V$. If $\mathcal{R}(V, K) = \{0\}$, then*

$$\dim W_q(V, K) \leq \binom{n+q-1}{q} \frac{(n-2)(n-q-3)}{q+2} \quad \text{for } q = 0, \dots, n-4.$$

Moreover, equality holds for all q if $\dim(K) = 2n - 3$.

The study of resonance varieties (and of the associated cohomology jumping loci) in Hodge theory has been initiated by Green and Lazarsfeld [14], and it has been actively pursued in the theory of hyperplane arrangements and topology. We refer to [6], [23] and the references therein for an overview, and for the

connection between topological invariants of groups and Koszul modules. Theorem 1.4 yields upper bounds for these invariants in a purely algebraic context, as explained in detail in [2].

The Cayley-Chow form of the Grassmannian of lines. Theorem 1.3 can be reformulated geometrically as follows. Recall that one defines the *Cayley–Chow form* of the Grassmannian $\mathbf{G} \subseteq \mathbb{P}(\wedge^2 V^\vee)$ in its Plücker embedding as the locus of linear subspaces of codimension $(2n - 3)$ in $\mathbb{P}(\wedge^2 V^\vee)$ that meet \mathbf{G} . In view of (6), this can be seen equivalently as the locus of $(2n - 3)$ -dimensional subspaces in $\wedge^2 V$ for which $W(V, K)$ has non-zero resonance, that is, as

$$\mathcal{C}\text{how}(\mathbf{G}) := \left\{ K \in \text{Gr}_{2n-3}(\wedge^2 V) : \mathbb{P}(K^\perp) \cap \mathbf{G} \neq \emptyset \right\}.$$

Note that $\mathcal{C}\text{how}(\mathbf{G})$ is a divisor in $\text{Gr}_{2n-3}(\wedge^2 V)$ of degree equal to $\deg(\mathbf{G})$, which is known to be computed by the Catalan number $\frac{1}{n-1} \cdot \binom{2n-4}{n-2}$. In view of the description (3) for the space $W_{n-3}(V, K)$, an equivalent reformulation of Theorem 1.3 (in the case when $\dim(K) = 2n - 3$) is the following:

Theorem 1.5. *Suppose that $\text{char}(\mathbf{k}) = 0$ or $\text{char}(\mathbf{k}) \geq n - 2$. The Cayley–Chow form $\mathcal{C}\text{how}(\mathbf{G})$ consists set-theoretically of those subspaces $K \subseteq \wedge^2 V$ with $\dim(K) = 2n - 3$ satisfying*

$$\left(K \otimes \text{Sym}^{n-3}(V) \right) \cap \text{Ker}(\delta_{2,n-3}) \neq 0.$$

In other words, the Cayley–Chow form of \mathbf{G} is the pull-back of the divisorial Schubert cycle defined by the Koszul space $\text{Ker}(\delta_{2,n-3} : \wedge^2 V \otimes \text{Sym}^{n-3} V \rightarrow V \otimes \text{Sym}^{n-2} V)$ under the map

$$\text{Gr}_{2n-3}(\wedge^2 V) \rightarrow \text{Gr}_{(2n-3)\binom{2n-4}{n-1}}(\wedge^2 V \otimes \text{Sym}^{n-3} V), \quad K \mapsto K \otimes \text{Sym}^{n-3} V.$$

Returning to the Generic Green Conjecture, our new approach and the proof of Theorem 1.1 is based on exhibiting a close relationship between the syzygies of the tangential variety \mathcal{T} and the graded components of a specific finite length Koszul module, whose study was initiated in [7, Section 3.I.B], and which is called a *Weyman module*. This relationship lies at the heart of this paper, and we explain it in the remaining part of the Introduction.

The syzygies of the tangential variety of the rational normal curve. We assume for the rest of the Introduction that $\text{char}(\mathbf{k}) \neq 2$. We prove in Theorem 5.1 that the homogeneous coordinate ring of \mathcal{T} is Gorenstein, with Castelnuovo–Mumford regularity 3. In other words, the dimensions $b_{i,j}$ of the Koszul cohomology groups $K_{i,j}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1))$ are encoded in a table having the following shape:

$$(8) \quad \begin{array}{c|ccccccc} & 0 & 1 & 2 & \cdots & g-4 & g-3 & g-2 \\ \hline 0 & 1 & - & - & \cdots & - & - & - \\ 1 & - & b_{1,1} & b_{2,1} & \cdots & b_{g-4,1} & b_{g-3,1} & - \\ 2 & - & b_{1,2} & b_{2,2} & \cdots & b_{g-4,2} & b_{g-3,2} & - \\ 3 & - & - & - & \cdots & - & - & 1 \end{array}$$

where a dash in position (i, j) indicates the vanishing of the corresponding Koszul cohomology group. We prove in Theorem 5.2 that:

Theorem 1.6. *If $p = \text{char}(\mathbf{k})$ satisfies $p = 0$ or $p \geq \frac{g+2}{2}$, then*

$$b_{i,2} \neq 0 \quad \text{if and only if} \quad \frac{g-2}{2} \leq i \leq g-3.$$

The Gorenstein property implies that $b_{i,1} = b_{g-2-i,2}$ for all i , so the vanishing statement in Theorem 1.1 is equivalent to $b_{g-2-\lfloor \frac{g}{2} \rfloor, 2} = 0$, which follows from Theorem 1.6. In Theorem 5.2 we also give a similar characterization for the (non-)vanishing of $b_{i,2}$ when $3 \leq p \leq \frac{g+1}{2}$, and in Section 5.9 we consider $p = 2$. To prove Theorem 5.2, we realize the groups $K_{i,2}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1))$ as graded components of Weyman modules as explained next, and then apply Theorem 1.3 and the explicit description (5) of the set-theoretic support of a Koszul module.

Weyman modules. We fix a \mathbf{k} -vector space U of dimension two, and let $\mathbb{P}U$ denote the projective space of one-dimensional subspaces of U . For each $d \geq 0$ we have $H^0(\mathbb{P}U, \mathcal{O}_{\mathbb{P}U}(d)) = \text{Sym}^d(U^\vee)$, and we define the d -th divided power of U by

$$(9) \quad D^d U := \left(\text{Sym}^d(U^\vee) \right)^\vee.$$

If $\text{char}(\mathbf{k}) = 0$, or more generally, if all the binomial coefficients $\binom{d}{i}$ are invertible in \mathbf{k} , then there exists a natural isomorphism $D^d U \cong \text{Sym}^d U$, as explained in Section 3.1. Readers interested only in characteristic zero can use this identification throughout the paper.

For each $d \geq 1$ we consider the *Gaussian–Wahl map* associated with the line bundle $\mathcal{O}_{\mathbb{P}U}(d)$ (see [30]):

$$\mu_1 : \bigwedge^2 H^0(\mathbb{P}U, \mathcal{O}_{\mathbb{P}U}(d)) \longrightarrow H^0(\mathbb{P}U, \omega_{\mathbb{P}U} \otimes \mathcal{O}_{\mathbb{P}U}(2d)).$$

We choose a basis $(1, y)$ for U^\vee , so that $\text{Sym}^d(U^\vee)$ can be identified with the space of polynomials of degree at most d in y . Making the identification $\omega_{\mathbb{P}U} \cong \mathcal{O}_{\mathbb{P}U}(-2)$, we can rewrite μ_1 explicitly as

$$(10) \quad \mu_1 : \bigwedge^2 \text{Sym}^d(U^\vee) \longrightarrow \text{Sym}^{2d-2}(U^\vee), \quad \mu_1(y^i \wedge y^j) = (i-j) \cdot y^{i+j-1} \text{ for } 0 \leq i, j \leq d.$$

In characteristic zero (or sufficiently large), we have the Clebsch–Gordan decomposition

$$\bigwedge^2 \text{Sym}^d(U^\vee) = \bigoplus_{j=0}^{\lfloor \frac{d-1}{2} \rfloor} \text{Sym}^{2d-2-4j}(U^\vee),$$

and the map μ_1 is simply the projection onto the factor $\text{Sym}^{2d-2}(U^\vee)$ (see [12, Exercise 11.30]).

Under the assumption that $\text{char}(\mathbf{k}) \neq 2$, the Wahl map (10) is surjective. Dualizing it, using (9), and writing $V^{(d)} = D^d U$ and $K^{(d)} = D^{2d-2} U$, we get an inclusion $\Delta_1 : K^{(d)} \hookrightarrow \bigwedge^2 V^{(d)}$, and define

$$W^{(d)} := W(V^{(d)}, K^{(d)}).$$

Following [7], we call $W^{(d)}$ a *Weyman module*. The key observation is then the following (Theorem 5.3):

Theorem 1.7. *If $\text{char}(\mathbf{k}) \neq 2$, then for each $i = 1, \dots, g-3$ we have a natural identification*

$$K_{i,2}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) = W_{g-3-i}^{(i+2)}.$$

This result reveals a peculiar property of the Weyman module $W^{(i+2)}$, for its graded pieces are isomorphic to the Koszul cohomology groups $K_{i,2}$ of generic curves of *varying* genera! The analysis of the Koszul cohomology groups $K_{i,j}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1))$ is the most technical part of our paper, and is explained in detail in Section 5. It relies on applications of the Kempf–Weyman technique for constructing syzygies, and a summary that avoids technicalities is presented in what follows. Some of the constructions that we use in Section 5 were already outlined in [7], but a formal verification of their correctness requires

quite a bit of care. An essential tool we employ in our arguments is an *explicit, characteristic-free version of Hermite reciprocity* which is, as far as we know, new. It consists of an explicit, natural isomorphism

$$\mathrm{Sym}^d(D^i U) \longrightarrow \bigwedge^i (\mathrm{Sym}^{d+i-1} U)$$

for all $d, i \geq 1$, which is described in terms of the change of bases between elementary symmetric polynomials and Schur polynomials. This isomorphism is explained in Section 3.4.

Syzygies of the tangent developable \mathcal{T} : the geometric intuition. We conclude our Introduction by discussing the maps and spaces involved in the identification in Theorem 1.7. We consider the diagram

$$(11) \quad \begin{array}{ccc} D^{2i+2} U \otimes \mathrm{Sym}^{g-3-i}(D^{i+2} U) & \longrightarrow & D^{2i+2} U \otimes \bigwedge^{i+2} \mathrm{Sym}^{g-2} U \\ \delta_2 \downarrow & & \downarrow \tilde{\delta}_2 \\ D^{i+2} U \otimes \mathrm{Sym}^{g-2-i}(D^{i+2} U) & \longrightarrow & D^{i+2} U \otimes \bigwedge^{i+2} \mathrm{Sym}^{g-1} U \\ \delta_1 \downarrow & & \downarrow \tilde{\delta}_1 \\ \mathrm{Sym}^{g-1-i}(D^{i+2} U) & \longrightarrow & \bigwedge^{i+2} \mathrm{Sym}^g U \end{array}$$

where the horizontal arrows are identifications provided by Hermite reciprocity, the left column is the 3-term complex whose homology defines $W_{g-3-i}^{(i+2)}$, while the maps $\tilde{\delta}_1$ and $\tilde{\delta}_2$ are the unique ones making the diagram commute. To prove Theorem 1.7, it suffices to explain why $K_{i,2}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1))$ may be identified with $\mathrm{Ker}(\tilde{\delta}_1)/\mathrm{Im}(\tilde{\delta}_2)$. This is achieved by describing geometrically the groups in the right column of (11).

Recall that $\Gamma \subseteq \mathbf{P}^g$ denotes the rational normal curve of degree g , which we view as the Veronese embedding of $\mathbf{P}^1 := \mathbb{P}(U^\vee)$, where U is a two-dimensional \mathbf{k} -vector space. We identify $\mathcal{O}_{\mathbf{P}^1}(g) = \mathcal{O}_\Gamma(1)$ and $H^0(\Gamma, \mathcal{O}_\Gamma(1)) \cong H^0(\mathbf{P}^g, \mathcal{O}_{\mathbf{P}^g}(1)) \cong \mathrm{Sym}^g U$. We introduce the *jet bundle* $\mathcal{J} := \mathcal{P}(\mathcal{O}_{\mathbf{P}^1}(g))$ of $\mathcal{O}_{\mathbf{P}^1}(g)$, which sits canonically in an exact sequence

$$0 \longrightarrow \omega_{\mathbf{P}^1} \otimes \mathcal{O}_{\mathbf{P}^1}(g) \longrightarrow \mathcal{J} \longrightarrow \mathcal{O}_{\mathbf{P}^1}(g) \longrightarrow 0.$$

We denote by $\mathbf{P}(\mathcal{J}) \rightarrow \mathbf{P}^1$ the corresponding projectivized tangent bundle. The Taylor homomorphism of the jet bundle is a surjective map of sheaves $\mathrm{Sym}^g U \otimes \mathcal{O}_{\mathbf{P}^1} \rightarrow \mathcal{J}$, defined by $f \otimes 1 \mapsto (df, f)$. It gives rise to a morphism $\tau : \mathbf{P}(\mathcal{J}) \rightarrow \mathbf{P}^g$, mapping $\mathbf{P}(\mathcal{J})$ birationally onto \mathcal{T} , so $\tau : \mathbf{P}(\mathcal{J}) \rightarrow \mathcal{T}$ provides a resolution of singularities for the tangent developable. There exists an exact sequence on \mathbf{P}^g

$$0 \longrightarrow \mathcal{O}_{\mathcal{T}} \longrightarrow \tau_* \mathcal{O}_{\mathbf{P}(\mathcal{J})} \longrightarrow \omega_\Gamma \rightarrow 0,$$

which induces a long exact sequence in Koszul cohomology

$$\begin{aligned} \cdots \longrightarrow K_{i+1,1}(\mathcal{T}, \tau_* \mathcal{O}_{\mathbf{P}(\mathcal{J})}, \mathcal{O}_{\mathcal{T}}(1)) &\xrightarrow{\zeta} K_{i+1,1}(\Gamma, \omega_\Gamma, \mathcal{O}_\Gamma(1)) \longrightarrow K_{i,2}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) \longrightarrow \\ &\longrightarrow K_{i,2}(\mathcal{T}, \tau_* \mathcal{O}_{\mathbf{P}(\mathcal{J})}, \mathcal{O}_{\mathcal{T}}(1)) \longrightarrow \cdots \end{aligned}$$

The desired identification of $K_{i,2}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) \cong \mathrm{Ker}(\tilde{\delta}_1)/\mathrm{Im}(\tilde{\delta}_2)$ follows if we can prove:

- (1) $K_{i+1,1}(\mathcal{T}, \tau_* \mathcal{O}_{\mathbf{P}(\mathcal{J})}, \mathcal{O}_{\mathcal{T}}(1)) \cong D^{2i+2} U \otimes \bigwedge^{i+2} \mathrm{Sym}^{g-2} U$ and $K_{i,2}(\mathcal{T}, \tau_* \mathcal{O}_{\mathbf{P}(\mathcal{J})}, \mathcal{O}_{\mathcal{T}}(1)) = 0$.
- (2) $K_{i+1,1}(\Gamma, \omega_\Gamma, \mathcal{O}_\Gamma(1)) \cong \mathrm{Ker}(\tilde{\delta}_1)$.
- (3) Under the above isomorphisms, the map ζ can be identified with $\tilde{\delta}_2$.

Step (1) is carried out in Proposition 5.6, where we establish the identifications

$$K_{i+1,1}(\mathcal{T}, \tau_* \mathcal{O}_{\mathbf{P}(\mathcal{J})}, \mathcal{O}_{\mathcal{T}}(1)) \cong H^1(\Gamma, \bigwedge^{i+2} \xi) \cong D^{2i+2} U \otimes \bigwedge^{i+2} \text{Sym}^{g-2} U,$$

with ξ being the kernel of the Taylor homomorphism. We also show that $K_{i,2}(\mathcal{T}, \tau_* \mathcal{O}_{\mathbf{P}(\mathcal{J})}, \mathcal{O}_{\mathcal{T}}(1)) = 0$. Concerning (2), using the fact that the kernel bundle $M_{\mathcal{O}_{\Gamma}(1)}$ is obtained by restricting the tautological subbundle on \mathbf{P}^g to Γ , it follows from [8, Theorem 5.8] that

$$K_{i+1,1}(\Gamma, \omega_{\Gamma}, \mathcal{O}_{\Gamma}(1)) = \text{Ker} \left\{ H^1(\Gamma, \bigwedge^{i+2} M_{\mathcal{O}_{\Gamma}(1)} \otimes \omega_{\Gamma}) \xrightarrow{\alpha} \bigwedge^{i+2} \text{Sym}^g U \otimes H^1(\Gamma, \omega_{\Gamma}) \right\},$$

Since $H^1(\Gamma, \bigwedge^{i+2} M_{\mathcal{O}_{\Gamma}(1)} \otimes \omega_{\Gamma}) \cong D^{i+2} U \otimes \bigwedge^{i+2} \text{Sym}^{g-1} U$ and $H^1(\Gamma, \omega_{\Gamma}) \cong \mathbf{k}$, the maps α and $\tilde{\delta}_1$ have the same source and the same target. It is plausible then that $\alpha = \tilde{\delta}_1$ and that (2) should hold (see Corollary 5.10). A verification of (3) would then conclude Theorem 1.7.

The geometric approach sketched above, however appealing, is difficult to execute in practice, particularly because it requires at each step a very careful bookkeeping of the identifications involved. Instead we take a more algebraic approach in Section 5, which goes beyond the scope of this Introduction, and whose extended summary is included in Section 5.2. The Koszul cohomology calculations sketched above are replaced in our approach by syzygy calculations involving the Kempf–Weyman technique.

Summary. The paper is organized as follows. In Section 2 we discuss Koszul modules of finite length, a characteristic-free instance of Bott's vanishing theorem, and we prove Theorems 1.3 and 1.4. In Section 3 we recall the characteristic-free construction of divided, exterior and symmetric powers, and prove our explicit version of Hermite reciprocity. In Section 4 we summarize some basic results on Koszul cohomology and minimal resolutions, and discuss the Kempf–Weyman technique for constructing syzygies. In Section 5 we analyze the syzygies of the tangential variety to a rational normal curve, and prove Theorems 1.1, 1.6, and 1.7. We end with Section 6 where we use moduli of pseudo-stable curves and Theorem 1.1 to deduce (in suitable characteristics) the Generic Green Conjecture (Theorem 1.2).

Acknowledgment. We acknowledge with thanks the contribution of A. Suciuc. This project, including the companion paper [2], started with the paper [23], and since then we benefited from numerous discussions with him. We warmly thank A. Beauville, L. Ein, D. Eisenbud, B. Klingler, P. Pirola, F.-O. Schreyer and C. Voisin for interesting discussions related to this circle of ideas. We are particularly grateful to R. Lazarsfeld who read an early version of the paper and suggested many improvements which significantly clarified the exposition.

Aprodu was partially supported by the Romanian Ministry of Research and Innovation, CNCS - UEFISCDI, grant PN-III-P4-ID-PCE-2016-0030, within PNCDI III. Farkas was supported by the DFG grant *Syzygien und Moduli*. Raicu was supported by the Alfred P. Sloan Foundation and by the NSF Grant No. 1600765. Weyman was partially supported by the Sidney Professorial Fund and the NSF grant No. 1802067.

2. KOSZUL MODULES IN ALGEBRAIC GEOMETRY

The goal of this section is to extend the main result of [2] to positive characteristic, by establishing Theorem 1.3. Once this theorem is proved, the arguments from [2, Section 3.3] carry over to deduce Theorem 1.4. We prove Theorem 1.3 using a suitable positive characteristic version of Bott vanishing, in conjunction with the hypercohomology spectral sequence that was featured in the work of Voisin on the Green conjecture [28]. We recall the basics on partitions and Schur functors in Section 2.1, and

use them to formulate appropriate versions of Bott's theorem for flag varieties in Section 2.2, and for Grassmannians in Section 2.3. The proof of the vanishing Theorem 1.3 is explained in Section 2.4.

2.1. Partitions and Schur functors. The reference for this part is [31, Chapter 2]. A partition λ is a sequence $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r)$ of non-negative integers, where the *parts* λ_j of λ are assumed to be eventually 0. The *size* of λ is written as $|\lambda| = \lambda_1 + \lambda_2 + \dots + \lambda_r$. For each partition λ we define the *conjugate partition* λ' by letting λ'_j be the number of parts λ_i satisfying $\lambda_i \geq j$. We often omit trailing zeros, and group parts of the same size together by writing (a^b) for the sequence (a, a, \dots, a) , where the entry a is repeated b times. For instance, we write $(4^3, 2^4, 1)$ for $(4, 4, 4, 2, 2, 2, 2, 1, 0, 0, 0, \dots)$.

We let \mathbf{k} be a field, and for each partition λ we consider the *Schur functor* \mathbb{L}_λ on the category of finite dimensional \mathbf{k} -vector spaces, defined as in [31, Section 2.1]: For each vector space V , we construct $\mathbb{L}_\lambda V$ as a suitable quotient

$$\bigwedge^{\lambda_1} V \otimes \bigwedge^{\lambda_2} V \otimes \dots \otimes \bigwedge^{\lambda_r} V \twoheadrightarrow \mathbb{L}_\lambda V,$$

satisfying the following properties:

- If $\lambda_1 > \dim(V)$, then $\mathbb{L}_\lambda(V) = 0$.
- If $\lambda_1 = r$ and $\lambda_i = 0$ for $i > 1$, then $\mathbb{L}_\lambda V = \bigwedge^r V$.
- If $\lambda = (1^d)$, then $\mathbb{L}_\lambda V = \text{Sym}^d V$.

More generally, \mathbb{L}_λ can be applied to any locally free sheaf \mathcal{E} on a variety B over \mathbf{k} , yielding a locally free sheaf $\mathbb{L}_\lambda \mathcal{E}$ on B .

We write $\mathbb{Z}_{\text{dom}}^n$ for the set of *dominant weights* in \mathbb{Z}^n , that is, tuples $\lambda = (\lambda_1, \dots, \lambda_n) \in \mathbb{Z}^n$ satisfying $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$, and note that any partition with $\lambda_{n+1} = 0$ can be identified with a dominant weight in $\mathbb{Z}_{\text{dom}}^n$. Given $\lambda \in \mathbb{Z}_{\text{dom}}^n$, for a locally free sheaf \mathcal{E} of rank n on B we set

$$\mathbb{S}_\lambda \mathcal{E} := \mathbb{L}_{\mu'} \mathcal{E} \otimes (\det \mathcal{E})^{\otimes \lambda_n}$$

where $\mu = (\lambda_1 - \lambda_n, \lambda_2 - \lambda_n, \dots, \lambda_{n-1} - \lambda_n)$, and refer to \mathbb{S}_λ also as a *Schur functor*. We have the identifications $\mathbb{S}_{(d, 0^{n-1})} V = \text{Sym}^d V$ and $\mathbb{S}_{(1^r, 0^{n-r})} V = \bigwedge^r V$ for $d \geq 0$ and $0 \leq r \leq n$.

2.2. Bott's Theorem for flag varieties [31, Ch. 3–4], [20, Sections II.4–II.5]. Let V be an n -dimensional vector space over an algebraically closed field \mathbf{k} . We denote by $\mathbf{F}(V)$ the variety of complete flags

$$V_\bullet : V = V_n \twoheadrightarrow V_{n-1} \twoheadrightarrow \dots \twoheadrightarrow V_1 \twoheadrightarrow V_0 = 0,$$

where V_p is a p -dimensional quotient of V for each $p = 0, \dots, n$. We write $\mathcal{Q}_p(V)$ for the tautological rank p quotient bundle on $\mathbf{F}(V)$, whose fiber over a flag V_\bullet is V_p . We consider the tautological line bundle $\mathcal{L}_p(V)$ on $\mathbf{F}(V)$ defined as the kernel of the quotient map $\mathcal{Q}_p(V) \twoheadrightarrow \mathcal{Q}_{p-1}(V)$. For $\lambda \in \mathbb{Z}^n$ we define

$$\mathcal{L}^\lambda(V) := \mathcal{L}_1(V)^{\otimes \lambda_1} \otimes \mathcal{L}_2(V)^{\otimes \lambda_2} \otimes \dots \otimes \mathcal{L}_n(V)^{\otimes \lambda_n}.$$

More generally, if B is a variety over \mathbf{k} and \mathcal{E} is a locally free sheaf of rank n on B we denote by $\mathbf{F}_B(\mathcal{E})$ the *relative flag variety*, equipped with a natural map

$$\pi : \mathbf{F}_B(\mathcal{E}) \longrightarrow B,$$

such that for every point $b \in B$ the fiber $\pi^{-1}(b)$ is isomorphic to the variety $\mathbf{F}(\mathcal{E}_b)$ of flags on the fiber $\mathcal{E}_b = \mathcal{E} \otimes \mathbf{k}(b)$. We define the tautological bundles $\mathcal{Q}_p(\mathcal{E})$ and $\mathcal{L}^\lambda(\mathcal{E})$ in analogy to the absolute case, and note that we have a tautological sequence of quotient maps on $\mathbf{F}_B(\mathcal{E})$

$$\pi^*(\mathcal{E}) = \mathcal{Q}_n(\mathcal{E}) \twoheadrightarrow \mathcal{Q}_{n-1}(\mathcal{E}) \twoheadrightarrow \dots \twoheadrightarrow \mathcal{Q}_1(\mathcal{E}) \twoheadrightarrow \mathcal{Q}_0(\mathcal{E}) = 0.$$

We need the following characteristic-free version of Bott's Theorem, which is due to Kempf (and often referred to as *Kempf vanishing*) – see [20, Section II.4] and [31, Theorem 4.1.10].

Theorem 2.1. *If \mathcal{E} is a locally free sheaf of rank n on B and $\lambda \in \mathbb{Z}_{\text{dom}}^n$, then*

$$(12) \quad R^k \pi_*(\mathcal{L}^\lambda(\mathcal{E})) = \begin{cases} \mathbb{S}_\lambda \mathcal{E} & \text{if } k = 0; \\ 0 & \text{if } k > 0. \end{cases}$$

In particular, if $B = \text{Spec}(\mathbf{k})$ and $\mathcal{E} = V$, then $\mathcal{L}^\lambda(V)$ has vanishing higher cohomology and

$$H^0(\mathbf{F}(V), \mathcal{L}^\lambda(V)) = \mathbb{S}_\lambda V.$$

We shall be interested more generally in the cohomology of line bundles $\mathcal{L}^\lambda(V)$ when λ is not dominant. To that end we follow the analysis described in [20, Section II.5]. We consider the action of the symmetric group on \mathbb{Z}^n defined by

$$s_i \bullet \lambda := (\lambda_1, \dots, \lambda_{i-1}, \lambda_{i+1} - 1, \lambda_i + 1, \lambda_{i+2}, \dots, \lambda_n),$$

where s_i denotes the transposition $(i, i + 1)$.

Proposition 2.2 ([20, Proposition II.5.4]). *Suppose that V is a \mathbf{k} -vector space with $\dim(V) = n$, consider a weight $\lambda \in \mathbb{Z}^n$, and fix $1 \leq i \leq n - 1$.*

- (a) *If $\lambda_i = \lambda_{i+1} - 1$, then $H^k(\mathbf{F}(V), \mathcal{L}^\lambda(V)) = 0$ for all k .*
- (b) *Suppose that $\lambda_i \geq \lambda_{i+1}$ and that either $\text{char}(\mathbf{k}) = 0$, or $\text{char}(\mathbf{k}) = p > 0$ and furthermore*

$$\lambda_i - \lambda_{i+1} \leq p - 1.$$

We have that for all $k \in \mathbb{Z}$

$$H^k(\mathbf{F}(V), \mathcal{L}^\lambda(V)) = H^{k+1}(\mathbf{F}(V), \mathcal{L}^{s_i \bullet \lambda}(V)).$$

We shall use Proposition 2.2 in order to prove the following vanishing result, which plays a key role in our vanishing theorem for Koszul modules.

Lemma 2.3. *Let V denote a \mathbf{k} -vector space of dimension $n \geq 3$, and assume that $p := \text{char}(\mathbf{k})$ satisfies either $p = 0$ or $p \geq n - 2$. Suppose $2 \leq r \leq 2n - 3$ and consider the weight*

$$\lambda = (n - 2 - r, 1 - r, 0^{n-2}) \in \mathbb{Z}^n.$$

We have that

$$(13) \quad H^{r-1}(\mathbf{F}(V), \mathcal{L}^\lambda(V)) = 0.$$

Proof. Our proof consists of several applications of Proposition 2.2, whose hypothesis is more restrictive when $p > 0$. It is then sufficient to consider the case $p \geq n - 2$ for the rest of the argument. We split our analysis into two cases. To simplify notation, we write \mathbf{F} for $\mathbf{F}(V)$ and \mathcal{L}^μ for $\mathcal{L}^\mu(V)$.

Suppose that $2 \leq r \leq n - 1$ and consider the sequence of partitions $\lambda^1, \dots, \lambda^{r-1}$ defined by

$$\lambda^1 = (n - 2 - r, (-1)^{r-1}, 0^{n-r}) \text{ and } \lambda^{j+1} := s_{r-j} \bullet \lambda^j, \text{ for } j = 1, \dots, r - 2.$$

Note that $\lambda^j = (n - 2 - r, (-1)^{r-1-j}, -j, 0^{n-r+j-1})$ for $1 \leq j \leq r - 1$, and that in particular $\lambda^{r-1} = \lambda$. Since $\lambda_r^1 = \lambda_{r+1}^1 - 1$, it follows from Proposition 2.2(a) with $i = r$ that

$$(14) \quad H^1(\mathbf{F}, \mathcal{L}^{\lambda^1}) = 0.$$

We have for $1 \leq j \leq r - 2$ that the following inequalities

$$0 \leq \lambda_{r-j}^j - \lambda_{r-j+1}^j = j - 1 \leq r - 3 \leq n - 4 < p - 1$$

hold, so it follows from Proposition 2.2(b) with $i = r - j$ that

$$(15) \quad H^j(\mathbf{F}, \mathcal{L}^{\lambda^j}) = H^{j+1}(\mathbf{F}, \mathcal{L}^{\lambda^{j+1}}) \text{ for } j = 1, \dots, r - 2.$$

Combining (14) with (15) and the fact that $\lambda^{r-1} = \lambda$, we obtain (13).

Suppose now that $n \leq r \leq 2n - 3$, and consider the partitions $\lambda^1, \dots, \lambda^{r-n+1}$ defined by

$$\lambda^1 = (-r, (-1)^{r-n+1}, 0^{2n-2-r}) \text{ and } \lambda^{j+1} := s_{r-n+2-j} \bullet \lambda^j \text{ for } j = 1, \dots, r - n.$$

Note that $\lambda^j = (-r, (-1)^{r-n+1-j}, -j, 0^{2n-3-r+j})$ for $1 \leq j \leq r - n + 1$. Since $\lambda_{r-n+2}^1 = \lambda_{r-n+3}^1 - 1$, it follows from Proposition 2.2(a) with $i = r - n + 2$ that

$$(16) \quad H^n(\mathbf{F}, \mathcal{L}^{\lambda^1}) = 0.$$

For $1 \leq j \leq r - n$ we obtain that the following inequalities

$$0 \leq \lambda_{r-n+2-j}^j - \lambda_{r-n+3-j}^j = j - 1 \leq r - n - 1 \leq n - 4 < p - 1$$

hold, so it follows from Proposition 2.2(b) with $i = r - n + 2 - j$ that

$$(17) \quad H^{n+j-1}(\mathbf{F}, \mathcal{L}^{\lambda^j}) = H^{n+j}(\mathbf{F}, \mathcal{L}^{\lambda^{j+1}}), \text{ for all } j = 1, \dots, r - n.$$

Combining (16) with (17) we conclude that

$$H^r(\mathbf{F}, \mathcal{L}^{\lambda^{r-n+1}}) = 0,$$

and note that $\lambda^{r-n+1} = (-r, n - 1 - r, 0^{n-2})$. We now observe that $\lambda^{r-n+1} = s_1 \bullet \lambda$ and that

$$0 \leq \lambda_1 - \lambda_2 = n - 3 \leq p - 1$$

so we can apply Proposition 2.2(b) to obtain the vanishing statements

$$H^{r-1}(\mathbf{F}, \mathcal{L}^\lambda) = H^r(\mathbf{F}, \mathcal{L}^{\lambda^{r-n+1}}) = 0,$$

concluding our proof. \square

2.3. Bott's Theorem for Grassmannians. Let $\mathbf{G}(V) = \text{Gr}_2(V^\vee)$ be the Grassmannian of 2-dimensional quotients of V , or equivalently 2-dimensional subspaces of V^\vee . We write $\mathbf{G} := \mathbf{G}(V)$ and consider the tautological exact sequence

$$(18) \quad 0 \rightarrow \mathcal{R} \rightarrow V \otimes \mathcal{O}_{\mathbf{G}} \rightarrow \mathcal{Q} \rightarrow 0,$$

where \mathcal{R} (respectively \mathcal{Q}) denotes the universal rank $n - 2$ subbundle (respectively rank 2 quotient bundle) of the trivial bundle $V \otimes \mathcal{O}_{\mathbf{G}}$. We write $\mathcal{O}_{\mathbf{G}}(1) := \bigwedge^2 \mathcal{Q}$ for the Plücker line bundle and note that for every $\alpha = (\alpha_1, \alpha_2) \in \mathbb{Z}_{\text{dom}}^2$ we have the identification

$$(19) \quad \mathbb{S}_\alpha \mathcal{Q} = \text{Sym}^{\alpha_1 - \alpha_2} \mathcal{Q} \otimes \mathcal{O}_{\mathbf{G}}(\alpha_2).$$

We wish to compute the cohomology groups of some of the sheaves $\mathbb{S}_\alpha \mathcal{Q}$. To that end we reduce the calculation to the case of line bundles on the complete flag variety $\mathbf{F}(V)$. We write $\mathbf{F} := \mathbf{F}(V)$ and consider the natural map

$$(20) \quad \psi : \mathbf{F} \rightarrow \mathbf{G}, \text{ given by } V_\bullet \mapsto V_2,$$

which enables us to realize \mathbf{F} as the fiber product $\mathbf{F}_{\mathbf{G}}(\mathcal{Q}) \times_{\mathbf{G}} \mathbf{F}_{\mathbf{G}}(\mathcal{R})$. Alternatively, let $Y := \mathbf{F}_{\mathbf{G}}(\mathcal{Q})$ be the *partial flag variety* parametrizing partial flags

$$V \twoheadrightarrow V_2 \twoheadrightarrow V_1 \twoheadrightarrow V_0 = 0$$

and write $\psi_1 : Y \rightarrow \mathbf{G}$ for the structure map when we think of Y as a relative flag variety over \mathbf{G} (which in this case is a \mathbb{P}^1 -bundle). We then identify \mathbf{F} with the relative flag variety $\mathbf{F}_Y(\psi_1^*\mathcal{R})$, and write $\psi_2 : \mathbf{F} \rightarrow Y$ for the structure map. In particular, we have a factorization

$$\psi = \psi_1 \circ \psi_2 : \mathbf{F} \rightarrow \mathbf{G},$$

where ψ_2 forgets the subspaces V_{n-1}, \dots, V_3 and ψ_1 forgets V_1 respectively. Recall that $\mathcal{L}^\lambda = \mathcal{L}^\lambda(V)$.

Theorem 2.4. *Let $\alpha \in \mathbb{Z}_{\text{dom}}^2$ and $\beta \in \mathbb{Z}_{\text{dom}}^{n-2}$, and let $\lambda = (\alpha|\beta) \in \mathbb{Z}^n$ denote their concatenation. Using notation (20) we have*

$$(21) \quad R^k \psi_* (\mathcal{L}^\lambda) = \begin{cases} \mathbb{S}_\alpha \mathcal{Q} \otimes \mathbb{S}_\beta \mathcal{R} & \text{if } k = 0; \\ 0 & \text{if } k > 0. \end{cases}$$

In particular, we have that

$$(22) \quad H^k(\mathbf{G}, \mathbb{S}_\alpha \mathcal{Q} \otimes \mathbb{S}_\beta \mathcal{R}) = H^k(\mathbf{F}, \mathcal{L}^\lambda) \quad \text{for all } k,$$

and furthermore:

- (1) If $q \geq 0$, then $H^0(\mathbf{G}, \text{Sym}^q \mathcal{Q}) = \text{Sym}^q V$ and $H^k(\mathbf{G}, \text{Sym}^q \mathcal{Q}) = 0$ for all $k > 0$.
- (2) For all $a, k \geq 0$, we have that $H^k(\mathbf{G}, \text{Sym}^a \mathcal{Q} \otimes \mathcal{R}) = 0$.
- (3) If $\text{char}(\mathbf{k}) = p$ satisfies either $p = 0$ or $p \geq n - 2$, then

$$(23) \quad H^{r-1}(\mathbf{G}, \mathbb{S}_{(n-2-r, 1-r)} \mathcal{Q}) = 0, \quad \text{for all } r \geq 2.$$

Proof. To prove (21) we first note that via the natural identifications above we have

$$\mathcal{L}^\lambda = \mathcal{L}^\lambda(V) \cong \psi_2^*(\mathcal{L}^\alpha(\mathcal{Q})) \otimes_{\mathcal{O}_{\mathbf{F}}} \mathcal{L}^\beta(\psi_1^*\mathcal{R}).$$

Using the projection formula for the higher direct images along ψ_2 , we obtain

$$R^k \psi_{2*}(\mathcal{L}^\lambda) = \mathcal{L}^\alpha(\mathcal{Q}) \otimes_{\mathcal{O}_Y} R^k \psi_{2*}(\mathcal{L}^\beta(\psi_1^*\mathcal{R})) \stackrel{(12)}{=} \begin{cases} \mathcal{L}^\alpha(\mathcal{Q}) \otimes_{\mathcal{O}_Y} \mathbb{S}_\beta(\psi_1^*\mathcal{R}) & \text{if } k = 0; \\ 0 & \text{if } k > 0. \end{cases}$$

From the compatibility of ψ_1^* with tensor constructions, it follows that $\mathbb{S}_\beta(\psi_1^*\mathcal{R}) = \psi_1^*(\mathbb{S}_\beta \mathcal{R})$. The projection formula along ψ_1 then yields

$$R^k \psi_{1*}(\mathcal{L}^\alpha(\mathcal{Q}) \otimes_{\mathcal{O}_Y} \psi_1^*(\mathbb{S}_\beta \mathcal{R})) = R^k \psi_{1*}(\mathcal{L}^\alpha(\mathcal{Q})) \otimes \mathbb{S}_\beta \mathcal{R} \stackrel{(12)}{=} \begin{cases} \mathbb{S}_\alpha \mathcal{Q} \otimes \mathbb{S}_\beta \mathcal{R} & \text{if } k = 0; \\ 0 & \text{if } k > 0. \end{cases}$$

Using the fact that $R\psi_* = R\psi_{1*} \circ R\psi_{2*}$, we obtain (21). The conclusion (22) follows from (21) and the Leray spectral sequence.

To establish conclusion (1) we apply (22) with $\alpha = (q, 0)$ and $\beta = (0^{n-2})$, together with Theorem 2.1. To prove conclusion (2) we apply (22) with $\alpha = (a, 0)$ and $\beta = (1, 0^{n-3})$, together with Proposition 2.2(a) with $i = 2$ (noting that $\lambda_2 = \lambda_3 - 1$). To prove conclusion (3) we apply (22) with $\alpha = (n - 2 - r, 1 - r)$ and $\beta = (0^{n-2})$, together with Lemma 2.3 when $r \leq 2n - 3$. Finally, when $r > 2n - 3$, then we have $r - 1 > 2n - 4 = \dim(\mathbf{G})$, therefore the vanishing (23) is a consequence of Grothendieck's vanishing theorem, see e.g. [16, Theorem III.2.7]. \square

2.4. Vanishing for Koszul modules. We start by describing a geometric construction of the kernel $\text{Ker}(\delta_1)$ of the Koszul differential $\delta_1 : V \otimes S \rightarrow S$. We let $\mathbf{G} = \text{Gr}_2(V^\vee) \subseteq \mathbb{P}(\bigwedge^2 V^\vee)$ as in Section 2.3, and consider

$$S := \text{Sym}_{\mathcal{O}_{\mathbf{G}}}(\mathcal{Q}) = \mathcal{O}_{\mathbf{G}} \oplus \mathcal{Q} \oplus \text{Sym}^2 \mathcal{Q} \oplus \dots$$

viewed as a sheaf of graded $\mathcal{O}_{\mathbf{G}}$ -algebras on \mathbf{G} . We define Koszul differentials

$$\delta_1^{\mathcal{Q}} : \mathcal{Q} \otimes_{\mathcal{O}_{\mathbf{G}}} S \rightarrow S \quad \text{and} \quad \delta_2^{\mathcal{Q}} : \bigwedge^2 \mathcal{Q} \otimes_{\mathcal{O}_{\mathbf{G}}} S \rightarrow \mathcal{Q} \otimes_{\mathcal{O}_{\mathbf{G}}} S$$

in the usual way. We write $\bigwedge^2 \mathcal{Q} \otimes_{\mathcal{O}_{\mathbf{G}}} S = S(1)$, since $\bigwedge^2 \mathcal{Q} = \mathcal{O}_{\mathbf{G}}(1)$ is the Plücker line bundle.

Lemma 2.5. *We have a commutative diagram*

$$(24) \quad \begin{array}{ccccc} \bigwedge^2 V \otimes S & \xrightarrow{\delta_2} & V \otimes S & \xrightarrow{\delta_1} & S \\ f_2 \downarrow & & f_1 \downarrow & & f_0 \downarrow \\ 0 \longrightarrow & H^0(\mathbf{G}, S(1)) & \xrightarrow{H^0(\mathbf{G}, \delta_2^{\mathcal{Q}})} & H^0(\mathbf{G}, \mathcal{Q} \otimes S) & \xrightarrow{H^0(\mathbf{G}, \delta_1^{\mathcal{Q}})} & H^0(\mathbf{G}, S) \end{array}$$

where the maps f_1 and f_0 are isomorphisms. Moreover, we have an isomorphism $\text{Ker}(\delta_1) \cong H^0(\mathbf{G}, S(1))$.

Proof. The proof is along the lines of [2, Lemma 3.4], which is set in characteristic 0 context. The only potential differences in positive characteristic come from the two calculations of sheaf cohomology on Grassmannians in the said proof, namely

$$H^0(\mathbf{G}, \text{Sym}^q \mathcal{Q}) = \text{Sym}^q V,$$

which is covered by conclusion (1) of Theorem 2.4, and

$$H^0(\mathbf{G}, \mathcal{R} \otimes \text{Sym}^q \mathcal{Q}) = H^1(\mathbf{G}, \mathcal{R} \otimes \text{Sym}^q \mathcal{Q}) = 0 \text{ for all } q \geq 0,$$

which follows from conclusion (2) of Theorem 2.4. The rest of the argument is manifestly characteristic independent, so we do not reproduce it here. \square

Using Lemma 2.5, we obtain the following geometric description of the Koszul module

$$(25) \quad W(V, K) = \text{Coker} \left\{ H^0(\mathbf{G}, K \otimes S) \xrightarrow{H^0(\mathbf{G}, \eta)} H^0(\mathbf{G}, S(1)) \right\},$$

where $\eta : K \otimes S \rightarrow S(1)$ is induced by

$$K \otimes \mathcal{O}_{\mathbf{G}} \hookrightarrow \bigwedge^2 V \otimes \mathcal{O}_{\mathbf{G}} \twoheadrightarrow \bigwedge^2 \mathcal{Q} = \mathcal{O}_{\mathbf{G}}(1).$$

With all these preparations in place, the proof of Theorem 1.3 follows by extending the argument for [2, Theorem 3.1] to positive characteristic, as follows.

Proof of Theorem 1.3. If $W_q(V, K) = 0$ for $q \geq n - 3$ then $\mathcal{R}(V, K) = \{0\}$ by (6). For the converse, it suffices to prove that $W_{n-3}(V, K)$: since the module $W(V, K)$ is generated in degree zero, this implies that $W_q(V, K) = 0$ for all $q \geq n - 3$. The assumption that $\mathcal{R}(V, K) = \{0\}$ yields using (6) that $(K, \mathcal{O}_{\mathbf{G}}(1))$ is base point free, and hence $K \otimes \mathcal{O}_{\mathbf{G}} \rightarrow \mathcal{O}_{\mathbf{G}}(1)$ is surjective. This gives rise to an exact Koszul complex

$$\mathcal{K}^\bullet : 0 \longrightarrow \bigwedge^m K \otimes \mathcal{O}_{\mathbf{G}}(1 - m) \longrightarrow \dots \longrightarrow \bigwedge^2 K \otimes \mathcal{O}_{\mathbf{G}}(-1) \longrightarrow K \otimes \mathcal{O}_{\mathbf{G}} \longrightarrow \mathcal{O}_{\mathbf{G}}(1) \longrightarrow 0$$

where $\mathcal{K}^{-i} = \bigwedge^i K \otimes \mathcal{O}_{\mathbf{G}}(1-i)$. Tensoring with $\mathrm{Sym}^{n-3} \mathcal{Q}$ we get a hypercohomology spectral sequence $E_1^{-i,j} = H^j(\mathbf{G}, \mathcal{K}^{-i} \otimes \mathrm{Sym}^{n-3} \mathcal{Q}) \implies 0$, where

$$(26) \quad E_1^{-i,j} = \bigwedge^i K \otimes H^j(\mathbf{G}, \mathbb{S}_{(n-2-i, 1-i)} \mathcal{Q}).$$

We have using (25) that

$$W_{n-3}(V, K) = \mathrm{Coker}(E_1^{-1,0} \longrightarrow E^{0,0})$$

and we suppose by contradiction that this map is not surjective. Since $E_{\infty}^{0,0} = 0$, there must be some non-zero differential

$$E_r^{-r, r-1} \longrightarrow E_r^{0,0} \text{ for } r \geq 2, \text{ or } E_r^{0,0} \longrightarrow E_r^{r, 1-r} \text{ for } r \geq 1.$$

Since $E_1^{r, 1-r} = 0$ for all $r \geq 1$ it follows that $E_r^{r, 1-r} = 0$ as well, so the latter case does not occur. To prove that the former case does not occur either and obtain a contradiction, it suffices to check that $E_1^{-r, r-1} = 0$ for all $r \geq 2$, which follows from (26) and from conclusion (3) of Theorem 2.4. \square

3. SYMMETRIC, DIVIDED AND EXTERIOR POWERS, AND HERMITE RECIPROCITY

The goal of this section is to study various tensor constructions in arbitrary characteristic, that will be used in our analysis of the syzygies of the tangent developable to a rational normal curve in Section 5. For a more thorough discussion, we refer the reader to [1]. In Section 3.1 we introduce the symmetric, divided and exterior powers of a vector space V , which we denote by $\mathrm{Sym}^{\bullet} V$, $D^{\bullet} V$, and $\bigwedge^{\bullet} V$ respectively. In Section 3.2 we explain how the natural ambient space of the Veronese embedding is defined in terms of divided powers. We then focus on tensor constructions for a vector space U of dimension 2, discuss a number of $\mathfrak{sl}(U)$ -equivariant maps in Section 3.3, and prove a version of Hermite reciprocity in arbitrary characteristic in Section 3.4: more precisely, we produce an explicit $\mathfrak{sl}(U)$ -equivariant isomorphism

$$\mathrm{Sym}^d(D^i U) \longrightarrow \bigwedge^i(\mathrm{Sym}^{d+i-1} U)$$

for all $d, i \geq 1$. We work over an arbitrary field \mathbf{k} .

3.1. Symmetric, divided and exterior powers. Let V denote a finite dimensional vector space over a field \mathbf{k} , and for $d > 0$ consider the tensor power $T^d V := V^{\otimes d} = V \otimes V \otimes \cdots \otimes V$ with the natural action of the symmetric group \mathfrak{S}_d by permuting the factors. The *divided power* $D^d V$ is defined as the set of symmetric tensors in $T^d V$, that is,

$$D^d V := \{\omega \in T^d V : \sigma(\omega) = \omega \text{ for all } \sigma \in \mathfrak{S}_d\}.$$

If we consider the subspace of $T^d V$ defined by

$$\Sigma_d := \mathrm{Span}\{\sigma(\omega) - \omega : \sigma \in \mathfrak{S}_d \text{ and } \omega \in T^d V\},$$

then the *symmetric power* $\mathrm{Sym}^d V$ is defined as the quotient $\mathrm{Sym}^d V := T^d V / \Sigma_d$. If V has a basis (x_1, \dots, x_n) , then $\mathrm{Sym}^d V$ identifies with the space of homogeneous polynomials of degree d in x_1, \dots, x_n , and as such it has a basis of monomials $x_1^{a_1} \cdots x_n^{a_n}$, where $a_1 + \cdots + a_n = d$. To get a basis for $D^d V$ we first consider the orbits

$$O_{a_1, \dots, a_n} := \mathfrak{S}_d \cdot x_1^{\otimes a_1} \otimes x_2^{\otimes a_2} \otimes \cdots \otimes x_n^{\otimes a_n}$$

and for $a_1 + \cdots + a_n = d$ consider the *divided power monomials*

$$x_1^{(a_1)} \cdots x_n^{(a_n)} := \sum_{\omega \in O_{a_1, \dots, a_n}} \omega.$$

They form a basis for $D^d(V)$, and in particular we have that $\dim(\text{Sym}^d V) = \dim(D^d V)$. By composing the inclusion of $D^d V$ into $T^d V$ with the projection onto $\text{Sym}^d V$ we obtain a natural map

$$(27) \quad D^d V \longrightarrow \text{Sym}^d V, \quad x_1^{(a_1)} \cdots x_n^{(a_n)} \mapsto \binom{d}{a_1, \dots, a_n} x_1^{a_1} \cdots x_n^{a_n}, \quad \text{where } \binom{d}{a_1, \dots, a_n} = \frac{d!}{a_1! \cdots a_n!}.$$

This map is an isomorphism in characteristic zero, or more generally when the multinomial coefficients are invertible. In general it is neither injective, nor surjective.

We next consider the subspace of $T^d V$ defined by

$$\Xi_d := \text{Span}\{v_1 \otimes \cdots \otimes v_n : v_i \in V \text{ and } v_i = v_j \text{ for some } i \neq j\}$$

and define the *exterior power* $\Lambda^d V$ as the quotient $\Lambda^d V := T^d V / \Xi_d$. If we write $v_1 \wedge \cdots \wedge v_d$ for the class of $v_1 \otimes \cdots \otimes v_d$ in the quotient, then $\Lambda^d V$ has a basis consisting of $x_{i_1} \wedge \cdots \wedge x_{i_d}$, where $1 \leq i_1 < \cdots < i_d \leq n$. There is a natural inclusion of $\Lambda^d V$ into $T^d V$ given by

$$(28) \quad v_1 \wedge \cdots \wedge v_d \longrightarrow \sum_{\sigma \in \mathfrak{S}_d} \text{sgn}(\sigma) \cdot \sigma(v_1 \otimes \cdots \otimes v_n).$$

This gives a splitting of the quotient map $T^d V \twoheadrightarrow \Lambda^d V$ if and only if $d!$ is invertible in \mathbf{k} .

There is a natural \mathfrak{S}_n -invariant perfect pairing

$$(29) \quad T^d(V) \times T^d(V^\vee) \longrightarrow \mathbf{k},$$

defined on pure tensors via

$$\langle v_1 \otimes \cdots \otimes v_d, f_1 \otimes \cdots \otimes f_d \rangle = f_1(v_1) f_2(v_2) \cdots f_d(v_d), \quad \text{where } v_i \in V \text{ and } f_j \in V^\vee.$$

This induces a perfect pairing between $\text{Sym}^d V$ and $D^d(V^\vee)$, giving a natural identification

$$(\text{Sym}^d V)^\vee \cong D^d(V^\vee).$$

If (e_1, \dots, e_n) is the basis of V^\vee dual to the basis (x_1, \dots, x_n) of V , then the divided power monomials $e_1^{(a_1)} \cdots e_n^{(a_n)}$ give a basis of $D^d(V^\vee)$ dual to the monomial basis of $\text{Sym}^d V$.

If we think of $\Lambda^d V$ as a quotient of $T^d V$, and of $\Lambda^d V^\vee$ as a subspace of $T^d(V^\vee)$ via (28), the pairing (29) induces a perfect pairing $\Lambda^d V \times \Lambda^d(V^\vee) \longrightarrow \mathbf{k}$, described on decomposable elements via

$$\langle v_1 \wedge \cdots \wedge v_d, f_1 \wedge \cdots \wedge f_d \rangle \longrightarrow \det[f_i(v_j)]_{1 \leq i, j \leq d}, \quad \text{where } v_i \in V \text{ and } f_j \in V^\vee.$$

We get a natural isomorphism $(\Lambda^d V)^\vee \cong \Lambda^d(V^\vee)$. We write $\Lambda^d V^\vee$ for either of the two spaces above. By contrast, we shall be careful in distinguishing between $\text{Sym}^d(V^\vee)$ and $(\text{Sym}^d V)^\vee \cong D^d(V^\vee)$!

If $V = U$ is of dimension two, then we get a natural isomorphism $U \cong \Lambda^2 U \otimes U^\vee$, which extends to tensor powers to an isomorphism $T^d(U) \cong (\Lambda^2 U)^{\otimes d} \otimes T^d(U^\vee)$. Restricting to \mathfrak{S}_n -invariant tensors, we obtain an isomorphism $D^d(U) \cong (\Lambda^2 U)^{\otimes d} \otimes D^d(U^\vee)$, which will be used in Sections 5.5 and 5.6.

3.2. The Veronese embedding. We fix a vector space U of dimension two and denote by $\mathbf{P}^1 := \mathbb{P}(U^\vee)$ the projective space of 1-dimensional subspaces of U^\vee . Then $H^0(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(d)) = \text{Sym}^d U$, which is the vector space of linear forms on $(\text{Sym}^d U)^\vee \cong D^d(U^\vee)$. Setting $\mathbf{P}^d := \mathbb{P}(D^d(U^\vee))$, the linear series $|\mathcal{O}_{\mathbf{P}^1}(d)|$ gives an embedding

$$\nu_d : \mathbf{P}^1 \longrightarrow \mathbf{P}^d, \quad [f] \mapsto [f \otimes \cdots \otimes f].$$

If U has a basis (x_1, x_2) , we consider the induced monomial bases for $\text{Sym}^d U$ and $D^d U$ as in Section 3.1. For simplicity, it will be convenient to use a *dehomogenized* notation, where

$$(30) \quad x_1 = x, \quad x_2 = 1, \quad x_1^i x_2^{d-i} = x^i \text{ and } x_1^{(i)} x_2^{(d-i)} = x^{(i)}.$$

If we write $(1, y)$ for the basis of U^\vee dual to $(1, x)$ and use the analogous conventions to (30), then the Veronese map $\nu_d : \mathbf{P}^1 \rightarrow \mathbf{P}^d$ is expressed as

$$[a + by] \rightarrow [(a + by) \otimes \cdots \otimes (a + by)] = [a^d \cdot y^{(0)} + a^{d-1}b \cdot y^{(1)} + a^{d-2}b^2 \cdot y^{(2)} + \cdots + b^d \cdot y^{(d)}],$$

that is, relative to the chosen bases, ν_d takes the familiar form $[a : b] \rightarrow [a^d : a^{d-1}b : a^{d-2}b^2 : \cdots : b^d]$.

3.3. Some \mathfrak{sl}_2 -equivariant maps. We let U as in the previous section, with \mathbf{k} -basis $(1, x)$ and corresponding bases $(1, x, \dots, x^d)$ and $(x^{(0)}, x^{(1)}, x^{(2)}, \dots, x^{(d)})$ for $\text{Sym}^d U$ and $D^d U$ respectively. We let $\mathfrak{sl}_2 := \mathfrak{sl}(U)$ denote the Lie algebra of \mathbf{k} -endomorphisms of U with trace 0. We write \mathfrak{L} (respectively \mathfrak{R}) for the *lowering* (respectively *raising*) operators in \mathfrak{sl}_2 , whose action on $\text{Sym}^d U$ and $D^d U$ is given by

$$\mathfrak{L} \cdot x^i = i \cdot x^{i-1}, \quad \mathfrak{L} \cdot x^{(i)} = (d - i + 1) \cdot x^{(i-1)}, \quad \mathfrak{R} \cdot x^i = (d - i) \cdot x^{i+1}, \quad \mathfrak{R} \cdot x^{(i)} = (i + 1) \cdot x^{(i+1)}.$$

Relative to the conventions (30), one can think of \mathfrak{L} as the operator $x_2 \cdot \frac{\partial}{\partial x_1}$, and of \mathfrak{R} as $x_1 \cdot \frac{\partial}{\partial x_2}$. The natural map (27) is described on the basis elements by

$$(31) \quad D^d U \rightarrow \text{Sym}^d U, \quad x^{(i)} \mapsto \binom{d}{i} \cdot x^i,$$

and can be checked to be \mathfrak{sl}_2 -equivariant. By abuse of notation, whenever we refer to $x^{(i)}$ as an element of $\text{Sym}^d U$ we will interpret it as $\binom{d}{i} \cdot x^i$.

For $a, b \geq 0$, we define two \mathfrak{sl}_2 -equivariant maps, namely the *multiplication* map

$$(32) \quad \mu = \mu_U : \text{Sym}^a U \otimes \text{Sym}^b U \rightarrow \text{Sym}^{a+b} U, \quad x^i \otimes x^j \rightarrow x^{i+j}$$

and the *co-multiplication* map

$$(33) \quad \Delta = \Delta_U : D^{a+b} U \rightarrow D^a U \otimes D^b U, \quad x^{(t)} \rightarrow \sum_{\substack{i+j=t \\ i \leq a, j \leq b}} x^{(i)} \otimes x^{(j)}.$$

We may abuse language and refer to Δ as the *dual* of μ : the correct interpretation of this statement is that via the isomorphism $D^d(U^\vee) \cong (\text{Sym}^d U)^\vee$ one has that μ_U is dual to Δ_{U^\vee} , and μ_{U^\vee} is dual to Δ_U . We shall need in Proposition 5.6 the map $\Delta^{(2)} : D^{a+2} U \rightarrow D^a U \otimes \text{Sym}^2 U$ obtained as the composition:

$$\Delta^{(2)} : D^{a+2} U \xrightarrow{(33)} D^a U \otimes D^2 U \xrightarrow{(31)} D^a U \otimes \text{Sym}^2 U.$$

It is important to observe that the following diagram is commutative:

$$(34) \quad \begin{array}{ccc} D^{a+2} U & \xrightarrow{\Delta^{(2)}} & D^a U \otimes \text{Sym}^2 U \\ \Delta \downarrow & & \uparrow \text{id}_{D^a U} \otimes \mu \\ D^{a+1} U \otimes U & \xrightarrow{\Delta \otimes \text{id}_U} & D^a U \otimes U \otimes U \end{array}$$

Lemma 3.1. *Let $a \geq 1$. There exist \mathfrak{sl}_2 -equivariant maps*

$$\mu_1 : \text{Sym}^a U \otimes \text{Sym}^a U \rightarrow \text{Sym}^{2a-2} U \text{ and } \Delta_1 : D^{2a-2} U \rightarrow D^a U \otimes D^a U$$

defined by $\mu_1(x^i \otimes x^j) = (i - j) \cdot x^{i+j-1}$ and respectively

$$(35) \quad \Delta_1(x^{(t)}) = \sum_{i+j=t+1} (i-j) \cdot x^{(i)} \otimes x^{(j)}$$

Moreover, if $\text{char}(\mathbf{k}) \neq 2$ then μ_1 is surjective and Δ_1 is injective.

Proof. Note that Δ_1 is the dual of μ_1 (in the same way that Δ is dual to μ), so it suffices to verify that μ_1 is well-defined, \mathfrak{sl}_2 -equivariant and surjective when $\text{char}(\mathbf{k}) \neq 2$. Note first that if $(i, j) \neq (0, 0), (a, a)$, then $0 \leq i+j-1 \leq 2a-2$, so $\mu_1(x^i \otimes x^j) \in \text{Sym}^{2a-2} U$. If $(i, j) = (0, 0)$ or $(i, j) = (a, a)$, then $\mu_1(x^i \otimes x^j) = 0$ is still well-defined. To check equivariance, it suffices to verify that μ_1 commutes with the operators \mathfrak{L} and \mathfrak{R} . The condition $\mu_1(\mathfrak{L}(x^i \otimes x^j)) = \mathfrak{L}(\mu_1(x^i \otimes x^j))$ is equivalent to

$$i \cdot [(i-1) - j] + j \cdot [i - (j-1)] = (i+j-1) \cdot (i-j),$$

which follows by inspection. Similarly, $\mu_1(\mathfrak{R}(x^i \otimes x^j)) = \mathfrak{R}(\mu_1(x^i \otimes x^j))$ follows from a direct calculation. Consider any $0 \leq t \leq 2a-2$. If $t = 2r$ then $\mu_1(x^{r+1} \otimes x^r) = x^t$, so $x^t \in \text{Im}(\mu_1)$. If $t = 2r+1$ then $\mu_1(x^{r+2} \otimes x^r) = 2x^t$, so $x^t \in \text{Im}(\mu_1)$ when 2 is invertible in \mathbf{k} . This proves the surjectivity of μ_1 . \square

From the definition of μ_1 , observe that $\mu_1(x^i \otimes x^i) = 0$ for all i , and $\mu_1(x^i \otimes x^j + x^j \otimes x^i) = 0$ for $i \neq j$. It follows that $\mu_1(v \otimes v) = 0$ for all $v \in \text{Sym}^a U$, so μ_1 factors through $\bigwedge^2 \text{Sym}^a U$. We get a map

$$(36) \quad \mu_1 : \bigwedge^2 \text{Sym}^a U \longrightarrow \text{Sym}^{2a-2} U,$$

which, as explained in the Introduction, can be thought of as the Gaussian–Wahl map of $\mathcal{O}_{\mathbb{P}^1}(a)$ (see (10)). Dually, the map Δ_1 in Lemma 3.1 can be viewed as a map

$$(37) \quad \Delta_1 : D^{2a-2} U \longrightarrow \bigwedge^2 D^a U.$$

3.4. A Hermite type identification in arbitrary characteristic. Hermite reciprocity [19] over a field \mathbf{k} of characteristic 0 asserts the existence of an \mathfrak{sl}_2 -isomorphism $\text{Sym}^d(\text{Sym}^i U) \cong \text{Sym}^i(\text{Sym}^d U)$, see [12, Exercise 11.34]. Under the same assumption, one has the isomorphism $\text{Sym}^d(\text{Sym}^i U) \cong \bigwedge^d(\text{Sym}^{d+i-1} U)$, or equivalently $\text{Sym}^d(\text{Sym}^i U) \cong \bigwedge^i(\text{Sym}^{d+i-1} U)$, see [12, Exercise 11.35]. These isomorphisms are usually proved non-constructively by identifying the characters of the two representations, and they no longer hold in positive characteristic. The goal of this section is to describe an explicit \mathfrak{sl}_2 -isomorphism

$$(38) \quad \psi_d^i : \text{Sym}^d(D^i U) \longrightarrow \bigwedge^i(\text{Sym}^{d+i-1} U)$$

in arbitrary characteristic.

Remark 3.2. Since $\bigwedge^2 U$ is isomorphic to the trivial \mathfrak{sl}_2 -representation, we get that $U \cong U^\vee$. More generally, if V is an \mathfrak{sl}_2 -representation with $N = \dim(V)$ then we have isomorphisms $\bigwedge^r V \cong \bigwedge^{N-r} V^\vee$. Combining this with $(\text{Sym}^d V)^\vee \cong D^d(V^\vee)$, we get a chain of \mathfrak{sl}_2 -isomorphisms

$$\text{Sym}^d(D^i U) \cong \bigwedge^i(\text{Sym}^{d+i-1} U) \cong \bigwedge^d(\text{Sym}^{d+i-1} U)^\vee \cong \bigwedge^d(D^{d+i-1}(U^\vee)) \cong \bigwedge^d(D^{d+i-1} U),$$

which is an arbitrary characteristic analogue of [12, Exercise 11.35]. Moreover,

$$\text{Sym}^d(D^i U) \cong \bigwedge^d(\text{Sym}^{d+i-1} U)^\vee \cong (\text{Sym}^i(D^d U))^\vee \cong D^i(\text{Sym}^d(U^\vee)) \cong D^i(\text{Sym}^d U)$$

is an arbitrary characteristic version of Hermite reciprocity [12, Exercise 11.34].

To construct (38) we identify both sides with an appropriate subspace of the ring of symmetric polynomials, and construct ψ_d^i from the change of basis between elementary symmetric and Schur polynomials.

For $i > 0$ we let \mathcal{P}_i be the collection of partitions with at most i parts: an element $\lambda \in \mathcal{P}_i$ is written as $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_i)$, with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_i \geq 0$. We let \mathcal{P}'_i denote the collection of partitions with parts of size at most i , write λ' for the conjugate of a partition λ , and note that $\lambda \in \mathcal{P}_i$ if and only if $\lambda' \in \mathcal{P}'_i$. We let $\Lambda_i \subseteq \mathbf{k}[z_1, \dots, z_i]$ denote the ring of symmetric polynomials in z_1, \dots, z_i , and consider the following two well-studied bases of Λ_i (see [22, Chapter I]), indexed by \mathcal{P}_i and \mathcal{P}'_i :

- For every $\lambda \in \mathcal{P}_i$, the *Schur polynomial* s_λ is defined via

$$s_\lambda(z_1, \dots, z_i) = \frac{\det(z_k^{\lambda_\ell + i - k})_{1 \leq k, \ell \leq i}}{\det(z_k^{i-k})_{1 \leq k, \ell \leq i}}.$$

The collection $\{s_\lambda : \lambda \in \mathcal{P}_i\}$ is a basis for Λ_i , which we call the *Schur basis*.

- The *elementary symmetric polynomials* $e_0 = 1, e_1, \dots, e_i$ are defined via the equality

$$e_0 + e_1 \cdot T + e_2 \cdot T^2 + \dots + e_i \cdot T^i = (1 + z_1 \cdot T) \cdots (1 + z_i \cdot T),$$

where T is an auxiliary variable. For each $\mu \in \mathcal{P}'_i$ let $e_\mu := e_{\mu_1} \cdot e_{\mu_2} \cdots$. The collection $\{e_\mu : \mu \in \mathcal{P}'_i\}$ is also a basis for Λ_i , which we call the *elementary symmetric basis*.

For each $j = 0, \dots, i$, one has $e_j = s_{(1^j)}$, where $(1^j) = (1, 1, \dots, 1)$. More generally, for $d \geq 1$ we consider the following collections of partitions

$$\mathcal{P}_i(d) := \{\lambda \in \mathcal{P}_i : \lambda_1 \leq d\}, \quad \mathcal{P}'_i(d) := \{\mu \in \mathcal{P}'_i : \mu_{d+1} = 0\}.$$

The correspondence $\lambda \longleftrightarrow \lambda'$ induces a bijection between $\mathcal{P}_i(d)$ and $\mathcal{P}'_i(d)$. Moreover, we have that $\{s_\lambda : \lambda \in \mathcal{P}_i(d)\}$ spans the same subspace of Λ_i as $\{e_\mu : \mu \in \mathcal{P}'_i(d)\}$. We denote this subspace by $\Lambda_i(d)$, and get in this way a filtration of Λ_i satisfying $\Lambda_i(d) \cdot \Lambda_i(d') = \Lambda_i(d + d')$, for all $d, d' \geq 0$.

The multiplication map on Λ_i is easy to describe with respect to the elementary symmetric basis, but it is more subtle (based on the Littlewood–Richardson rule) with respect to the Schur basis. For our purposes it is sufficient to consider products of the form $s_\lambda \cdot s_{(1^j)} = s_\lambda \cdot e_j$, where the multiplication is described by Pieri's rule. To describe it we first make some conventions. We write $[i] := \{1, \dots, i\}$ and let $\binom{[i]}{k}$ be the collection of k -element subsets of $[i]$.

- We extend the definition of s_λ for all $\lambda \in \mathbb{Z}_{\geq 0}^i$ by setting $s_\lambda = 0$ when $\lambda \notin \mathcal{P}_i$, that is, when there exists some index j for which $\lambda_j < \lambda_{j+1}$.
- If $I \subseteq [i]$, we write (1^I) for the element $\lambda \in \mathbb{Z}_{\geq 0}^i$ with $\lambda_j = 1$ for $j \in I$ and $\lambda_j = 0$ for $j \notin I$. When $I = [k]$, for $k \leq i$ we have $(1^I) = (1^k)$.

With these conventions, Pieri's rule can be stated as follows:

$$(39) \quad s_\lambda \cdot e_j = \sum_{I \in \binom{[i]}{j}} s_{\lambda + (1^I)} = \sum_{I \in \binom{[i]}{j}, \lambda + (1^I) \in \mathcal{P}_i} s_{\lambda + (1^I)}.$$

Even though \mathfrak{sl}_2 does not act a priori on $\Lambda_i(d)$, we construct an \mathfrak{sl}_2 -equivariant vector space isomorphism ψ_d as in (38) by identifying both sides with $\Lambda_i(d)$. We define the *elementary symmetric basis* of $\text{Sym}^d(\mathbb{D}^i U)$ by associating to each element $\mu \in \mathcal{P}'_i(d)$ the basis element $e_\mu(x) \in \text{Sym}^d(\mathbb{D}^i U)$ defined by

$$(40) \quad e_\mu(x) = x^{(\mu_1)} \cdot x^{(\mu_2)} \cdots x^{(\mu_d)}.$$

This choice of basis induces a vector space isomorphism

$$\epsilon_d^i : \text{Sym}^d(\mathbb{D}^i U) \longrightarrow \Lambda_i(d), \quad \epsilon_d^i(e_\mu(x)) = e_\mu, \text{ for all } \mu \in \mathcal{P}'_i(d).$$

Thinking of $\text{Sym}(\mathbb{D}^i U)$ as a polynomial ring on $x^{(0)}, x^{(1)}, \dots, x^{(i)}$, we can define a ring homomorphism $\text{Sym}(\mathbb{D}^i U) \rightarrow \Lambda_i$ by sending $x^{(0)} \mapsto 1$, $x^{(j)} \mapsto e_j$, for $j = 1, \dots, i$. The map ϵ_d^i is then obtained by restricting this homomorphism to $\text{Sym}^d(\mathbb{D}^i U)$. It follows that

$$(41) \quad \epsilon_{d+1}^i(m \cdot x^{(j)}) = \epsilon_d^i(m) \cdot \epsilon_1^i(x^{(j)}) \text{ for all } m \in \text{Sym}^d(\mathbb{D}^i U) \text{ and } j = 0, \dots, i.$$

We next define the *Schur basis* of $\bigwedge^i(\text{Sym}^{d+i-1} U)$ by associating to an element $\lambda \in \mathcal{P}_i(d)$ the basis element $s_\lambda(x) \in \bigwedge^i(\text{Sym}^{d+i-1} U)$ defined by

$$(42) \quad s_\lambda(x) := x^{\lambda_1+i-1} \wedge x^{\lambda_2+i-2} \wedge \dots \wedge x^{\lambda_i}.$$

This choice of basis induces a vector space isomorphism

$$\sigma_d^i : \Lambda_i(d) \longrightarrow \bigwedge^i(\text{Sym}^{d+i-1} U), \quad \sigma_d^i(s_\lambda) = s_\lambda(x), \text{ for all } \lambda \in \mathcal{P}_i(d).$$

The maps σ_d^i satisfy a compatibility analogous to (41) as explained next. For all \mathbf{k} -vector spaces T and T' we have a natural inclusion, see [1, Theorem III.2.4] and [31, p.96].

$$\left(\bigwedge^i T \right) \otimes \mathbb{D}^i(T') \longrightarrow \bigwedge^i (T \otimes T').$$

Applying this when $T = \text{Sym}^{d+i-1} U$ and $T' = U$, we obtain an \mathfrak{sl}_2 -equivariant inclusion followed by an \mathfrak{sl}_2 -equivariant map induced by $\mu : \text{Sym}^{d+i-1} U \otimes U \rightarrow \text{Sym}^{d+i} U$:

$$\begin{array}{c} (\bigwedge^i \text{Sym}^{d+i-1} U) \otimes \mathbb{D}^i U \longrightarrow \bigwedge^i(\text{Sym}^{d+i-1} U \otimes U) \longrightarrow \bigwedge^i(\text{Sym}^{d+i} U) \\ \searrow \hspace{15em} \nearrow \\ \nu_d^i \end{array}$$

The map ν_d^i can be expressed concretely as follows: for every $\lambda \in \mathcal{P}_i(d)$ and every $j = 0, \dots, i$, we have

$$(43) \quad \nu_d^i(s_\lambda(x) \otimes x^{(j)}) = \sum_{I \in \binom{[i]}{j}} s_{\lambda+(1^I)}(x),$$

where in order for the right hand side to make sense, we have to extend the definition (42) for arbitrary $\lambda \in \mathbb{Z}_{\geq 0}^i$. Notice however that if $\mu = \lambda + (1^I)$ appears in (43) and $\mu \in \mathbb{Z}_{\geq 0}^i \setminus \mathcal{P}_i(d+1)$ then there exists some index $k \in [i-1]$ with $\mu_{k+1} = \mu_k + 1$, and therefore $\mu_k + i - k = \mu_{k+1} + i - (k+1)$, which yields

$$s_\mu(x) = x^{\mu_1+i-1} \wedge \dots \wedge x^{\mu_k+i-k} \wedge x^{\mu_{k+1}+i-(k+1)} \wedge \dots \wedge x^{\mu_i} = 0.$$

We conclude from (39) and (43) that

$$(44) \quad \sigma_{d+1}^i(f \cdot e_j) = \nu_d^i(\sigma_d(f) \otimes x^{(j)}), \text{ for all } f \in \Lambda_i(d) \text{ and } j = 0, \dots, i.$$

Lemma 3.3. *For each $d \geq 0$, the vector space isomorphism*

$$\psi_d^i : \text{Sym}^d(\mathbb{D}^i U) \longrightarrow \bigwedge^i(\text{Sym}^{d+i-1} U)$$

defined by $\psi_d^i = \sigma_d^i \circ \epsilon_d^i$ is \mathfrak{sl}_2 -equivariant.

Proof. We prove this by induction on d . In the case $d = 0$ we have

$$\psi_0^i : \mathbf{k} \longrightarrow \bigwedge^i (\text{Sym}^{i-1} U), \quad 1 \mapsto x^{i-1} \wedge x^{i-2} \wedge \cdots \wedge 1,$$

which is an \mathfrak{sl}_2 -equivariant isomorphism. Assume now $d > 0$ and consider the commutative diagram

$$(45) \quad \begin{array}{ccccc} \text{Sym}^d(\mathbb{D}^i U) \otimes \mathbb{D}^i U & \xrightarrow{\epsilon_d^i \otimes \epsilon_1^i} & \Lambda_i(d) \otimes \Lambda_i(1) & \xrightarrow{\sigma_d^i \otimes (\epsilon_1^i)^{-1}} & \bigwedge^i (\text{Sym}^{d+i-1} U) \otimes \mathbb{D}^i U \\ \downarrow & & \downarrow & & \downarrow \nu_d^i \\ \text{Sym}^{d+1}(\mathbb{D}^i U) & \xrightarrow{\epsilon_{d+1}^i} & \Lambda_i(d+1) & \xrightarrow{\sigma_{d+1}^i} & \bigwedge^i (\text{Sym}^{d+i} U) \end{array}$$

where the left square commutes by (41) while the right square commutes by (44). The left vertical map and ν_d^i are \mathfrak{sl}_2 -equivariant. The composition on the top line is $(\sigma_d^i \otimes (\epsilon_1^i)^{-1}) \circ (\epsilon_d^i \otimes \epsilon_1^i) = \psi_d^i \otimes \text{id}$, which is \mathfrak{sl}_2 -equivariant since ψ_d^i is \mathfrak{sl}_2 -equivariant by induction. It follows that ψ_{d+1}^i , which is the composition of the maps on the bottom line of the diagram, is \mathfrak{sl}_2 -equivariant, concluding our induction. \square

For later use, we record that the Koszul differential gives rise to an \mathfrak{sl}_2 -equivariant inclusion

$$k_i^d : \bigwedge^i \text{Sym}^d U \longrightarrow \bigwedge^{i-1} \text{Sym}^d U \otimes \text{Sym}^d U, \text{ given by}$$

$$(46) \quad k_i^d(s_\lambda(x)) = \sum_{j=1}^i (-1)^{j-1} s_{\lambda_j}(x) \otimes x^{\lambda_j+i-j}, \text{ for } \lambda \in \mathcal{P}_i(d-i+1), \text{ where}$$

$$(47) \quad \lambda^{\hat{j}} = (\lambda_1 + 1, \dots, \lambda_{j-1} + 1, \lambda_{j+1}, \dots, \lambda_i) \in \mathcal{P}_{i-1}(d-i+2).$$

4. SYZYGIES AND THE KEMPF–WEYMAN GEOMETRIC TECHNIQUE

The goal of this section is to summarize a number of standard results that relate to syzygies, minimal resolutions, and Koszul cohomology groups, and to establish some relevant notation that will be used in the rest of the article. In Section 4.1 we discuss a relationship between Koszul cohomology groups and kernel bundles, which will be important in Section 6. In Section 4.2 we describe the minimization of the mapping cone for certain morphisms of complexes, that will be used in the proof of Corollary 5.10 and in that of Theorem 5.4. Finally, in Section 4.3 we recall the basic aspects of the Kempf–Weyman geometric technique for constructing syzygies, that will be featured several times in Sections 5.5 and 5.6.

4.1. Syzygies and Koszul cohomology. We begin by recalling a few basic facts and notation regarding Koszul cohomology groups, and we refer the reader to [13] and [3] for more details. Let X be a projective variety over a field \mathbf{k} , let L be a very ample line bundle on X and consider the associated embedding $\varphi_L : X \longrightarrow \mathbb{P}(H^0(X, L)^\vee) \cong \mathbb{P}^r$. For a sheaf \mathcal{F} on X , we consider the module

$$\Gamma_X(\mathcal{F}, L) := \bigoplus_{n \geq 0} H^0(X, \mathcal{F} \otimes L^n),$$

which can be regarded as a graded module over the polynomial algebra $S := \text{Sym } H^0(X, L)$. For $i, j \geq 0$, the *Koszul cohomology group of i -th order syzygies of weight j* is defined by

$$K_{i,j}(X, \mathcal{F}, L) := \text{Tor}_i^S(\Gamma_X(\mathcal{F}, L), \mathbf{k})_{i+j},$$

or equivalently as the middle homology of the 3-term complex

$$(48) \quad \bigwedge^{i+1} H^0(X, L) \otimes \Gamma_X(\mathcal{F}, L)_{j-1} \longrightarrow \bigwedge^i H^0(X, L) \otimes \Gamma_X(\mathcal{F}, L)_j \longrightarrow \bigwedge^{i-1} H^0(X, L) \otimes \Gamma_X(\mathcal{F}, L)_{j+1}.$$

When $\mathcal{F} = \mathcal{O}_X$, we write as usual $\Gamma_X(L) := \Gamma_X(\mathcal{O}_X, L)$ and $K_{i,j}(X, L) := K_{i,j}(X, \mathcal{O}_X, L)$. The Koszul cohomology groups give rise to the *minimal graded free S -resolution* of $\Gamma_X(\mathcal{F}, L)$: this is a complex F_\bullet of graded S -modules, unique up to isomorphism, producing an exact sequence

$$\cdots \longrightarrow F_{i+1} \longrightarrow F_i \longrightarrow \cdots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow \Gamma_X(\mathcal{F}, L) \longrightarrow 0,$$

and satisfying $F_i = \bigoplus_{j \geq 0} S(-i-j) \otimes K_{i,j}(X, \mathcal{F}, L)$ for all $i \geq 0$. Here $S(-d)$ denotes a free graded S -module of rank one, generated in degree d . We define the *kernel bundle* M_L associated to the pair (X, L) , as the kernel of the evaluation map on sections, that is, via the exact sequence

$$(49) \quad 0 \longrightarrow M_L \longrightarrow H^0(X, L) \otimes \mathcal{O}_X \longrightarrow L \longrightarrow 0,$$

where the surjectivity on the right follows from the global generation of L . Applying the construction (49) to the pair $(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(1))$, we get a short exact sequence

$$(50) \quad 0 \longrightarrow M_{\mathcal{O}_{\mathbb{P}^r}(1)} \longrightarrow H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(1)) \otimes \mathcal{O}_{\mathbb{P}^r} \longrightarrow \mathcal{O}_{\mathbb{P}^r}(1) \longrightarrow 0.$$

In fact (49) is the pull-back of (50) along φ_L . Moreover, (50) can be identified with the twist by $\mathcal{O}_{\mathbb{P}^r}(1)$ of the Euler sequence on \mathbb{P}^r . The proof of the following result appears in [3, Proposition 2.8].

Lemma 4.1. *Let L be a very ample line bundle on X . Then for $i \geq 0$ we have that*

$$(51) \quad K_{i+1,1}(X, L) \cong \text{Ker} \left\{ H^0(\mathbb{P}^r, \bigwedge^i M_{\mathcal{O}_{\mathbb{P}^r}(1)}(2)) \xrightarrow{\alpha} H^0(X, \bigwedge^i M_L \otimes L^2) \right\},$$

where α is the natural map induced by the identifications $\varphi_L^*(\mathcal{O}_{\mathbb{P}^r}(1)) = L$ and $\varphi_L^*(M_{\mathcal{O}_{\mathbb{P}^r}(1)}) = M_L$.

4.2. Minimizations and mapping cones. Let S be a graded polynomial ring over a field \mathbf{k} . If A is any graded S -module, we write $\bar{A} = A \otimes_S \mathbf{k}$ for the reduction of A modulo the maximal homogeneous ideal. If $\phi : A \rightarrow B$ is a map of S -modules then $\bar{\phi} : \bar{A} \rightarrow \bar{B}$ denotes the induced map $\phi \otimes_S \text{id}_{\mathbf{k}}$. Similarly, if A_\bullet is any complex of graded S -modules then $\bar{A}_\bullet = A_\bullet \otimes_S \mathbf{k}$.

Recall that any complex F_\bullet of finitely-generated, graded, free S -modules is a direct sum of a minimal complex M_\bullet and an exact complex. The complex M_\bullet in question is unique up to isomorphism and is called *the minimization of F_\bullet* . Explicitly, the terms of M_\bullet are computed by

$$(52) \quad M_i = H_i(\bar{F}_\bullet) \otimes_{\mathbf{k}} S = \bigoplus_{j \in \mathbb{Z}} H_i(\bar{F}_\bullet)_j \otimes_{\mathbf{k}} S,$$

where $H_i(\bar{F}_\bullet)_j$ is a graded vector space concentrated in degree j , so that $H_i(\bar{F}_\bullet)_j \otimes_{\mathbf{k}} S$ is a direct sum of copies of $S(-j)$. On several occasions in Section 5, we shall need to compute the minimization of a mapping cone, and in each case the following result will apply.

Lemma 4.2. *Let F_\bullet and G_\bullet be two minimal complexes of finitely-generated, free, graded S -modules, and let $f_\bullet : F_\bullet \rightarrow G_\bullet$ be a morphism of complexes such that for all i , all the entries of the matrix representing f_i are constant. The terms of the minimization M_\bullet of the mapping cone of f_\bullet are $M_i = \text{Ker}(f_{i-1}) \oplus \text{Coker}(f_i)$.*

Proof. We write d_i^F (resp. d_i^G) for the differentials in F_\bullet (resp. G_\bullet), and let $C(f)_\bullet$ denote the mapping cone of f_\bullet . We have that $C(f)_i = F_{i-1} \oplus G_i$, and the i -th differential in $C(f)_\bullet$ is given by the matrix

$$\partial_i = \begin{pmatrix} d_{i-1}^F & 0 \\ f_{i-1} & -d_i^G \end{pmatrix}.$$

Since F_\bullet and G_\bullet are minimal, the reduction of $C(f)_\bullet$ modulo the maximal homogeneous ideal of S yields a complex $\overline{C(f)}_\bullet$, whose terms are $\overline{C(f)}_i = \overline{F}_{i-1} \oplus \overline{G}_i$ and with differentials

$$\overline{\partial}_i = \begin{pmatrix} 0 & 0 \\ \overline{f}_{i-1} & 0 \end{pmatrix}.$$

It follows that $H_i(\overline{C(f)}_\bullet) = \text{Ker}(\overline{f}_{i-1}) \oplus \text{Coker}(\overline{f}_i)$ for all i .

Our assumption on the maps f_i implies that $f_i = \overline{f}_i \otimes_{\mathbf{k}} S$ is obtained by base change from a map of graded vector spaces. Since this base change is flat, we conclude that

$$\text{Ker}(f_i) = \text{Ker}(\overline{f}_i) \otimes_{\mathbf{k}} S \text{ and } \text{Coker}(f_i) = \text{Coker}(\overline{f}_i) \otimes_{\mathbf{k}} S \text{ for all } i.$$

The desired conclusion about the terms M_i follows now from the description (52) of the minimization:

$$M_i = H_i(\overline{C(f)}_\bullet) \otimes_{\mathbf{k}} S = \text{Ker}(\overline{f}_{i-1}) \otimes_{\mathbf{k}} S \oplus \text{Coker}(\overline{f}_i) \otimes_{\mathbf{k}} S = \text{Ker}(f_{i-1}) \oplus \text{Coker}(f_i). \quad \square$$

4.3. The geometric technique for constructing syzygies. Let Y be a projective variety over a field \mathbf{k} , and let V be a finite dimensional \mathbf{k} -vector space. Suppose we have a short exact sequence

$$(53) \quad 0 \longrightarrow \xi \longrightarrow V \otimes \mathcal{O}_Y \longrightarrow \eta \longrightarrow 0,$$

of vector bundles over Y and set $r := \text{rk}(\xi)$. We view $S := \text{Sym}(V)$ as the coordinate ring of the affine space V^\vee . Let $X := \text{Spec}_Y(\mathcal{O}_Y \otimes S) = V^\vee \times Y$ denote the trivial bundle on Y with fiber V^\vee and write $\pi : X \longrightarrow V^\vee$ and $p : X \longrightarrow Y$ for the projections. We consider the sheaf $\mathcal{S} := \text{Sym}_{\mathcal{O}_Y}(\eta)$, and let

$$Z := \text{Spec}_Y(\mathcal{S}),$$

be the total space of the vector bundle associated with η . The surjection in (53) makes Z a subbundle of the trivial bundle X and we get the following commutative diagram:

$$(54) \quad \begin{array}{ccccc} Z & \hookrightarrow & X & \xrightarrow{p} & Y \\ \downarrow \pi|_Z & & \downarrow \pi & & \\ \pi(Z) & \hookrightarrow & V^\vee & & \end{array}$$

The sheaf \mathcal{O}_Z is resolved over X by the following Koszul complex

$$\mathcal{K}(\xi)_\bullet : \quad 0 \longrightarrow \bigwedge^r p^*(\xi) \longrightarrow \cdots \longrightarrow p^*(\xi) \longrightarrow \mathcal{O}_X \left[\longrightarrow \mathcal{O}_Z \longrightarrow 0 \right].$$

If \mathcal{V} denotes a locally free sheaf on Y , we can tensor $\mathcal{K}(\xi)_\bullet$ with $p^*(\mathcal{V})$ to obtain a locally free \mathcal{O}_X -resolution of $\mathcal{M}(\mathcal{V}) := p^*(\mathcal{V}) \otimes \mathcal{O}_Z$, given by

$$\mathcal{K}(\xi, \mathcal{V})_\bullet : \quad 0 \longrightarrow p^*(\mathcal{V}) \otimes \bigwedge^r p^*(\xi) \longrightarrow \cdots \longrightarrow p^*(\mathcal{V}) \otimes p^*(\xi) \longrightarrow p^*(\mathcal{V}) \left[\longrightarrow \mathcal{M}(\mathcal{V}) \longrightarrow 0 \right].$$

The derived direct image of $\mathcal{M}(\mathcal{V})$ along π , or equivalently, that of the complex $\mathcal{K}(\xi, \mathcal{V})_\bullet$, is represented by a minimal complex of graded S -modules constructed as follows (see [31, Theorem 5.1.2]).

Theorem 4.3. *Let \mathcal{V} be a locally free sheaf on Y and set $\mathcal{M}(\mathcal{V}) := p^*(\mathcal{V}) \otimes \mathcal{O}_Z$. There exists a minimal complex $F(\mathcal{V})_\bullet$ of graded S -modules, whose terms are defined by*

$$F(\mathcal{V})_i = \bigoplus_{j \geq 0} H^{i+j}(Y, \bigwedge^j \xi \otimes \mathcal{V}) \otimes_{\mathbf{k}} S(-i-j) \text{ for all } i \in \mathbb{Z}.$$

This complex is exact in positive homological degrees, and its homology groups in negative degrees are given by

$$H_{-i}(F(\mathcal{V})_\bullet) = H^i(Z, \mathcal{M}(\mathcal{V})) = \bigoplus_{d \geq 0} H^i(Y, \mathcal{V} \otimes \text{Sym}^d \eta) \quad \text{for all } i \geq 0.$$

If $\mathcal{M}(\mathcal{V})$ has vanishing higher cohomology, then the complex $F(\mathcal{V})_\bullet$ gives the minimal free resolution of $H^0(Z, \mathcal{M}(\mathcal{V}))$, as will be the case in the proof of Propositions 5.6. We also apply Theorem 4.3 in the proof of Proposition 5.8, where we construct a complex with two non-vanishing homology groups.

5. SYZYGIES OF THE TANGENT DEVELOPABLE TO A RATIONAL NORMAL CURVE

The goal of this section is to establish a link, following [7], between the syzygies of the tangent developable \mathcal{T} to a rational normal curve of degree g , and Koszul modules of a particular kind. This link, together with the general results on finite length Koszul modules that we established in Section 2, allows us to completely characterize the (non-)vanishing behavior for the syzygies of \mathcal{T} .

Throughout this section we fix a field \mathbf{k} , a 2-dimensional \mathbf{k} -vector space U and set $\mathbf{P}^1 := \mathbb{P}(U^\vee)$. We fix $g \geq 3$, let $\mathbf{P}^g := \mathbb{P}(D^g(U^\vee))$, and consider as in Section 3.2 the degree g Veronese embedding $\nu : \mathbf{P}^1 \rightarrow \mathbf{P}^g$, whose image is the *rational normal curve* Γ of degree g . We let $\mathcal{T} \subseteq \mathbf{P}^g$ denote the *tangential variety* (or *tangent developable*) of the curve Γ , and set

- $S := \text{Sym}(\text{Sym}^g U) \cong \text{Sym } H^0(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1))$, the homogeneous coordinate ring of \mathbf{P}^g .
- $I := I_{\mathcal{T}} \subseteq S$, the homogeneous ideal defining \mathcal{T} .
- $R := S/I$, the homogeneous coordinate ring of \mathcal{T} .

The goal of this section is to describe the minimal free resolution of R as an S -module, and in particular to establish Theorem 1.1. Our first result describes the shape of this minimal resolution.

Theorem 5.1. *If $\text{char}(\mathbf{k}) \neq 2$ then R is Gorenstein with Castelnuovo–Mumford regularity $\text{reg}(R) = 3$.*

Since R is the homogeneous coordinate ring of the surface \mathcal{T} , it follows that $\dim(R) = 3$. Theorem 5.1 implies then that the Betti table of the S -module R is described by (8), where

$$b_{i,j} := \dim K_{i,j}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)).$$

Moreover, the Gorenstein property implies that $b_{i,1} = b_{g-2-i,2}$, for $i = 1, \dots, g-3$. Based on this equality, we characterize the (non-)vanishing behavior of the Betti numbers of \mathcal{T} .

Theorem 5.2. *Let $p = \text{char}(\mathbf{k}) \neq 2$. If $p = 0$ or $p \geq \frac{g+2}{2}$, then*

$$(55) \quad b_{i,2} \neq 0 \iff \frac{g-2}{2} \leq i \leq g-3.$$

If $3 \leq p \leq \frac{g+1}{2}$, then

$$(56) \quad b_{i,2} \neq 0 \iff p-2 \leq i \leq g-3.$$

The vanishing Theorem 1.1 in the Introduction is now a direct consequence of Theorem 5.2. Indeed, it can be rephrased as $b_{\lfloor \frac{g}{2} \rfloor, 1} = 0$ if $p = \text{char}(\mathbf{k})$ satisfies $p = 0$ or $p \geq \frac{g+2}{2}$. Using the Gorenstein property of R and that $g-2 - \lfloor \frac{g}{2} \rfloor = \lfloor \frac{g-3}{2} \rfloor$, we have $b_{\lfloor \frac{g}{2} \rfloor, 1} = b_{\lfloor \frac{g-3}{2} \rfloor, 2}$. The vanishing of $b_{\lfloor \frac{g-3}{2} \rfloor, 2}$ follows from (55).

The key to proving Theorem 5.2 is to interpret the Koszul cohomology groups $K_{i,2}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1))$ in the framework of Koszul modules. To that end, we consider for $a \geq 1$ the \mathfrak{sl}_2 -equivariant map (37), which is an inclusion if $\text{char}(\mathbf{k}) \neq 2$ (see Lemma 3.1), in which case we denote by $W^{(a)}$ the associated Koszul module $W(D^a U, D^{2a-2} U)$ (see (58)). This module was considered in [7, Section 3.I.B] and was called a *Weyman module*. The following is the crucial result of this section.

Theorem 5.3. *If $\text{char}(\mathbf{k}) \neq 2$, then for each $i = 1, \dots, g - 3$, we have an identification*

$$K_{i,2}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) = W_{g-3-i}^{(i+2)}.$$

Our analysis of the syzygies of \mathcal{T} is for the most part characteristic-free, and we will be careful to point out when the assumption that $\text{char}(\mathbf{k}) \neq 2$ is used (see Section 5.9 for the case $\text{char}(\mathbf{k}) = 2$). In particular, our analysis yields the following characteristic-free description of the groups $K_{i,1}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1))$.

Theorem 5.4. *If $\text{char}(\mathbf{k})$ is arbitrary, then for $i = 0, \dots, g - 2$, we have an identification*

$$(57) \quad K_{i,1}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) = \text{Ker} \left\{ \delta_2 : D^{2i} U \otimes \text{Sym}^{g-2-i}(D^{i+1} U) \longrightarrow D^{i+1} U \otimes \text{Sym}^{g-1-i}(D^{i+1} U) \right\},$$

where δ_2 is the composition of the Koszul differential

$$\bigwedge^2 (D^{i+1} U) \otimes \text{Sym}^{g-2-i}(D^{i+1} U) \longrightarrow D^{i+1} U \otimes \text{Sym}^{g-1-i}(D^{i+1} U)$$

with the map induced by (37)

$$D^{2i} U \otimes \text{Sym}^{g-2-i}(D^{i+1} U) \longrightarrow \bigwedge^2 (D^{i+1} U) \otimes \text{Sym}^{g-2-i}(D^{i+1} U).$$

Before delving into the proofs, we summarize the contents of this section. We begin by defining Weyman modules in Section 5.1, and explaining how Theorem 5.2 follows from Theorems 5.1, 5.3, and from the results in Section 2. In Section 5.2 we give a roadmap to the proofs of Theorems 5.1, 5.3, and 5.4, which occupy the rest of the section. In Section 5.3, we summarize the \mathfrak{sl}_2 description of cohomology of line bundles on \mathbf{P}^1 , which will be used throughout. In Section 5.4 we explain how to parametrize the affine cone over \mathcal{T} using the bundle of principal parts. Sections 5.5 – 5.7 describe the technical parts of our arguments: they rely heavily on the preliminaries discussed in Sections 3 and 4, particularly on Hermite reciprocity and the Kempf–Weyman technique, and they conclude with a proof of Theorem 5.4. Using the foundation laid in the preceding sections, we give a quick proof of Theorems 5.1 and 5.3 in Section 5.8. We end in Section 5.9 with a concrete discussion of the resolution of \mathcal{T} in characteristic 2.

5.1. Weyman modules in characteristic different from 2. For $n \geq 2$ we define

$$(58) \quad W^{(n-1)} := W(D^{n-1} U, D^{2n-4} U)$$

where $D^{2n-4} U$ is regarded as a subspace of $\bigwedge^2(D^{n-1} U)$ via the map Δ_1 (see (37) and Lemma 3.1 with $a = n - 1$). We define $\mathcal{R}^{(n-1)}$ to be the resonance variety $\mathcal{R}(D^{n-1} U, D^{2n-4} U)$ (see (5)).

Lemma 5.5. *Let $p = \text{char}(\mathbf{k})$. If $p = 0$ or $p \geq n$, then $\mathcal{R}^{(n-1)} = \{0\}$. If $3 \leq p \leq n - 1$ then $\mathcal{R}^{(n-1)} \neq \{0\}$.*

Proof. Consider the map $\mu_1 : \bigwedge^2 \text{Sym}^{n-1}(U^\vee) \longrightarrow \text{Sym}^{2n-4}(U^\vee)$ (defined in (36) and Lemma 3.1), which is dual to Δ_1 . The condition $\mathcal{R}^{(n-1)} \neq \{0\}$ is equivalent to the fact that $\text{Ker}(\mu_1)$ contains a non-zero decomposable form $f_1 \wedge f_2$. We choose a basis $(1, y)$ for U^\vee , so that for each $d \geq 0$ we can identify $\text{Sym}^d(U^\vee)$ with the space of polynomials of degree at most d in y .

If $3 \leq p \leq n - 1$ then $f_1 = 1$ and $f_2 = y^p$ belong to $\text{Sym}^{n-1}(U^\vee)$, and $\mu_1(f_1 \wedge f_2) = \mu_1(1 \wedge y^p) = 0$. Since $1 \wedge y^p \neq 0$, we conclude using (5) that $\mathcal{R}^{(n-1)} \neq \{0\}$. Suppose now by contradiction that $\text{Ker}(\mu_1)$ contains a non-zero decomposable form $f_1 \wedge f_2$, and that $p = 0$ or $p \geq n$. By rescaling $f_1 \wedge f_2$, we may assume that f_1, f_2 are monic polynomials in y , say

$$f_1 = y^r + b_{r-1}y^{r-1} + \dots + b_0, \quad f_2 = y^s + c_{s-1}y^{s-1} + \dots + c_0.$$

If $r \neq s$, we may assume $r > s$ and we get $\mu_1(f_1 \wedge f_2) = (r-s)y^{r+s-1} + h(y)$, where $\deg(h) < r+s-1$ and $0 < r-s \leq n-1$. It follows that p cannot divide $r-s$, hence $\mu_1(f_1 \wedge f_2) \neq 0$, a contradiction. Suppose finally that $r = s$, and let i be the maximal index for which $b_i \neq c_i$, which exists since $f_1 \wedge f_2 \neq 0$. We get

$$\mu_1(f_1 \wedge f_2) = (r-i) \cdot (b_i - c_i) \cdot y^{r+i-1} + h(y), \text{ where } \deg(h) < r+i-1.$$

Since $0 < r-i \leq n-1$ and $b_i - c_i \neq 0$, it follows again that $f_1 \wedge f_2 \notin \text{Ker}(\mu_1)$, yielding a contradiction. \square

We can now give a quick proof of Theorem 5.2, assuming the statements of Theorems 5.1 and 5.3.

Proof of Theorem 5.2. By Theorem 5.1, the only potentially non-zero $b_{i,2}$ occur for $1 \leq i \leq g-3$, so we fix one such index i . Since $p \neq 2$, Theorem 5.3 applies to give

$$(59) \quad b_{i,2} = \dim(W_{g-3-i}^{(i+2)}).$$

Let $n = i+3$, so that $W^{(i+2)} = W^{(n-1)}$, to which we apply Lemma 5.5. Suppose first $3 \leq p \leq n-1 = i+2$, so the support $\mathcal{R}^{(n-1)}$ of $W^{(n-1)}$ is positive dimensional, in particular $W_q^{(n-1)} \neq 0$ for all $q \geq 0$. Taking $q = g-3-i$ we obtain $b_{i,2} \neq 0$, in particular the implication “ \Leftarrow ” in (56) holds when $3 \leq p \leq \frac{g+1}{2}$. If $p \geq \frac{g+2}{2}$, using the assumption $p \leq i+2$, we conclude that $i \geq \frac{g-2}{2}$, so conditioned on $3 \leq p \leq i+2$, the equivalence in (55) holds.

Suppose now that $p = 0$ or $p \geq n = i+3$, so that $\mathcal{R}^{(n-1)} = \{0\}$. From Theorems 1.3 and 1.4 we obtain the equivalence $W_q^{(n-1)} \neq 0$ if and only if $0 \leq q \leq n-4$, which, in view of (59), implies that

$$(60) \quad b_{i,2} \neq 0 \iff 0 \leq g-3-i \leq i-1 \iff \frac{g-2}{2} \leq i \leq g-3.$$

If $3 \leq p \leq \frac{g+1}{2}$ then $i \leq p-3 \leq \frac{g-5}{2}$, proving based on (60) that $b_{i,2} = 0$ and establishing the remaining part of the equivalence (56). If $p = 0$ or $p \geq \frac{g+2}{2}$ then (60) agrees with (55). \square

5.2. The roadmap to the minimal free resolution of R . For the convenience of the reader, we summarize here our analysis of the minimal free resolution of R as an S -module, which concludes with the proofs of Theorems 5.1, 5.3, and 5.4. The key players in our proof are four minimal complexes of free graded S -modules, denoted $F_\bullet, J_\bullet, C_\bullet$ and K_\bullet , the first two of which are constructed via the Kempf–Weyman geometric technique (see Section 4.3). For visualization purposes, we record their Betti diagrams below: our convention is that for a complex A_\bullet , a “ $*$ ” in row j and column i indicates that the free module A_i has generators of degree $i+j$, while a “ $-$ ” indicates that no such generators exist. We will also write “1” instead of “ $*$ ” when $A_i \cong S(-i-j)$ is a free module of rank one, generated in degree $i+j$. With these conventions, our four complexes take the following shapes:

$$\begin{array}{c}
 F_\bullet : \quad \begin{array}{c|cccccc}
 & 0 & 1 & 2 & \cdots & g-3 & g-2 \\
 0 & 1 & - & - & \cdots & - & - \\
 1 & * & * & * & \cdots & * & *
 \end{array}
 \end{array}
 \quad
 \begin{array}{c}
 C_\bullet : \quad \begin{array}{c|ccccccc}
 & 0 & 1 & 2 & \cdots & g-3 & g-2 & g-1 \\
 0 & - & - & - & \cdots & - & - & - \\
 1 & * & * & * & \cdots & * & * & - \\
 2 & - & - & - & \cdots & - & - & 1
 \end{array}
 \end{array}$$

$$\begin{array}{c}
 J_\bullet : \quad \begin{array}{c|cccccc}
 & 0 & 1 & 2 & \cdots & g-1 & g \\
 0 & 1 & * & * & \cdots & * & * \\
 1 & * & * & * & \cdots & * & *
 \end{array}
 \end{array}
 \quad
 \begin{array}{c}
 K_\bullet : \quad \begin{array}{c|cccccc}
 & 0 & 1 & 2 & \cdots & g-1 & g \\
 0 & 1 & * & * & \cdots & * & 1
 \end{array}
 \end{array}$$

The complex F_\bullet . The affine cone $\widehat{\mathcal{T}}$ over \mathcal{T} admits a dominant map from the total space Z of the bundle of principal parts associated with the sheaf $\mathcal{O}_{\mathbb{P}^1}(g)$ (see Section 5.4). As such, R is naturally a subring of

$\tilde{R} = H^0(Z, \mathcal{O}_Z)$. The complex F_\bullet gives the minimal free resolution of \tilde{R} (see Section 5.5). In characteristic different from two, the map $Z \rightarrow \hat{\mathcal{T}}$ is birational, and \tilde{R} is the normalization of R .

The complex C_\bullet . The affine cone $\hat{\Gamma}$ over the rational normal curve is Cohen–Macaulay, so it has a canonical module which we denote by C . The complex C_\bullet is the minimal resolution of C (see Section 5.6).

The complex J_\bullet . The complex J_\bullet is the only one that is not a resolution (see Proposition 5.8): it is a linear complex with two non-zero cohomology groups, namely $H_0(J_\bullet) = \mathbf{k}$, and $H_1(J_\bullet) = C$.

The complex K_\bullet . The complex K_\bullet is the Koszul complex resolving the residue field $\mathbf{k} \cong S/\mathfrak{m}$.

We now describe the various links between the complexes F_\bullet , J_\bullet , C_\bullet and K_\bullet .

The relationship between C_\bullet , J_\bullet and K_\bullet . There is a degree preserving map $p_\bullet : J_\bullet \rightarrow K_\bullet$, constructed in Proposition 5.9, inducing an isomorphism $\mathbf{k} = H_0(J_\bullet) \cong H_0(K_\bullet) = \mathbf{k}$. The mapping cone of p_\bullet gives rise after minimization (as in Section 4.2) and a homological shift, to a minimal resolution of C , and is therefore quasi-isomorphic to C_\bullet . Precisely, $C_i = \text{Ker}(p_{i+1})$ for $i = 0, \dots, g-2$ (see Corollary 5.10).

The relationship between F_\bullet , J_\bullet and (57). There is a map $q_\bullet : F_\bullet \rightarrow J_{\bullet+1}$ of complexes, which is constructed in Proposition 5.12, inducing a surjection $\tilde{R} = H_0(F_\bullet) \twoheadrightarrow H_1(J_\bullet) = C$. Using the explicit Hermite reciprocity discussed in Section 3.4, it follows that if we restrict the map q_i to the minimal generators of degree $i+1$ of F_i and J_{i+1} , then we get precisely the map δ_2 in (57), so

$$(61) \quad \text{Ker}(\delta_2) = \text{Ker}((q_i \otimes \mathbf{k})_{i+1}) \text{ for } i = 0, \dots, g-2.$$

The relationship between q_\bullet and the first linear strand of the resolution of R . The morphism q_0 sends the unique generator of F_0 of degree 0 to zero, and therefore R is contained in $R' := \text{Ker}(\tilde{R} \rightarrow C)$. The quotient R'/R is generated in degree ≥ 2 , so the resolutions of R and R' have the same first linear strand. To construct the minimal resolution of R' , we show in Corollary 5.13 that $q_i \circ p_{i+1} = 0$ for $i = 0, \dots, g-2$, so q_i sends F_i into $C_i \subseteq J_{i+1}$. By restricting the range, this induces a map of complexes $\tilde{q}_\bullet : F_\bullet \rightarrow C_\bullet$, lifting the quotient map $\tilde{R} \rightarrow C$. Minimizing the mapping cone of \tilde{q}_\bullet as in Lemma 4.2, we get a minimal resolution of R' , with

$$(62) \quad \text{Tor}_i^S(R', \mathbf{k})_{i+1} = \text{Ker}((\tilde{q}_i \otimes \mathbf{k})_{i+1}), \quad \text{Tor}_i^S(R', \mathbf{k})_{i+2} = \text{Coker}((\tilde{q}_{i+1} \otimes \mathbf{k})_{i+2}).$$

We conclude in Section 5.7 that

$$K_{i,1}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) = \text{Tor}_i^S(R, \mathbf{k})_{i+1} = \text{Tor}_i^S(R', \mathbf{k})_{i+1} = \text{Ker}((\tilde{q}_i \otimes \mathbf{k})_{i+1}) = \text{Ker}((q_i \otimes \mathbf{k})_{i+1}).$$

It then follows from (61) that $K_{i,1}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) = \text{Ker}(\delta_2)$ as in (57), proving Theorem 5.4.

The minimal resolution of R for $\text{char}(\mathbf{k}) \neq 2$. By construction, we have $\text{Coker}(\tilde{q}_i) = \text{Ker}(p_{i+1})/\text{Im}(q_i)$, which allows us to realize $\text{Coker}(\tilde{q}_i)$ as the middle homology of a 3-term complex

$$F_i \xrightarrow{q_i} J_{i+1} \xrightarrow{p_{i+1}} K_{i+1}.$$

Restricting this sequence to minimal generators of degree $(i+1)$ we obtain via Hermite reciprocity the 3-term complex defining the degree $g-2-i$ component of the Weyman module $W^{(i+1)}$, and conclude in Lemma 5.14 that $\text{Coker}(\tilde{q}_i) \otimes_S \mathbf{k} = W_{g-2-i}^{(i+1)}$. Taking $i = 0$ we find that \tilde{q}_1 is surjective, and deduce that $R' = R$ in Corollary 5.15. Combining this with (62) proves Theorem 5.3. To prove Theorem 5.1 one is left with verifying that \tilde{q}_{g-2} is injective, which follows from one final application of Hermite reciprocity and is explained at the end of Section 5.8.

The minimal resolution of R for $\text{char}(\mathbf{k}) = 2$. It follows from the discussion in Section 5.4 that \mathcal{T} is a variety of minimal degree in characteristic two. We show in Section 5.9 that it is a rational normal scroll, and conclude that its minimal resolution is given by an Eagon–Northcott complex.

5.3. Cohomology on \mathbf{P}^1 . Let U be a two dimensional vector space as before and recall that $\mathbf{P}^1 = \mathbb{P}(U^\vee)$. The tautological quotient map $U \otimes \mathcal{O}_{\mathbf{P}^1} \rightarrow \mathcal{O}_{\mathbf{P}^1}(1)$ gives rise to an exact Koszul complex

$$(63) \quad 0 \longrightarrow \bigwedge^2 U \otimes \mathcal{O}_{\mathbf{P}^1}(-1) \longrightarrow U \otimes \mathcal{O}_{\mathbf{P}^1} \longrightarrow \mathcal{O}_{\mathbf{P}^1}(1) \longrightarrow 0,$$

and using (49) we get a natural identification $M_{\mathcal{O}_{\mathbf{P}^1}(1)} \cong \bigwedge^2 U \otimes \mathcal{O}_{\mathbf{P}^1}(-1)$. The canonical bundle $\omega_{\mathbf{P}^1}$ is naturally identified with $\bigwedge^2 U \otimes \mathcal{O}_{\mathbf{P}^1}(-2)$. The cohomology groups of twists of $\mathcal{O}_{\mathbf{P}^1}$ are then given by:

$$H^0(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(a)) = \begin{cases} \text{Sym}^a U & \text{if } a \geq 0, \\ 0 & \text{otherwise,} \end{cases} \quad H^1(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(a)) = \begin{cases} \bigwedge^2 U^\vee \otimes D^{-a-2}(U^\vee) & \text{if } a \leq -2, \\ 0 & \text{otherwise.} \end{cases}$$

Thinking of (63) as a two-step resolution of $\mathcal{O}_{\mathbf{P}^1}(1)$, the sheaf $\mathcal{O}_{\mathbf{P}^1}(a) \cong \text{Sym}^a(\mathcal{O}_{\mathbf{P}^1}(1))$ is then resolved by the a -th symmetric power of the said two-step resolution. This gives an exact sequence

$$(64) \quad 0 \longrightarrow \text{Sym}^{a-1} U \otimes \bigwedge^2 U \otimes \mathcal{O}_{\mathbf{P}^1}(-1) \xrightarrow{\iota_a} \text{Sym}^a U \otimes \mathcal{O}_{\mathbf{P}^1} \longrightarrow \mathcal{O}_{\mathbf{P}^1}(a) \longrightarrow 0,$$

where the second map is given by evaluation of sections, and the first is induced by the inclusion in (63), and by the multiplication $\mu : \text{Sym}^{a-1} U \otimes U \rightarrow \text{Sym}^a U$. Using (49) we get from (64) an identification

$$(65) \quad M_{\mathcal{O}_{\mathbf{P}^1}(a)} \cong \text{Sym}^{a-1} U \otimes \bigwedge^2 U \otimes \mathcal{O}_{\mathbf{P}^1}(-1).$$

5.4. Jet bundles and the Gauss map. Recall that Γ is the rational normal curve of degree g in \mathbf{P}^g , obtained as the image of the embedding of \mathbf{P}^1 via the complete linear system $|\mathcal{O}_{\mathbf{P}^1}(g)|$, and that $\mathcal{T} \subseteq \mathbf{P}^g$ denotes the tangential variety of Γ . The first jet bundle $\mathcal{J} := \mathcal{P}^1(\mathcal{O}_{\mathbf{P}^1}(g))$ has the explicit description

$$(66) \quad \mathcal{J} = \begin{cases} U \otimes \mathcal{O}_{\mathbf{P}^1}(g-1) & \text{if } p \nmid g; \\ \bigwedge^2 U \otimes \mathcal{O}_{\mathbf{P}^1}(g-2) \oplus \mathcal{O}_{\mathbf{P}^1}(g) & \text{if } p|g. \end{cases}$$

The vector bundle \mathcal{J} sits in an exact sequence

$$(67) \quad 0 \longrightarrow \left(\bigwedge^2 U\right)^{\otimes 2} \otimes \text{Sym}^{g-2} U \otimes \mathcal{O}_{\mathbf{P}^1}(-2) \xrightarrow{\iota} \text{Sym}^g U \otimes \mathcal{O}_{\mathbf{P}^1} \longrightarrow \mathcal{J} \longrightarrow 0,$$

where $\left(\bigwedge^2 U\right)^{\otimes 2} \otimes \text{Sym}^{g-2} U \otimes \mathcal{O}_{\mathbf{P}^1}(-2) \cong \mathcal{N}_{\Gamma/\mathbf{P}^g}^\vee(g)$, see for instance [5, Corollary 2.5] for a characteristic free calculation. The inclusion map ι in (67) is induced by composing the map ι_g in (64) with $\text{id}_{\bigwedge^2 U} \otimes \iota_{g-1}(-1)$. We denote by $\nu : \mathbf{P}(\mathcal{J}) \rightarrow \mathbf{P}^g$ the map induced by the subspace of sections

$$\text{Sym}^g U \subseteq H^0(\mathbf{P}^1, \mathcal{J}) \cong H^0(\mathbf{P}(\mathcal{J}), \mathcal{O}_{\mathbf{P}(\mathcal{J})}(1)).$$

The image of ν is the tangential variety \mathcal{T} and τ is a resolution of singularities if $\text{char}(\mathbf{k}) \neq 2$. Let

$$(68) \quad \mathcal{S} := \text{Sym}_{\mathcal{O}_{\mathbf{P}^1}}(\mathcal{J}) \quad \text{and} \quad Z := \underline{\text{Spec}}_{\mathbf{P}^1}(\mathcal{S}) \hookrightarrow D^g(U^\vee) \times \mathbf{P}^1 =: X.$$

If we let \mathbb{G} denote the Grassmannian of two dimensional subspaces of $D^g(U^\vee)$, then the surjection in (67) gives rise to the *Gauss map* $\gamma : \Gamma \rightarrow \mathbb{G}$, whose image we denote by Γ^* . We have a diagram (see [21])

$$(69) \quad \begin{array}{ccccc} \Gamma \cong \mathbf{P}^1 & \xleftarrow{\rho} & Z & \hookrightarrow & X \\ \downarrow \gamma & & \downarrow \pi|_Z & & \downarrow \pi \\ \mathbb{G} & \longleftarrow & \Gamma^* & & \widehat{\mathcal{T}} \hookrightarrow D^g(U^\vee) \end{array}$$

where $\widehat{\mathcal{T}}$ is the affine cone over \mathcal{T} . We recall that $R = H^0(\widehat{\mathcal{T}}, \mathcal{O}_{\widehat{\mathcal{T}}})$ denotes the homogeneous coordinate ring of \mathcal{T} , and let $\widetilde{R} = H^0(Z, \mathcal{O}_Z)$. Since π maps Z onto $\widehat{\mathcal{T}}$, the pull-back along $\pi|_Z$ leads to an inclusion of R into \widetilde{R} , which will be analyzed later on. Applying [21, Lemma 1.6, Corollary 1.7, Remark 1.9], we conclude that $\deg(\mathcal{T}) = \deg(\Gamma^*)$ and $\deg(\gamma) = \deg(\pi|_Z)$. Moreover, γ can be written in an affine chart as

$$(70) \quad t \xrightarrow{\gamma} \text{row Span} \begin{bmatrix} 1 & t & t^2 & \cdots & t^g \\ 0 & 1 & 2t & \cdots & gt^{g-1} \end{bmatrix}.$$

Suppose first that $\text{char}(\mathbf{k}) \neq 2$. Since the 2×2 minors of the above matrix span the space of polynomials in t of degree up to $2g - 2$, it follows that when viewed in its Plücker embedding, Γ^* is a rational normal curve (in its span) of degree $2g - 2$. Moreover, the map γ is an isomorphism. We conclude that $\deg(\Gamma^*) = 2g - 2$ and $\deg(\gamma) = 1$, and therefore $\deg(\mathcal{T}) = 2g - 2$ and $\pi|_Z$ is birational. It follows that the map $\pi|_Z$ gives a resolution of singularities for $\widehat{\mathcal{T}}$, and in particular \widetilde{R} is the normalization of R .

Suppose now that $\text{char}(\mathbf{k}) = 2$. The 2×2 minors of (70) span the space of polynomials in t^2 of degree $g - 1$, so γ can be regarded as the composition of the Frobenius map with a Veronese embedding of degree $g - 1$. It follows that $\deg(\mathcal{T}) = \deg(\Gamma^*) = g - 1$ and $\deg(\pi|_Z) = \deg(\gamma) = 2$.

5.5. The resolution of \widetilde{R} . Recall that $S = \text{Sym}(\text{Sym}^g U) \cong \text{Sym} H^0(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1))$ denotes the homogeneous coordinate ring of \mathbf{P}^g , let X, Z as in (68), and let $\widetilde{R} := H^0(Z, \mathcal{O}_Z) = \bigoplus_{n \geq 0} H^0(\mathbf{P}(\mathcal{J}), \mathcal{O}_{\mathbf{P}(\mathcal{J})}(n))$. We describe the minimal free resolution of \widetilde{R} as an S -module.

Proposition 5.6. *The S -module $\widetilde{R} := H^0(Z, \mathcal{O}_Z)$ has an \mathfrak{sl}_2 -equivariant minimal free resolution given by*

$$F_\bullet : \quad F_{g-2} \longrightarrow \cdots \longrightarrow F_1 \longrightarrow F_0 \longrightarrow \widetilde{R} \longrightarrow 0,$$

where $F_0 = S \oplus \text{Sym}^{g-2} U \otimes S(-1)$, and

$$F_i = \left(D^{2i} U \otimes \bigwedge^{i+1} \text{Sym}^{g-2} U \right) \otimes S(-i-1) \text{ for } i = 1, \dots, g-2.$$

The linear part of the differential $\partial_i^F : F_i \rightarrow F_{i-1}$ is defined via an \mathfrak{sl}_2 -equivariant map

$$D^{2i} U \otimes \bigwedge^{i+1} \text{Sym}^{g-2} U \longrightarrow \left(D^{2i-2} U \otimes \bigwedge^i \text{Sym}^{g-2} U \right) \otimes \text{Sym}^g U$$

induced by

$$\Delta^{(2)} : D^{2i} U \xrightarrow{(34)} D^{2i-2} U \otimes \text{Sym}^2 U, \quad k_{i+1}^{g-2} : \bigwedge^{i+1} \text{Sym}^{g-2} U \xrightarrow{(46)} \bigwedge^i \text{Sym}^{g-2} U \otimes \text{Sym}^{g-2} U,$$

$$\text{and } \mu : \text{Sym}^2 U \otimes \text{Sym}^{g-2} U \xrightarrow{(32)} \text{Sym}^g U.$$

If we ignore the quadratic part of the differential $\partial_1^F : F_1 \rightarrow F_0$, then we have the formula

$$(71) \quad \partial_i^F(x^{(t)} \otimes s_\lambda(x)) = \sum_{\substack{u+s=t \\ u \leq 2, s \leq 2i-2}} \sum_{j=1}^{i+1} (-1)^{j-1} \cdot x^{(s)} \otimes s_{\lambda^j}(x) \otimes \left(\binom{2}{u} \cdot x^u \cdot x^{\lambda_j+i+1-j} \right)$$

for all $0 \leq t \leq 2i$ and $\lambda \in \mathcal{P}_{i+1}(g-2-i)$.

Remark 5.7. In the language of Koszul cohomology, Proposition 5.6 yields the \mathfrak{sl}_2 -identification

$$K_{i,1}(\mathcal{T}, \tau_* \mathcal{O}_{\mathcal{P}(\mathcal{J})}, \mathcal{O}_{\mathcal{T}}(1)) \cong D^{2i}U \otimes \bigwedge^{i+1} \text{Sym}^{g-2}U, \text{ for } i = 1, \dots, g-2.$$

Proof. We let $\xi = (\bigwedge^2 U)^{\otimes 2} \otimes \text{Sym}^{g-2}U \otimes \mathcal{O}_{\mathbf{P}^1}(-2)$, $V = \text{Sym}^g U$ and $\eta = \mathcal{J}$, so that (67) is a special instance of (53). We let $\mathcal{V} = \mathcal{O}_{\mathbf{P}^1}$ and apply Theorem 4.3 to construct a complex F_\bullet with

$$(72) \quad F_i = \bigoplus_{j \geq 0} H^j(\mathbf{P}^1, \bigwedge^{i+j} \xi) \otimes S(-i-j).$$

It follows from (66) and Section 5.3 that $\mathcal{M}(\mathcal{V}) = \mathcal{O}_Z$ has no higher cohomology, so F_\bullet is a minimal free resolution of \tilde{R} . We have that $\bigwedge^{i+j} \xi = (\bigwedge^2 U)^{\otimes(2i+2j)} \otimes \bigwedge^{i+j} \text{Sym}^{g-2}U \otimes \mathcal{O}_{\mathbf{P}^1}(-2i-2j)$ and therefore

$$(73) \quad H^j(\mathbf{P}^1, \bigwedge^{i+j} \xi) = \begin{cases} \mathbf{k} & \text{if } i = j = 0; \\ \bigwedge^2 U \otimes D^{2i}U \otimes \bigwedge^{i+1} \text{Sym}^{g-2}U & \text{if } 1 \leq i \leq g-2 \text{ and } j = 1; \\ 0 & \text{otherwise.} \end{cases}$$

Indeed, if $i+j=0$ then $\bigwedge^{i+j} \xi = \mathcal{O}_{\mathbf{P}^1}$, whose only non-vanishing cohomology is $H^0(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}) = \mathbf{k}$. Since $\text{rk}(\xi) = g-1$, we may assume that $0 < i+j \leq g-1$, and since $\mathcal{O}_{\mathbf{P}^1}(-2i-2j)$ has no global sections, we may assume that $j=1$. It follows from Section 5.3 that

$$H^1(\mathbf{P}^1, \bigwedge^{i+1} \xi) = (\bigwedge^2 U)^{\otimes(2i+2)} \otimes \bigwedge^{i+1} \text{Sym}^{g-2}U \otimes \bigwedge^2 U^\vee \otimes D^{2i}(U^\vee).$$

Using $\bigwedge^2 U \otimes \bigwedge^2 U^\vee \cong \mathbf{k}$ and $(\bigwedge^2 U)^{\otimes 2i} \otimes D^{2i}(U^\vee) \cong D^{2i}U$, we get (73).

Since we are only interested in the \mathfrak{sl}_2 -structure, we have that $\bigwedge^2 U \cong \mathbf{k}$, so (72) and (73) give the desired formula for the terms in the resolution F_\bullet . The formula (71) follows by combining (32), (33), (34) and (46), and recalling that $x^{(u)} \in \text{Sym}^2 U$ should be interpreted as $\binom{2}{u} \cdot x^u$ for $0 \leq u \leq 2$. \square

Since R is a quotient of S and a subring of \tilde{R} , it is isomorphic to S/I , where I is the image of the quadratic part of the differential $\partial_1^F : F_1 \rightarrow F_0$. It is not clear what this is from the construction above, but the construction tells us that I is generated by quadrics. In characteristic 2, it is easy to identify a subcomplex of F_\bullet which gives a resolution of R , which we do in Section 5.9. In the general case we need one more auxiliary construction.

5.6. The minimal resolution of the canonical module C of the affine cone $\hat{\Gamma}$. We construct a minimal resolution of C as the minimization of a mapping cone. We show that it admits an explicit map from the resolution of \tilde{R} , inducing a surjection $\tilde{R} \twoheadrightarrow C$. In Section 5.8 we show that this map gives rise to an isomorphism $C \cong \tilde{R}/R$ provided that $\text{char}(\mathbf{k}) \neq 2$.

Recall from (49) and (65) that $M_{\mathcal{O}_{\mathbf{P}^1}(g)} = \text{Sym}^{g-1} U \otimes \wedge^2 U \otimes \mathcal{O}_{\mathbf{P}^1}(-1)$. We define, in analogy with (68)

$$(74) \quad S' := \text{Sym}_{\mathcal{O}_{\mathbf{P}^1}}(\mathcal{O}_{\mathbf{P}^1}(g)) \quad \text{and} \quad Z' := \underline{\text{Spec}}_{\mathbf{P}^1}(S') \hookrightarrow \mathbb{D}^g(U^\vee) \times \mathbf{P}^1 = X,$$

and we consider the canonical module of $\widehat{\Gamma}$, given as

$$(75) \quad C := H^0(Z', \mathcal{O}_{Z'}(g-2)) = \bigoplus_{n \geq 0} H^0(\Gamma, \omega_\Gamma \otimes \mathcal{O}_\Gamma(n)).$$

We note that C is generated in degree one by $\text{Sym}^{g-2}(U)$. We let C_\bullet denote the minimal free resolution of C , which is dual to that of the homogeneous coordinate ring of Γ . Using [8, Corollary 6.2] it follows that for $i = 0, \dots, g-2$, the free module C_i is generated in degree $i+1$, and $C_{g-1} \cong S(-g-1)$. Our goal is to realize C_\bullet as the minimization of a mapping cone. To that end, we start with the following.

Proposition 5.8. *There exists an \mathfrak{sl}_2 -equivariant complex of free S -modules*

$$J_\bullet : \quad 0 \longrightarrow J_g \longrightarrow \dots \longrightarrow J_1 \longrightarrow J_0, \quad \text{where}$$

$$J_i = \left(D^i U \otimes \bigwedge^i \text{Sym}^{g-1} U \right) \otimes S(-i), \quad \text{for } i = 0, \dots, g,$$

and whose homology is given by $H_0(J_\bullet) = \mathbf{k} \cong S/\mathfrak{m}$, $H_1(J_\bullet) = C$ and $H_i(J_\bullet) = 0$, for $i > 1$. Moreover, the differential $\partial_i^J : J_i \longrightarrow J_{i-1}$ is defined at the level of generators via an \mathfrak{sl}_2 -equivariant map

$$D^i U \otimes \bigwedge^i \text{Sym}^{g-1} U \longrightarrow \left(D^{i-1} U \otimes \bigwedge^{i-1} \text{Sym}^{g-1} U \right) \otimes \text{Sym}^g U$$

induced by

$$\Delta : D^i U \xrightarrow{(33)} D^{i-1} U \otimes U, \quad k_i^{g-1} : \bigwedge^i \text{Sym}^{g-1} U \xrightarrow{(46)} \bigwedge^{i-1} \text{Sym}^{g-1} U \otimes \text{Sym}^{g-1} U,$$

$$\text{and } \mu : U \otimes \text{Sym}^{g-1} U \xrightarrow{(32)} \text{Sym}^g U.$$

More precisely, for all $0 \leq t \leq i$ and $\lambda \in \mathcal{P}_i(g-i)$, we have

$$(76) \quad \partial_i^J(x^{(t)} \otimes s_\lambda(x)) = \sum_{\substack{u+s=t \\ u \leq 1, s \leq i-1}} \sum_{j=1}^i (-1)^{j-1} \cdot x^{(s)} \otimes s_{\lambda^j}(x) \otimes x^{\lambda_j+i-j+u}.$$

Proof. We use (53) with $\xi = M_{\mathcal{O}_{\mathbf{P}^1}(g)}$, $V = \text{Sym}^g U$, $\eta = \mathcal{O}_{\mathbf{P}^1}(g)$, and let $\mathcal{V} = \mathcal{O}_{\mathbf{P}^1}(-2)$. It follows that $\mathcal{M}(\mathcal{V}) = \mathcal{O}_{Z'}(-2)$, whose only non-zero cohomology groups are

$$H^0(X, \mathcal{O}_{Z'}(-2)) = C \quad \text{and} \quad H^1(X, \mathcal{O}_{Z'}(-2)) = H^1(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(-2)) = \bigwedge^2 U^\vee \cong \mathbf{k},$$

where \mathbf{k} is a graded S -module concentrated in degree 0. Theorem 4.3 yields a minimal complex G_\bullet with

$$G_i = \bigoplus_{j \geq 0} H^j(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(-2) \otimes \bigwedge^{i+j} M_{\mathcal{O}_{\mathbf{P}^1}(g)}) \otimes S(-i-j),$$

whose non-zero homology is given by $H_0(G_\bullet) = C$ and $H_{-1}(G_\bullet) = \mathbf{k}$. Note that

$$H^j(\mathbf{P}^1, \mathcal{O}_{\mathbf{P}^1}(-2) \otimes \bigwedge^{i+j} M_{\mathcal{O}_{\mathbf{P}^1}(g)}) = \begin{cases} D^{i+1} U \otimes \bigwedge^{i+1} \text{Sym}^{g-1} U & \text{if } -1 \leq i \leq g-1 \text{ and } j = 1; \\ 0 & \text{otherwise.} \end{cases}$$

In the above calculation we have used the description of cohomology from Section 5.3 and the isomorphism $(\wedge^2 U)^{\otimes i+1} \otimes D^{i+1}(U^\vee) \cong D^{i+1}(U)$. Letting $J_\bullet = G_{\bullet+1}$ we get a complex J_\bullet with the desired properties. We get (76) by combining (32), (33) and (46), and noting that $x^{(u)} = x^u \in U$ for $u = 0, 1$. \square

To build a mapping cone resolution for C , we need a map of complexes between J_\bullet and the Koszul complex resolving the residue field \mathbf{k} . We achieve this as follows.

Proposition 5.9. *Consider the Koszul complex resolving the residue field \mathbf{k} as an S -module*

$$K_\bullet : \quad 0 \longrightarrow K_{g+1} \longrightarrow \cdots \longrightarrow K_1 \longrightarrow K_0 [\longrightarrow \mathbf{k} \longrightarrow 0],$$

where

$$K_i = \left(\wedge^i \text{Sym}^g U \right) \otimes S(-i), \text{ for } i = 0, \dots, g+1,$$

and the differential given by the maps k_i^g in (46). Using the notation (43) and letting $\bar{p}_i = \nu_{g-i}^i$, the maps

$$\bar{p}_i : D^i U \otimes \wedge^i \text{Sym}^{g-1} U \longrightarrow \wedge^i \text{Sym}^g U$$

induce a map of complexes $p_\bullet : J_\bullet \longrightarrow K_\bullet$, with $\bar{p}_i = p_i \otimes \text{id}_{\mathbf{k}}$.

Proof. We have to verify that for each $i = 1, \dots, g+1$ we have a commutative diagram

$$\begin{array}{ccc} D^i U \otimes \wedge^i \text{Sym}^{g-1} U & \xrightarrow{\partial_i^J} & D^{i-1} U \otimes \wedge^{i-1} \text{Sym}^{g-1} U \otimes \text{Sym}^g U \\ \bar{p}_i \downarrow & & \downarrow \bar{p}_{i-1} \otimes \text{id}_{\text{Sym}^g U} \\ \wedge^i \text{Sym}^g U & \xrightarrow{\partial_i^K} & \wedge^{i-1} \text{Sym}^g U \otimes \text{Sym}^g U \end{array}$$

The space $D^i U \otimes \wedge^i \text{Sym}^{g-1} U$ has a basis consisting of elements of the form $x^{(t)} \otimes s_\lambda(x)$, where $0 \leq t \leq i$ and $\lambda \in \mathcal{P}_i(g-i)$. It follows from (76) that

$$(77) \quad \partial_i^J(x^{(t)} \otimes s_\lambda(x)) = \left(\sum_{j=1}^i (-1)^{j-1} x^{(t)} \otimes s_{\lambda \hat{j}}(x) \otimes x^{\lambda_j + i - j} \right) + \left(\sum_{j=1}^i (-1)^{j-1} x^{(t-1)} \otimes s_{\lambda \hat{j}}(x) \otimes x^{\lambda_j + i - j + 1} \right),$$

where the second term vanishes identically when $t = 0$, while the first one disappears when $t = i$. Using (43) and the fact that $\partial_i^K = k_i^g$ as defined in (46) we get

$$(78) \quad (\partial_i^K \circ \bar{p}_i)(x^{(t)} \otimes s_\lambda(x)) = \sum_{j=1}^i (-1)^{j-1} \cdot \left(\sum_{I \in \binom{[i]}{t}} s_{(\lambda + (1^I)) \hat{j}}(x) \otimes x^{(\lambda + (1^I))_j + i - j} \right).$$

Comparing (77) with (78), it follows that in order to prove the commutativity of the diagram it suffices to check that for fixed $j \in \{1, \dots, i\}$ we have

$$(79) \quad \begin{aligned} \bar{p}_{i-1}(x^{(t)} \otimes s_{\lambda \hat{j}}(x)) &= \sum_{j \notin I \in \binom{[i]}{t}} s_{(\lambda + (1^I)) \hat{j}}(x) \text{ and} \\ \bar{p}_{i-1}(x^{(t-1)} \otimes s_{\lambda \hat{j}}(x)) &= \sum_{j \in I \in \binom{[i]}{t}} s_{(\lambda + (1^I)) \hat{j}}(x). \end{aligned}$$

Note that $\bar{p}_{i-1}(x^{(t)} \otimes s_{\lambda^{\hat{j}}}(x)) = \sum_{I' \in \binom{[i-1]}{t}} s_{\lambda^{\hat{j}} + (1^{I'})}(x)$ and that the correspondence

$$(80) \quad I' \longrightarrow I = \{k : k \in I' \text{ and } k < j\} \cup \{k+1 : k \in I' \text{ and } k \geq j\}$$

establishes a bijection between the collection of sets $I' \in \binom{[i-1]}{t}$ and that of sets $I \in \binom{[i]}{t}$ with $j \notin I$, which has the property that $(\lambda + (1^I))^{\hat{j}} = \lambda^{\hat{j}} + (1^{I'})$. This proves the first equality in (79). For the second equality we use

$$\bar{p}_{i-1}(x^{(t-1)} \otimes s_{\lambda^{\hat{j}}}(x)) = \sum_{I' \in \binom{[i-1]}{t-1}} s_{\lambda^{\hat{j}} + (1^{I'})}(x)$$

and note the bijective correspondence between sets $I' \in \binom{[i-1]}{t}$ and $I \in \binom{[i]}{t}$ with $j \in I$, given by

$$(81) \quad I' \longrightarrow I = \{k : k \in I' \text{ and } k < j\} \cup \{j\} \cup \{k+1 : k \in I' \text{ and } k \geq j\}.$$

Moreover, this correspondence has the property that $(\lambda + (1^I))^{\hat{j}} = \lambda^{\hat{j}} + (1^{I'})$, which shows the second equality in (79) and concludes our proof. \square

Corollary 5.10. *Let $p_{\bullet} : J_{\bullet} \longrightarrow K_{\bullet}$ be the map of complexes constructed in Proposition 5.9, and let C_{\bullet} denote the minimal resolution of the module C in (75). We have*

$$(82) \quad C_i = \text{Ker}(p_{i+1}), \text{ for } i = 0, \dots, g-2.$$

Proof. Since $H_0(p_{\bullet}) : H_0(J_{\bullet}) \longrightarrow H_0(K_{\bullet})$ is an isomorphism, it follows that by applying the mapping cone construction to the map of complexes $p_{\bullet} : J_{\bullet} \longrightarrow K_{\bullet}$ and minimizing the resulting complex as in Lemma 4.2, we obtain a minimal complex M_{\bullet} whose terms are

$$M_i = \text{Ker}(p_{i-1}) \oplus \text{Coker}(p_i), \text{ for } i = 0, \dots, g+1,$$

and whose only non-zero homology is $H_2(M_{\bullet}) = H_1(J_{\bullet}) = C$. Since the map p_0 is an isomorphism and p_1 is surjective, we get that $M_0 = M_1 = 0$ so $M_{\bullet+2}$ is in fact a minimal resolution of C , from which we deduce that $M_{i+2} \cong C_i$, for $i = 0, \dots, g-1$.

We have that C_i is a free module with generators of degree $i+1$, so the same must be true about M_{i+2} . Since $\text{Ker}(p_{i+1})$ is a free module generated in degree $i+1$ and $\text{Coker}(p_{i+2})$ is a free module generated in degree $i+2$, it follows that $\text{Coker}(p_{i+2}) = 0$ for all $i = 0, \dots, g-2$ and therefore p_i is surjective for all $i = 0, \dots, g$. It follows that $C_i = M_{i+2} = \text{Ker}(p_{i+1})$, for $i = 0, \dots, g-2$. \square

Remark 5.11. Keeping the notation above, we have an isomorphism $C_{g-1} \cong M_{g+1}$ of free modules of rank one generated in degree $g+1$, and we conclude that $\text{Ker}(p_g) = 0$ and $M_{g+1} = \text{Coker}(p_{g+1}) = K_{g+1}$.

5.7. The first linear strand of the resolution of R . The goal of this section is to complete the proof of Theorem 5.4. Our strategy is summarized in the following steps:

(1) We construct a map of complexes $\tilde{q}_{\bullet} : F_{\bullet} \longrightarrow C_{\bullet}$ such that the induced map

$$H_0(\tilde{q}_{\bullet}) : H_0(F_{\bullet}) \longrightarrow H_0(C_{\bullet})$$

is a surjection $\tilde{R} \longrightarrow C$.

(2) If we let $R' := \text{Ker}(\tilde{R} \rightarrow C)$, then $R \subseteq R'$ and the mapping cone of \tilde{q}_{\bullet} gives a resolution of R' . Minimizing it, we get that the terms in the first linear strand of the minimal resolution of R' can be described as kernels of the Koszul maps (57).

(3) The quotient R'/R is a module generated in degree at least 2, so the first strands of the minimal resolutions of R and R' coincide.

Proposition 5.12. For $i = 0, \dots, g-2$, we consider the \mathfrak{sl}_2 -equivariant map

$$\bar{q}_i : D^{2i} U \otimes \bigwedge^{i+1} \text{Sym}^{g-2} U \longrightarrow D^{i+1} U \otimes \bigwedge^{i+1} \text{Sym}^{g-1} U$$

induced by the map $\Delta_1 : D^{2i} U \longrightarrow D^{i+1} U \otimes D^{i+1} U$ in Lemma 3.1, followed by

$$\nu_{g-2-i}^{i+1} : D^{i+1} U \otimes \bigwedge^{i+1} \text{Sym}^{g-2} U \xrightarrow{(43)} \bigwedge^{i+1} \text{Sym}^{g-1} U.$$

The maps \bar{q}_i define a map of complexes $q_\bullet : F_\bullet \longrightarrow J_{\bullet+1}$ (if we take q_0 to send the summand $S \subseteq F_0$ to zero), such that the induced map $H_0(F_\bullet) \longrightarrow H_1(J_\bullet)$ is a surjection $\tilde{R} \longrightarrow C$ containing R in the kernel.

Proof. The map $\Delta_1 : D^0 U \longrightarrow D^1 U \otimes D^1 U$ can be identified with the \mathfrak{sl}_2 -equivariant inclusion of $\mathfrak{k} \cong \bigwedge^2 U \longrightarrow U \otimes U$, so the map $q_0 : F_0 \longrightarrow J_1$ is given by

$$S \oplus \text{Sym}^{g-2} U \otimes S(-1) \xrightarrow{q_0} (U \otimes \text{Sym}^{g-1} U) \otimes S(-1),$$

where S is sent to 0, and $\text{Sym}^{g-2} U \cong \bigwedge^2 U \otimes \text{Sym}^{g-2} U \longrightarrow U \otimes \text{Sym}^{g-1} U$ is an inclusion given by the Koszul differential. Since $\partial_1^J : J_1 \longrightarrow J_0$ is induced by the multiplication map $U \otimes \text{Sym}^{g-1} U \longrightarrow \text{Sym}^g U$, it follows that $\partial_1^J \circ q_0 = 0$. Assuming $q_\bullet : F_\bullet \longrightarrow J_{\bullet+1}$ is indeed a map of complexes (which we shall prove shortly), to see that $H_0(q_\bullet)$ induces a surjection $\tilde{R} \rightarrow C$ containing R in its kernel, it suffices to note that q_0 sends the generators $\text{Sym}^{g-2} U \subseteq \tilde{R}$ isomorphically onto the generators $\text{Sym}^{g-2} U$ of C , and that the generator $1 \in R$ is sent to 0 since $q_0(S) = 0$.

To prove that $q_\bullet : F_\bullet \longrightarrow J_{\bullet+1}$ is a map of complexes we have to verify that for every $i \geq 1$ we get a commutative diagram (note that since $q_0(S) = 0$, we can ignore the quadratic part of the differential ∂_1^F)

$$\begin{array}{ccc} D^{2i} U \otimes \bigwedge^{i+1} \text{Sym}^{g-2} U & \xrightarrow{\bar{q}_i} & D^{i+1} U \otimes \bigwedge^{i+1} \text{Sym}^{g-1} U \\ \partial_i^F \downarrow & & \downarrow \partial_{i+1}^J \\ (D^{2i-2} U \otimes \bigwedge^i \text{Sym}^{g-2} U) \otimes \text{Sym}^g U & \xrightarrow{\bar{q}_{i-1} \otimes \text{id}} & (D^i U \otimes \bigwedge^i \text{Sym}^{g-1} U) \otimes \text{Sym}^g U \end{array}$$

The space $D^{2i} U \otimes \bigwedge^{i+1} \text{Sym}^{g-2} U$ has a basis consisting of elements $x^{(t)} \otimes s_\lambda(x)$ where $0 \leq t \leq 2i$ and $\lambda \in \mathcal{P}_{i+1}(g-i-2)$. Since

$$\Delta_1(x^{(t)}) = \sum_{\substack{t'+t''=t+1 \\ t', t'' \leq i+1}} (t' - t'') \cdot x^{(t')} \otimes x^{(t'')}$$

it follows that

$$\bar{q}_i(x^{(t)} \otimes s_\lambda(x)) = \sum_{\substack{t'+t''=t+1 \\ t', t'' \leq i+1}} (t' - t'') \cdot x^{(t')} \otimes \left(\sum_{I \in \binom{[i+1]}{t''}} s_{\lambda+(1^I)}(x) \right)$$

and using (76) we get that $(\partial_{i+1}^J \circ \bar{q}_i)(x^{(t)} \otimes s_\lambda(x))$ is computed by

$$\sum_{\substack{t'+t''=t+1 \\ t', t'' \leq i+1}} \sum_{I \in \binom{[i+1]}{t''}} \sum_{\substack{u'+s'=t' \\ u' \leq 1, s' \leq i}} \sum_{j=1}^{i+1} (-1)^{j-1} \cdot (t' - t'') \cdot x^{(s')} \otimes s_{(\lambda+(1^I))_j}(x) \otimes x^{(\lambda+(1^I))_j+i+1-j+u'}$$

$$(83) = \sum_{\substack{u'+s'+t''=t+1 \\ u' \leq 1, s' \leq i, t'' \leq i+1}} \sum_{I \in \binom{[i+1]}{u'}} \sum_{j=1}^{i+1} (-1)^{j-1} \cdot (u' + s' - t'') \cdot x^{(s')} \otimes s_{(\lambda+(1^I))_j}(x) \otimes x^{(\lambda+(1^I))_j+i+1-j+u'}.$$

On the other hand we have using (71) that

$$\partial_i^F(x^{(t)} \otimes s_\lambda(x)) = \sum_{\substack{u+s=t \\ u \leq 2, s \leq 2i-2}} \sum_{j=1}^{i+1} (-1)^{j-1} \cdot x^{(s)} \otimes s_{\lambda_j}(x) \otimes (x^{(u)} \cdot x^{\lambda_j+i+1-j}),$$

where $x^{(u)} = x^u$ when $u = 0, 2$, and $x^{(1)} = 2x$. Thus $((\bar{q}_{i-1} \otimes \text{id}_{\text{Sym}^g U}) \circ \partial_i^F)(x^{(t)} \otimes s_\lambda(x))$ is computed by

$$(84) = \sum_{\substack{u+s=t \\ u \leq 2, s \leq 2i-2}} \sum_{j=1}^{i+1} \sum_{\substack{s'+s''=s+1 \\ s', s'' \leq i}} \sum_{I' \in \binom{[i]}{s''}} (-1)^{j-1} \cdot (s' - s'') \cdot x^{(s')} \otimes s_{\lambda^{\hat{j}}+(1^{I'})}(x) \otimes (x^{(u)} \cdot x^{\lambda_j+i+1-j})$$

$$= \sum_{\substack{u+s'+s''=t+1 \\ u \leq 2, s', s'' \leq i}} \sum_{j=1}^{i+1} \sum_{I' \in \binom{[i]}{s''}} (-1)^{j-1} \cdot (s' - s'') \cdot x^{(s')} \otimes s_{\lambda^{\hat{j}}+(1^{I'})}(x) \otimes (x^{(u)} \cdot x^{\lambda_j+i+1-j})$$

To conclude the proof we show that we can identify (83) with (84). We fix j and s' with $1 \leq j \leq i+1$ and $0 \leq s' \leq i$. It is then enough to verify that

$$(85) \quad \sum_{\substack{u'+t''=t+1-s' \\ u' \leq 1, t'' \leq i+1}} \sum_{I \in \binom{[i+1]}{u'}} (u' + s' - t'') \cdot s_{(\lambda+(1^I))_j}(x) \otimes x^{(\lambda+(1^I))_j+i+1-j+u'}$$

$$= \sum_{\substack{u+s''=t+1-s' \\ u \leq 2, s'' \leq i}} \sum_{I' \in \binom{[i]}{s''}} (s' - s'') \cdot s_{\lambda^{\hat{j}}+(1^{I'})}(x) \otimes (x^{(u)} \cdot x^{\lambda_j+i+1-j}).$$

We fix u, s'' with $u + s'' = t + 1 - s'$, $u \leq 2$, $s'' \leq i$, and fix $I' \in \binom{[i]}{s''}$. We consider three separate cases:

$u = 0$: We let $u' = 0$, $t'' = s''$, and define I via (80). The term in (85) corresponding to this choice of parameters is equal to $(s' - s'') \cdot s_{\lambda^{\hat{j}}+(1^{I'})}(x) \otimes (x^{(u)} \cdot x^{\lambda_j+i+1-j})$ since $u' - t'' = -s''$, $(\lambda + (1^I))_j^{\hat{j}} = \lambda^{\hat{j}} + (1^{I'})$, $x^{(u)} = 1$ and $(\lambda + (1^I))_j + i + 1 - j + u' = \lambda_j + i + 1 - j$, where the last equality follows from the fact that $j \notin I$ and $u' = 0$.

$u = 2$: We let $u' = 1$, $t'' = s'' + 1$, and define I via (81). The term in (85) corresponding to this choice of parameters is equal to $(s' - s'') \cdot s_{\lambda^{\hat{j}}+(1^{I'})}(x) \otimes (x^{(u)} \cdot x^{\lambda_j+i+1-j})$ since $u' - t'' = -s''$, $(\lambda + (1^I))_j^{\hat{j}} = \lambda^{\hat{j}} + (1^{I'})$, $x^{(u)} = x^2$ and $(\lambda + (1^I))_j + i + 1 - j + u' = 2 + \lambda_j + i + 1 - j$, where the last equality follows from the fact that $j \in I$ and $u' = 1$.

$u = 1$: In this case there are two terms in (85) whose sum is

$$(s' - s'') \cdot s_{\lambda^{\hat{j}}+(1^{I'})}(x) \otimes (x^{(u)} \cdot x^{\lambda_j+i+1-j}) = 2 \cdot (s' - s'') \cdot s_{\lambda^{\hat{j}}+(1^{I'})}(x) \otimes x^{\lambda_j+i+2-j}$$

and they correspond to the following choice of parameters. If we let $u' = 0$, $t'' = s'' + 1$, and define I via (81) then we get

$$(u' + s' - t'') \cdot s_{(\lambda+(1^I))_j}(x) \otimes x^{(\lambda+(1^I))_j+i+1-j+u'} = (s' - s'' - 1) \cdot s_{\lambda^{\hat{j}}+(1^{I'})}(x) \otimes x^{\lambda_j+i+2-j},$$

while if we take $u' = 1, t'' = s''$, and define I via (80) then we get

$$(u' + s' - t'') \cdot s_{(\lambda+(1^I))j}(x) \otimes x^{(\lambda+(1^I))j+i+1-j+u'} = (1 + s' - s'') \cdot s_{\lambda j+(1^{I'})}(x) \otimes x^{\lambda j+i+2-j}.$$

To finish the proof, we make sure that all the terms in (85) are accounted for in the above case analysis:

- If $u' = 0$ and $j \in I$ then take $u = 1, s'' = t'' - 1$ and define I' by reversing (81).
- If $u' = 0$ and $j \notin I$ then take $u = 0, s'' = t''$ and define I' by reversing (80).
- If $u' = 1$ and $j \in I$ then take $u = 2, s'' = t'' - 1$ and define I' by reversing (81).
- If $u' = 1$ and $j \notin I$ then take $u = 1, s'' = t''$ and define I' by reversing (80). □

The map q_\bullet constructed in Proposition 5.12 induces a map of complexes $F_\bullet \rightarrow C_\bullet$ as follows

Corollary 5.13. *For $i = 0, \dots, g-2$ we have $p_{i+1} \circ q_i = 0$, therefore $q_i(F_i) \subseteq C_i = \text{Ker}(p_{i+1})$, by applying Corollary 5.10. The induced map of complexes $\tilde{q}_\bullet : F_\bullet \rightarrow C_\bullet$ has the property that*

$$H_0(\tilde{q}_\bullet) : H_0(F_\bullet) = \tilde{R} \rightarrow H_0(C_\bullet) = C \text{ is a surjective map.}$$

Proof. If we replace i by $i+1$, let $d = g-2-i$, and tensor the commutative diagram (45) with $D^{i+1}U$ then we obtain a commutative diagram

$$\begin{array}{ccc} (D^{i+1}U)^{\otimes 2} \otimes \text{Sym}^{g-2-i}(D^{i+1}U) & \xrightarrow{\text{id}^{\otimes 2} \otimes \psi_{g-2-i}} & (D^{i+1}U)^{\otimes 2} \otimes \wedge^{i+1}(\text{Sym}^{g-2}U) \\ \downarrow & & \downarrow \text{id} \otimes \nu_{g-2-i}^{i+1} \\ D^{i+1}U \otimes \text{Sym}^{g-1-i}(D^{i+1}U) & \xrightarrow{\text{id} \otimes \psi_{g-1-i}} & D^{i+1}U \otimes \wedge^{i+1}(\text{Sym}^{g-1}U) \end{array}$$

Restricting the top row of the diagram above along the chain of maps

$$D^{2i}U \xrightarrow{\Delta_1} \bigwedge^2 D^{i+1}U \subseteq (D^{i+1}U)^{\otimes 2}$$

and recalling that q_i was defined by restricting along the same maps, we obtain the commutativity of the top square in the diagram

$$(86) \quad \begin{array}{ccc} D^{2i}U \otimes \text{Sym}^{g-2-i}(D^{i+1}U) & \xrightarrow{\text{id} \otimes \psi_{g-2-i}} & D^{2i}U \otimes \wedge^{i+1}(\text{Sym}^{g-2}U) \\ \delta_2 \downarrow & & \downarrow \bar{q}_i \\ D^{i+1}U \otimes \text{Sym}^{g-1-i}(D^{i+1}U) & \xrightarrow{\text{id} \otimes \psi_{g-1-i}} & D^{i+1}U \otimes \wedge^{i+1}(\text{Sym}^{g-1}U) \\ \delta_1 \downarrow & & \downarrow \bar{p}_{i+1} \\ \text{Sym}^{g-i}(D^{i+1}U) & \xrightarrow{\psi_{g-i}} & \wedge^{i+1}(\text{Sym}^g U) \end{array}$$

Note that the bottom square of this diagram also commutes, being another instance of (45), namely that where we replace i by $i+1$, let $d = g-1-i$, and recall that $\bar{p}_{i+1} = \nu_{g-i-1}^{i+1}$.

In the diagram (86) the maps in the left column are induced by Koszul differentials, so their composition must be zero. Since the horizontal maps are isomorphisms the maps in the right column must compose to zero as well, hence $\bar{p}_{i+1} \circ \bar{q}_i = 0$ and therefore $p_{i+1} \circ q_i = 0$. Using the fact that $C_i = \text{Ker}(p_{i+1})$ (see (82)), it follows that q_\bullet induces (by restricting the range) a map of complexes $\tilde{q}_\bullet : F_\bullet \rightarrow C_\bullet$. Just as in the proof of Proposition 5.12 it follows by inspecting the generators that $H_0(\tilde{q}_\bullet)$ is surjective. □

Proof of Theorem 5.4. We consider the minimization N_\bullet of the mapping cone of \tilde{q}_\bullet , whose terms are computed by Lemma 4.2 via

$$(87) \quad N_i = \text{Ker}(\tilde{q}_{i-1}) \oplus \text{Coker}(\tilde{q}_i).$$

We have that $N_0 = 0$ and $N_1 = S \oplus \text{Coker}(\tilde{q}_1)$, where $\text{Coker}(\tilde{q}_1)$ is a free module generated in degree 2 (if non-zero). From the long exact sequence for the homology of the mapping cone, we obtain that N_\bullet has only one non-zero homology group, namely $H_1(N_\bullet) = \text{Ker}\{H_0(\tilde{q}_\bullet) : H_0(F_\bullet) \rightarrow H_0(C_\bullet)\} = R' \supseteq R$. It follows that N_\bullet is quasi-isomorphic to the minimal free resolution of R' , shifted by 1. Since the summand $S \subseteq N_1$ maps onto R , the quotient R'/R is generated in degree 2 (if non-zero), in particular

$$(88) \quad \text{Tor}_i^S(R, \mathbf{k})_{i+1} = \text{Tor}_i^S(R', \mathbf{k})_{i+1} \text{ for all } i \geq 0.$$

Since $\text{Ker}(\tilde{q}_i)$ is a free module generated in degree $i+1$ for $i \geq 1$, and $\text{Coker}(\tilde{q}_i)$ is a free module generated in degree $i+1$ for $1 \leq i \leq g-2$, and in degree $g+1$ when $i = g-1$, we conclude that

$$(89) \quad \text{Tor}_i^S(R', \mathbf{k})_{i+1} = \text{Tor}_0^S(N_{i+1}, \mathbf{k})_{i+1} = \text{Ker}(\tilde{q}_i)_{i+1}, \text{ for all } i \geq 0.$$

Recall from Proposition 5.9 and (82) that C_i is a direct summand of J_{i+1} . From Proposition 5.12 and Corollary 5.13 we recall that the map of complexes $\tilde{q}_\bullet : F_\bullet \rightarrow C_\bullet$ is induced by a map of complexes $q_\bullet : F_\bullet \rightarrow J_{\bullet+1}$. It follows that $\text{Ker}(\tilde{q}_i) = \text{Ker}(q_i)$ and in particular

$$(90) \quad \text{Ker}(\tilde{q}_i)_{i+1} = \text{Ker}\left\{ D^{2i} U \otimes \bigwedge^{i+1} (\text{Sym}^{g-2} U) \xrightarrow{\bar{q}_i} D^{i+1} U \otimes \bigwedge^{i+1} (\text{Sym}^{g-1} U) \right\},$$

where the map \bar{q}_i was defined in Proposition 5.12. Using the commutative diagram

$$\begin{array}{ccc} D^{2i} U \otimes \text{Sym}^{g-2-i}(D^{i+1} U) & \xrightarrow{\text{id} \otimes \psi_{g-2-i}} & D^{2i} U \otimes \bigwedge^{i+1} (\text{Sym}^{g-2} U) \\ \delta_2 \downarrow & & \downarrow \bar{q}_i \\ D^{i+1} U \otimes \text{Sym}^{g-1-i}(D^{i+1} U) & \xrightarrow{\text{id} \otimes \psi_{g-1-i}} & D^{i+1} U \otimes \bigwedge^{i+1} (\text{Sym}^{g-1} U) \end{array}$$

from the proof of Corollary 5.13, together with (88), (89) and (90) we obtain

$$K_{i,1}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) = \text{Tor}_i^S(R, \mathbf{k})_{i+1} = \text{Tor}_i^S(R', \mathbf{k})_{i+1} = \text{Ker}(\bar{q}_i) = \text{Ker}(\delta_2),$$

concluding the proof of the theorem. \square

5.8. The minimal resolution of R . Throughout this section we assume that $\text{char}(\mathbf{k}) \neq 2$, and we analyze more carefully the constructions from the previous sections in order to prove Theorems 5.1 and 5.3. The main observation is the following.

Lemma 5.14. *For $i = 1, \dots, g-2$, we have that $\text{Coker}(\tilde{q}_i)$ is isomorphic to $W_{g-2-i}^{(i+1)} \otimes_{\mathbf{k}} S(-i-1)$.*

Proof. Since F_i and C_i are free S -modules generated in degree $i+1$ for $1 \leq i \leq g-2$, and since \tilde{q}_i is a homogeneous map, it follows that $\text{Coker}(\tilde{q}_i)$ is a free module generated in degree $i+1$. To identify the minimal generators of $\text{Coker}(\tilde{q}_i)$, we recall that \tilde{q}_i is induced by the map q_i , whose image lands inside $C_i = \text{Ker}(p_{i+1})$ by Corollary 5.13. It follows that the minimal generators of $\text{Coker}(\tilde{q}_i)$ are described as

$$\frac{\text{Ker}(\bar{p}_{i+1})}{\text{Im}(\bar{q}_i)} \stackrel{(86)}{\cong} \frac{\text{Ker}(\delta_1)}{\text{Im}(\delta_2)} = W_{g-2-i}^{(i+1)}. \quad \square$$

Using this observation, we can now deduce Theorems 5.1 and 5.3. We start with:

Corollary 5.15. *We have $R' = R$ and Theorem 5.3 holds.*

Proof. The complex $N_{\bullet+1}$ gives the minimal free resolution of R' . Since $R \subseteq R'$ is the image of the summand S of N_1 in (87), in order to prove that $R' = R$, it suffices to check that $\text{Coker}(\tilde{q}_1) = 0$, which by Lemma 5.14 is equivalent to the vanishing $W_{g-3}^{(2)} = 0$. It is then enough to prove that $W^{(2)}$ is identically zero, which is equivalent to the fact that the inclusion $\Delta_1 : D^2 U \hookrightarrow \bigwedge^2(D^2 U)$ is an equality. Since $\dim(D^2 U) = 3$, the source and target of Δ_1 have both dimension three, hence Δ_1 is indeed an equality.

Since the complex $N_{\bullet+1}$ gives the minimal free resolution of $R' = R$, we obtain for $i = 1, \dots, g-3$

$$K_{i,2}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) = \text{Tor}_i^S(R, \mathbf{k})_{i+2} = (N_{i+1} \otimes_S \mathbf{k})_{i+2} \stackrel{(87)}{=} \text{Coker}(\tilde{q}_{i+1}) \otimes_S \mathbf{k} = W_{g-3-i}^{(i+2)}. \quad \square$$

We end this section by verifying Theorem 5.1.

Proof of Theorem 5.1. We first prove that R is Cohen–Macaulay (see also [24, Proposition 6.1] for a different proof in characteristic zero). Since $\dim(R) = \dim(\mathcal{T}) + 1 = 3$, and $\dim(S) = g + 1$, it suffices to check that R has projective dimension (at most) $g - 2$ as an S -module. Since $N_{\bullet+1}$ is the minimal resolution of R , this is equivalent to the fact that $N_i = 0$ for $i \geq g$. We obtain this conclusion using (87), since the source of \tilde{q}_{i-1} is $F_{i-1} = 0$ for $i \geq g$, and the target of \tilde{q}_i is $C_i = 0$ for $i \geq g$.

To prove that R is Gorenstein of regularity 3, it is then enough to check that

$$\text{Tor}_{g-2}^S(R, \mathbf{k}) = N_{g-1} \otimes_S \mathbf{k}$$

is one-dimensional, concentrated in degree $g + 1$. Since $\text{Coker}(\tilde{q}_{g-1}) = C_{g-1} \cong S(-g - 1)$, it follows from (87) that it is enough to check that \tilde{q}_{g-2} is injective, or equivalently, that \bar{q}_{g-2} is injective. Using the commutativity of the top square in (86) with $i = g - 2$, this is further equivalent to the injectivity of

$$D^{2(g-2)} U \xrightarrow{\delta_2} D^{g-1} U \otimes D^{g-1} U.$$

Since δ_2 is given in this case by the map Δ_1 in Lemma 3.1 with $a = g - 1$, the conclusion follows. \square

We record one more consequence of the previous results:

Corollary 5.16. *If $\text{char}(\mathbf{k}) \neq 2$, the dualizing sheaf $\omega_{\mathcal{T}}$ is trivial, and $H^1(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(d)) = 0$ for all d .*

Proof. It follows from Theorem 5.1 that $\text{Ext}_S^{g-2}(R, S) = R(g + 1)$. By sheafification this implies that $\mathcal{E}xt_{\mathbf{P}^g}^{g-2}(\mathcal{O}_{\mathcal{T}}, \mathcal{O}_{\mathbf{P}^g}) = \mathcal{O}_{\mathcal{T}}(g + 1)$. Using $\omega_{\mathbf{P}^g} \cong \mathcal{O}_{\mathbf{P}^g}(-g - 1)$ and [16, Proposition III.7.5], it follows that

$$\omega_{\mathcal{T}} = \mathcal{E}xt_{\mathbf{P}^g}^{g-2}(\mathcal{O}_{\mathcal{T}}, \omega_{\mathbf{P}^g}) = \mathcal{E}xt_{\mathbf{P}^g}^{g-2}(\mathcal{O}_{\mathcal{T}}, \mathcal{O}_{\mathbf{P}^g})(-g - 1) = \mathcal{O}_{\mathcal{T}},$$

as desired. The vanishing $H^1(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(d)) = 0$ follows from the fact that \mathcal{T} is arithmetically Cohen–Macaulay (combine for instance [8, Propositions A1.11 and A1.16]). \square

One consequence of the Cohen–Macaulay property of R is the isomorphism $R \cong \Gamma_{\mathcal{T}}(\mathcal{O}_{\mathcal{T}}(1))$, which amounts to $\Gamma_{\mathcal{T}}(\mathcal{O}_{\mathcal{T}}(1))$ being generated in degree one.

5.9. The resolution of R in characteristic 2. The goal of this section is to prove that (unlike in the general case) the resolution of R consists of a single linear strand when $\text{char}(\mathbf{k}) = 2$, which is obtained as a subcomplex of F_{\bullet} as explained below. We identify $S = \mathbf{k}[z_0, \dots, z_g]$, where z_i is dual to $x^{(i)}$, and claim that the 2×2 minors of the matrix

$$M = \begin{bmatrix} z_0 & z_1 & \cdots & z_{g-2} \\ z_2 & z_3 & \cdots & z_g \end{bmatrix}$$

vanish on \mathcal{T} . Since \mathcal{T} is irreducible, it is enough to check the vanishing on some affine chart. Choosing the chart as in (70) and evaluating M on an element $f \in \mathcal{T}$ we obtain

$$M(f) = u \cdot \begin{bmatrix} 1 & t & t^2 & t^3 & \cdots & t^{g-2} \\ t^2 & t^3 & t^4 & t^5 & \cdots & t^g \end{bmatrix} + v \cdot \begin{bmatrix} 0 & t & 0 & t^3 & \cdots & (g-2) \cdot t^{g-2} \\ 0 & t^3 & 0 & t^5 & \cdots & g \cdot t^g \end{bmatrix}$$

for some scalars $u, v, t \in \mathbf{k}$. Since the second row of $M(f)$ is obtained from the first by multiplying by t^2 , the 2×2 minors of $M(f)$ vanish on \mathcal{T} , as desired. The 2×2 minors of M define a rational normal scroll of dimension 2 and degree $g - 1$, containing \mathcal{T} , see [8, Section A2H]. Carrying out in characteristic 2 the local analysis in Section 5.4, we obtain that $\deg(\mathcal{T}) = g - 1$, hence \mathcal{T} equals the scroll. We conclude that the minimal resolution of R is given by an Eagon–Northcott complex.

To identify this as a subcomplex of F_\bullet , we let for $i \geq 1$

$$(91) \quad D_{\text{odd}}^{2i} U = \bigoplus_{1 \leq t \leq i} \mathbf{k} \cdot x^{(2t-1)} = \{f \in D^{2i} : \mathfrak{L} \cdot f = \mathfrak{R} \cdot f = 0\} \subseteq D^{2i} U.$$

We define $F_0^{\text{odd}} = S$ and

$$(92) \quad F_i^{\text{odd}} = \left(D_{\text{odd}}^{2i} U \otimes \bigwedge^{i+1} \text{Sym}^{g-2} U \right) \otimes S(-i-1), \text{ for } i = 1, \dots, g-2.$$

Since $\binom{2}{1} = 0$ in \mathbf{k} , it follows from (71) that F_\bullet^{odd} determines a subcomplex of F_\bullet . Taking into account that $\text{rk}(F_1^{\text{odd}}) = \binom{g-1}{2} = \dim(I_2)$, we get that F_1^{odd} surjects onto the quadratic part of the ideal I . Since I is generated in degree 2, we get that $H_0(F_\bullet^{\text{odd}}) = R$. We can now define D_{even}^{2i} and F_\bullet^{even} in analogy with (91) and (92), and observe that F_\bullet^{even} is also a subcomplex of F_\bullet : it follows from (71) that the linear part of the differentials ∂_\bullet^F send F_\bullet^{even} to itself, so we only need to worry about the quadratic part of ∂_1^F ; since F_1^{odd} already gives the minimal generators of I , the quadratic part of ∂_1^F restricts to 0 on F_1^{even} , showing that F_\bullet^{even} is indeed a subcomplex. It follows that $F_\bullet = F_\bullet^{\text{odd}} \oplus F_\bullet^{\text{even}}$, and in particular $H_i(F_\bullet^{\text{odd}}) = 0$ for $i > 0$, showing that F_\bullet^{odd} is the minimal resolution of R .

Remark 5.17. A similar local calculation shows that if $3 \leq \text{char}(\mathbf{k}) = p \leq g$, then \mathcal{T} is contained in a rational normal scroll of codimension $g - p$, defined by the vanishing of the 2×2 minors of the matrix

$$\begin{bmatrix} z_0 & z_1 & \cdots & z_{g-p} \\ z_p & z_{p+1} & \cdots & z_g \end{bmatrix}.$$

The (linear) syzygies of the scroll are then going to be contained inside the linear syzygies of \mathcal{T} . Since the Eagon–Northcott resolution of the scroll has length $g - p$, we conclude that $K_{i,1}(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(1)) \neq 0$ for $1 \leq i \leq g - p$. In the range $3 \leq p \leq \frac{g+1}{2}$ this non-vanishing is sharp: indeed, we have $b_{i,1} = b_{g-2-i,2}$, so the conclusion $b_{i,1} \neq 0$ if and only if $1 \leq i \leq g - p$ follows from (56).

6. MODULI OF PSEUDO-STABLE CURVES AND GREEN'S CONJECTURE

In this section we explain how Theorem 1.1 implies Green's Conjecture for general curves of genus g , that is, Theorem 1.2. The key idea is to consider a moduli space of curves of genus g which contains among its points the cuspidal curves that are hyperplane sections of the tangential variety $\mathcal{T} \subseteq \mathbb{P}^g$. We fix throughout an algebraically closed field \mathbf{k} of characteristic $p \neq 2, 3$. This assumption excludes the cases $p = 3$ and $g = 3, 4$ from Theorem 1.2, but Green's Conjecture can be verified directly in these cases (see also [25] for a proof of Green's Conjecture in small genera).

Definition 6.1. A *pseudo-stable* curve C of genus g is a connected curve over \mathbf{k} of arithmetic genus g with only nodes and cusps as singularities such that (i) its dualizing sheaf ω_C is ample and (ii) each irreducible component of arithmetic genus 1 of C intersects the rest of the curve in at least two points.

Schubert [26] constructs the coarse moduli space of pseudo-stable curves of genus g as the Chow quotient of 3-canonically embedded curves. The construction and geometry of $\overline{\mathcal{M}}_g^{\text{ps}}$ is described in [26] and [18]. In particular, one has a divisorial contraction

$$\pi : \overline{\mathcal{M}}_g \rightarrow \overline{\mathcal{M}}_g^{\text{ps}},$$

which at the level of geometric points replaces each elliptic tail of a stable curve with a cusp. More precisely, if $[C] \in \overline{\mathcal{M}}_g$ is a stable curve having elliptic tails E_1, \dots, E_ℓ with $E_i \cap (\overline{C \setminus E_i}) = \{p_i\}$, for $i = 1, \dots, \ell$, then $[C'] := \pi([C])$ is described as follows: if we denote by D the union of the components of C different from the elliptic tails then the pseudo-stable curve C' is birational to D , and there exists a morphism $\nu : C \rightarrow C'$ contracting each tail E_i to a cusp $q_i \in C'$ (see [18, Proposition 3.1] for details).

The following result is hinted at in [27].

Proposition 6.2. *Pseudo-stable curves of genus $g \geq 3$ form a Deligne-Mumford stack $\overline{\mathcal{M}}_g^{\text{ps}}$ over $\text{Spec } \mathbb{Z}[\frac{1}{6}]$.*

Proof. To conclude that the algebraic stack of pseudo-stable curves of genus g is of Deligne-Mumford type, it suffices to show that a pseudo-stable curve C of genus $g \geq 3$ admits no infinitesimal deformations, that is, $H^0(C, \Omega_C^\vee) = 0$. We denote by $p_1, \dots, p_\ell \in C$ the cusps of C , by $\nu : D \rightarrow C$ the normalization map at the cusps and set $q_i := \nu^{-1}(p_i)$, for $i = 1, \dots, \ell$. Using [27, Proposition 2.3], we have $\Omega_C^\vee = \nu_*(\Omega_D^\vee(-q_1 - \dots - q_\ell))$, in particular $H^0(C, \Omega_C^\vee) \cong H^0(D, \omega_D^\vee(-q_1 - \dots - q_\ell))$. Condition (ii) in Definition 6.1 implies that the ℓ -pointed curve $[D, q_1, \dots, q_\ell]$ is stable. Since infinitesimal deformations of $[D, p_1, \dots, p_\ell]$ are parametrized by $H^0(D, \Omega_D^\vee(-q_1 - \dots - q_\ell))$ this space vanishes and the conclusion follows. \square

Remark 6.3. In characteristic 2 and 3 the stack $\overline{\mathcal{M}}_g^{\text{ps}}$ is not Deligne-Mumford, for in these cases cuspidal curves may have infinitesimal automorphisms. Precisely, if C is a pseudo-stable with a cusp at the point q and $\nu : D \rightarrow C$ the normalization at q , then it is no longer the case that the inclusion $\Omega_{C,q}^\vee \subseteq (\nu_*\Omega_D)_q$ holds, that is, there exist regular vector fields on C that do not extend to a regular vector field on D , as explained in [27, Example 1].

Assume $g \geq 3$ and let \mathcal{M}_g^\sharp be the open substack of $\overline{\mathcal{M}}_g^{\text{ps}}$ classifying irreducible non-hyperelliptic pseudo-stable curves of genus g . If C is a non-hyperelliptic irreducible pseudo-stable curve, it follows from [17, Theorem 1.6] that ω_C is very ample. We show that the condition $K_{[\frac{g}{2}],1}(C, \omega_C) \neq 0$ can be described as the degeneracy locus of a morphism of vector bundles on \mathcal{M}_g^\sharp , and in particular that it is a closed condition. We will then use Theorem 1.1 to produce examples of curves in \mathcal{M}_g^\sharp for which $K_{[\frac{g}{2}],1}(C, \omega_C) = 0$, and conclude that this vanishing holds generically.

Let M_{ω_C} be the kernel bundle associated with (C, ω_C) , as defined in (49) and $\mathbb{P}_C := \mathbb{P}(H^0(C, \omega_C)^\vee) \cong \mathbb{P}^{g-1}$ and denote by M_C the kernel bundle associated with $(\mathbb{P}_C, \mathcal{O}_{\mathbb{P}_C}(1))$. We recall from (51) that

$$(93) \quad K_{i+1,1}(C, \omega_C) = \text{Ker} \left\{ H^0(\mathbb{P}_C, \bigwedge^i M_C(2)) \xrightarrow{\alpha_C} H^0(C, \bigwedge^i M_{\omega_C} \otimes \omega_C^2) \right\}$$

We construct vector bundles \mathcal{A} and \mathcal{B} on \mathcal{M}_g^\sharp , whose fibres at $[C] \in \mathcal{M}_g^\sharp$ are

$$(94) \quad \mathcal{A}([C]) = H^0(\mathbb{P}_C, \bigwedge^i M_C(2)) \quad \text{and} \quad \mathcal{B}([C]) = H^0(C, \bigwedge^i M_{\omega_C} \otimes \omega_C^2),$$

and a morphism $\alpha : \mathcal{A} \rightarrow \mathcal{B}$ whose restriction to the fiber over $[C]$ is the map α_C in (93). To construct \mathcal{B} , we start with the following.

Proposition 6.4. *If C is an irreducible pseudo-stable curve of genus g , then*

$$H^1\left(C, \bigwedge^i M_{\omega_C} \otimes \omega_C^2\right) = 0 \text{ for } i < g - 1.$$

Proof. We have that $\det(M_{\omega_C}) = \omega_C^\vee$, which implies that $\bigwedge^i M_{\omega_C} \otimes \omega_C^2 \cong \bigwedge^{g-1-i} M_{\omega_C}^\vee \otimes \omega_C$. Combining this with Serre duality, it follows that

$$H^1\left(C, \bigwedge^i M_{\omega_C} \otimes \omega_C^2\right) \cong H^0\left(C, \bigwedge^{g-1-i} M_{\omega_C}\right)^\vee.$$

Thinking of (51) as a two-step resolution of the kernel bundle, it follows that $\bigwedge^{g-1-i} M_{\omega_C}$ is resolved by the $(g-1-i)$ -th exterior power of this resolution, which is the Koszul-type complex

$$\bigwedge^{g-1-i} H^0(C, \omega_C) \otimes \mathcal{O}_C \rightarrow \bigwedge^{g-2-i} H^0(C, \omega_C) \otimes \omega_C \rightarrow \dots$$

Using the left exactness of the global sections functor, we get

$$H^0\left(C, \bigwedge^{g-1-i} M_{\omega_C}\right) = \text{Ker}\left\{\bigwedge^{g-1-i} H^0(C, \omega_C) \rightarrow \bigwedge^{g-2-i} H^0(C, \omega_C) \otimes H^0(C, \omega_C)\right\},$$

which vanishes since $g-1-i > 0$ and the Koszul differential is an injection. \square

We now let $f : \mathcal{C} \rightarrow \mathcal{M}_g^\sharp$ denote the universal pseudo-stable curve, we let ω_f denote the relative dualizing sheaf, and define the vector bundle \mathcal{M} on \mathcal{M}_g^\sharp by

$$(95) \quad \mathcal{M} := \text{Ker}(f^* f_*(\omega_f) \rightarrow \omega_f),$$

where surjectivity follows because ω_C is globally generated for $[C] \in \mathcal{M}_g^\sharp$. Let $\mathcal{B} := f_*(\bigwedge^i \mathcal{M} \otimes \omega_f^2)$. Using Grauert's theorem [16, Corollary III.12.9] and Proposition 6.4, we conclude that \mathcal{B} is locally free, and by construction, the fiber $\mathcal{B}([C])$ is given as in (94).

To construct \mathcal{A} , we proceed in a way similar to [11]. We begin by observing as in the proof of Proposition 6.4 that $\bigwedge^i M_C(2)$ is resolved by a Koszul type complex

$$\bigwedge^i H^0(\mathbb{P}_C, \mathcal{O}_{\mathbb{P}_C}(1)) \otimes \mathcal{O}_{\mathbb{P}_C}(2) \rightarrow \bigwedge^{i-1} H^0(\mathbb{P}_C, \mathcal{O}_{\mathbb{P}_C}(1)) \otimes \mathcal{O}_{\mathbb{P}_C}(3) \rightarrow \dots$$

By taking global sections and using the fact that $H^0(\mathbb{P}_C, \mathcal{O}_{\mathbb{P}_C}(1)) = H^0(C, \omega_C)$, it follows that

$$H^0\left(\mathbb{P}_C, \bigwedge^i M_C(2)\right) = \text{Ker}\left\{\bigwedge^i H^0(C, \omega_C) \otimes \text{Sym}^2 H^0(C, \omega_C) \rightarrow \bigwedge^{i-1} H^0(C, \omega_C) \otimes \text{Sym}^3 H^0(C, \omega_C)\right\}$$

We consider the locally free sheaf $\mathcal{H} = f_*(\omega_f)$, whose fiber over $[C]$ is $H^0(C, \omega_C)$, and define \mathcal{A} to be the sheaf resolved by the Koszul-type complex

$$\bigwedge^i \mathcal{H} \otimes \text{Sym}^2 \mathcal{H} \rightarrow \bigwedge^{i-1} \mathcal{H} \otimes \text{Sym}^3 \mathcal{H} \rightarrow \dots \rightarrow \text{Sym}^{i+2} \mathcal{H} \rightarrow 0.$$

Since this is a finite locally free complex which is exact except at the beginning, it follows that \mathcal{A} is locally free. Moreover, restricting to the fiber over $[C]$ we get that $\mathcal{A}([C])$ is as in (94).

To construct the map $\alpha : \mathcal{A} \rightarrow \mathcal{B}$, it is enough by adjunction to define a map $f^*\mathcal{A} \rightarrow \bigwedge^i \mathcal{M} \otimes \omega_f^2$. Since f^* commutes with tensor constructions, it follows that $f^*\mathcal{A}$ is resolved by the Koszul-type complex

$$(96) \quad \bigwedge^i (f^*\mathcal{H}) \otimes \mathrm{Sym}^2(f^*\mathcal{H}) \longrightarrow \bigwedge^{i-1} (f^*\mathcal{H}) \otimes \mathrm{Sym}^3(f^*\mathcal{H}) \longrightarrow \cdots \longrightarrow \mathrm{Sym}^{i+2}(f^*\mathcal{H}) \longrightarrow 0.$$

Moreover, it follows from (95) that $\bigwedge^i \mathcal{M} \otimes \omega_f^2$ is resolved by the Koszul-type complex

$$(97) \quad \bigwedge^i (f^*\mathcal{H}) \otimes \omega_f^2 \longrightarrow \bigwedge^{i-1} (f^*\mathcal{H}) \otimes \omega_f^3 \longrightarrow \cdots \longrightarrow \omega_f^{i+2} \longrightarrow 0.$$

The morphism $f^*\mathcal{H} \rightarrow \omega_f$ induces a map between the complexes (96) and (97), which in turn induces a map between the 0-th homology groups, which is the desired map $f^*\mathcal{A} \rightarrow \bigwedge^i \mathcal{M} \otimes \omega_f^2$. Defining α by adjunction and restricting to the fiber over $[C]$ we obtain the natural map α_C in (93), as desired.

We are now in a position to conclude that Theorem 1.1 implies Green's Conjecture for generic curves of genus g in characteristic 0, or characteristic $p \geq \frac{g+2}{2}$.

Proof of Theorem 1.2. Since $\mathrm{char}(\mathbf{k}) = 0$ or $\mathrm{char}(\mathbf{k}) \geq \frac{g+2}{2}$, it follows from Theorem 1.1 that if C is a generic linear section of the tangential variety \mathcal{T} then $K_{[\frac{g}{2}],1}(C, \mathcal{O}_C(1)) = 0$. Such a curve C is an irreducible, g -cuspidal rational curve, where the cusps correspond to the intersection of the generic hyperplane with the rational normal curve. Necessarily, C has maximal gonality $\lfloor \frac{g+3}{2} \rfloor$, therefore C defines a point $[C] \in \mathcal{M}_g^\sharp$. By adjunction, $\omega_C \cong \omega_{\mathcal{T}}(1)|_C \cong \mathcal{O}_C(1)$, by Corollary 5.16. We conclude that C is a canonically embedded pseudo-stable curve, and that $K_{[\frac{g}{2}],1}(C, \omega_C) = 0$. If we take $i = \lfloor \frac{g}{2} \rfloor - 1$ and construct the vector bundles \mathcal{A}, \mathcal{B} and the map $\alpha : \mathcal{A} \rightarrow \mathcal{B}$ as above, it follows that the map α_C is injective, so the same is true for the map $\alpha_{C'}$ for a general point $[C'] \in \mathcal{M}_g^\sharp$. We conclude that the vanishing $K_{[\frac{g}{2}],1}(C', \omega_{C'}) = 0$ holds for a general curve $[C'] \in \mathcal{M}_g$, as desired. \square

Remark 6.5. The proof above shows that every g -cuspidal rational curve satisfies Green's conjecture in suitable characteristics, regardless of the position of the cusps. Indeed, taking any g distinct points on the rational normal curve $\Gamma \subseteq \mathbf{P}^g$ defines a rational g -cuspidal hyperplane section of \mathcal{T} with the same syzygies as \mathcal{T} , since $H^1(\mathcal{T}, \mathcal{O}_{\mathcal{T}}(d)) = 0$ for all $d \geq 0$, see [13, Theorem 3.b.7].

Remark 6.6. Using Corollary 5.16, we observe that the tangent developable $\mathcal{T} \subseteq \mathbb{P}^g$ can be viewed as a degenerate $K3$ surface in the following sense. We denote by \mathcal{F}_g the moduli space of quasi-polarized $K3$ surfaces $[X, H]$, where $H \in \mathrm{Pic}(X)$ is the polarization class with $H^2 = 2g - 2$. Inside \mathcal{F}_g , we consider the unigonal divisor D_g^{ug} consisting of $K3$ surfaces $[X, H]$, such that $H = gE + \Gamma$, where $\Gamma^2 = -2$, $E^2 = 0$ and $\Gamma \cdot E = 0$. The linear system $|H|$ has the rational curve Γ as fixed component and each element of $|H|$ is a flag curve of the form $\Gamma + E_1 + \cdots + E_g$, where $E_i \in |E|$ are elliptic curves. Let \mathcal{H}_g be the Hilbert scheme of quasi-polarized $K3$ surfaces $X \subseteq \mathbb{P}^g$ with $\mathrm{deg}(X) = 2g - 2$. Then $[\mathcal{T}] \in \mathcal{H}_g$. We let $\mathfrak{F}_g := \mathfrak{H}_g // SL(g+1)$ be the corresponding GIT quotient. There is a birational map $\phi : \mathcal{F}_g \dashrightarrow \mathfrak{F}_g$, which blows down the unigonal divisor D_g^{un} to a single point corresponding to the tangential variety $\mathcal{T} \subseteq \mathbb{P}^g$.

REFERENCES

- [1] K. Akin, D. Buchsbaum, J. Weyman, *Schur functors and Schur complexes*, *Advances in Math.* **44** (1982), 207–278.
- [2] M. Aprodu, G. Farkas, Ş. Papadima, C. Raicu, J. Weyman, *Topological invariants of groups and Koszul modules*, arXiv:1806.01702.
- [3] M. Aprodu, J. Nagel, *Koszul cohomology and algebraic geometry*, *AMS University Lecture Series* **52** (2010).
- [4] C. Bopp, F.-O. Schreyer, *A version of Green's Conjecture in positive characteristics*, arXiv:1803.10481.
- [5] I. Coskun, E. Riedl, *Normal bundles of rational curves on complete intersections*, arXiv:1705.08441.

- [6] A. Dimca, S. Papadima, A. Suciu, *Topology and geometry of cohomology jump loci*, Duke Math. J. **148** (2009), 405–457.
- [7] D. Eisenbud, *Green's conjecture: An orientation for algebraists*, in: Free resolutions in commutative algebra and algebraic geometry (Sundance 1990), Res. Notes Math. **2**, Jones and Bartlett, Boston (1992), 51–78.
- [8] D. Eisenbud, *The geometry of syzygies. A second course in commutative algebra and algebraic geometry*. Graduate Texts in Mathematics **229**, Springer-Verlag, New York, 2005.
- [9] D. Eisenbud, J. Harris, *Limit linear series: Basic theory*, Inventiones Math. **85** (1986), 337–371.
- [10] D. Eisenbud, F.-O. Schreyer, *Equations and syzygies of K3 carpets and unions of scrolls*, arXiv:1804.08011.
- [11] G. Farkas, *Koszul divisors on moduli spaces of curves*, American Journal of Math. **131** (2009), 819–869.
- [12] W. Fulton, J. Harris, *Representation theory. A first course*. Graduate Texts in Mathematics **129**, Springer-Verlag, New York (1991).
- [13] M. Green, *Koszul cohomology and the geometry of projective varieties*, J. Differential Geometry **19** (1984), 125–171.
- [14] M. Green, R. Lazarsfeld, *Higher obstructions to deforming cohomology groups of line bundles*, Journal of the American Math. Society **4** (1991), 87–103.
- [15] J. Harris, *Algebraic geometry. A first course*. Springer Graduate Texts in Mathematics, No. **133**, Springer-Verlag (1992).
- [16] R. Hartshorne, *Algebraic geometry*, Springer Graduate Texts in Mathematics, No. **52**, Springer-Verlag (1977).
- [17] R. Hartshorne *Generalized divisors on Gorenstein curves and a theorem of Noether*, J. Math. Kyoto Univ. **26** (1986), 375–386.
- [18] B. Hassett, D. Hyeon, *Log canonical models for the moduli space of curves: The first divisorial contraction*, Transactions of the American Math. Society **361** (2009), 4471–4489.
- [19] C. Hermite, *Sur la theorie des fonctions homogenes à deux indéterminées*, Cambridge and Dublin Math. J. **9** (1854), 172–217.
- [20] J. C. Jantzen, *Representations of algebraic groups*, Pure and Applied Mathematics, No. **131**, Academic Press (1987).
- [21] H. Kaji, *On the tangentially degenerate curves*, J. London Math. Society **33** (1986), 430–440.
- [22] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Math. Monographs (1995).
- [23] S. Papadima, A. Suciu, *Vanishing resonance and representations of Lie algebras*, J. Reine Angew. Math. **706** (2015), 83–101.
- [24] F.-O. Schreyer, *Syzygies of curves with special pencils*, Thesis (Ph.D.)—Brandeis University (1983).
- [25] F.-O. Schreyer, *Syzygies of canonical curves and special linear series*, Mathematische Annalen **275** (1986), 105–137.
- [26] D. Schubert, *A new compactification of the moduli space of curves*, Compositio Math. **78** (1991), 297–313.
- [27] D. Smyth, *Modular compactifications of the space of pointed elliptic curves I*, Compositio Math. **147** (2011), 877–913.
- [28] C. Voisin, *Green's generic syzygy conjecture for curves of even genus lying on a K3 surface*, Journal of the European Math. Society **4** (2002), 363–404.
- [29] C. Voisin, *Green's canonical syzygy conjecture for generic curves of odd genus*, Compositio Math. **141** (2005), 1163–1190.
- [30] J. Wahl, *Gaussian maps on algebraic curves*, Journal of Differential Geometry, **32** (1990), 77–98.
- [31] J. Weyman, *Cohomology of vector bundles and syzygies*, Cambridge Tracts in Mathematics (2003).

MARIAN APRODU: SIMION STOILow INSTITUTE OF MATHEMATICS
 P.O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA &
 FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY OF BUCHAREST
E-mail address: marian.aprodu@imar.ro & marian.aprodu@fmi.unibuc.ro

GAVRIL FARKAS: INSTITUT FÜR MATHEMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN
 UNTER DEN LINDEN 6, 10099 BERLIN, GERMANY
E-mail address: farkas@math.hu-berlin.de

ȘTEFAN PAPADIMA: SIMION STOILow INSTITUTE OF MATHEMATICS
 P.O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA

CLAUDIU RAICU: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME
 255 HURLEY NOTRE DAME, IN 46556, USA, AND
 SIMION STOILow INSTITUTE OF MATHEMATICS,
 P.O. BOX 1-764, RO-014700 BUCHAREST, ROMANIA
E-mail address: craicu@nd.edu

JERZY WEYMAN: DEPARTMENT OF MATHEMATICS UNIVERSITY OF CONNECTICUT
 STORRS, CT, 06269-3009, USA
E-mail address: jerzy.weyman@uconn.edu