

Topology II

Exercise sheet 1

Exercise 1.

Consider the category $\mathcal{A}_{\mathbb{Z}}$ of \mathbb{Z} -graded abelian groups, the category \mathcal{C} of chain complexes and the homotopy category \mathcal{H} , defined as follows:

- Objects of $\mathcal{A}_{\mathbb{Z}}$ are \mathbb{Z} -graded abelian groups, i.e. $G_* = \bigoplus_{n \in \mathbb{Z}} G_n$, where G_n is an abelian group, and morphisms from G_* to H_* are group homomorphisms $\phi: G_* \rightarrow H_*$ satisfying $\phi(G_n) \subset H_n$ for every $n \in \mathbb{Z}$.
- Objects of \mathcal{C} are chain complexes (C_*, ∂) and morphisms from (C_*, ∂_C) to (D_*, ∂_D) are chain maps $\Phi: (C_*, \partial_C) \rightarrow (D_*, \partial_D)$.
- Objects of \mathcal{H} are topological spaces and morphisms from X to Y are homotopy classes of continuous maps $X \rightarrow Y$.

- Verify that these define categories.
- The homology of a chain complex defines a functor H_* from \mathcal{C} to $\mathcal{A}_{\mathbb{Z}}$.
- Let F be some functor from \mathcal{H} to the category of groups \mathcal{G} such that $F(D^n) = 0$ and $F(S^{n-1}) \neq 0$. Conclude from the existence of F the Brouwer fix point theorem.

Exercise 2.

Let (X, x_0) be a pointed topological space. We denote by $\pi_k(X, x_0)$ the homotopy classes of maps $(I^k, \partial I^k) \rightarrow (X, x_0)$.

- Complete the proof of Theorem 2.1 from the lecture, i.e. verify that $f * g$ as defined in the lecture really defines a group structure on $\pi_k(X, x_0)$.

Extra task: Describe the same group structure on the homotopy classes of maps $(S^k, N) \rightarrow (X, x_0)$.

- Prove that $\pi_k(X, x_0)$ is abelian (for $k \geq 2$) by explicitly writing down a homotopy between $f * g$ and $g * f$.
- Let γ be a path from x_0 to x_1 in X . Verify that the map $\gamma_{\#}$ as defined in the lecture induces an isomorphism from $\pi_k(X, x_0)$ to $\pi_k(X, x_1)$.
- The homotopy groups π_k define a covariant functor from the category \mathcal{H} of homotopy classes of pointed topological spaces to the category \mathcal{G} of groups.
- A map $f: (S^k, N) \rightarrow (X, x_0)$ extends to a map $F: D^{k+1} \rightarrow X$ if and only if $[f] = 0$ in $\pi_k(X, x_0)$.

Extra task: Is there a similar statement for relative homotopy groups?